

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	An Open-Label, Multicenter Phase 2 Basket Study to Evaluate the Antitumor Activity and Safety of Lenvatinib in Children, Adolescents, and Young Adults With Relapsed or Refractory Solid Malignancies
Short Title:	A Study of Lenvatinib (MK-7902) in Pediatric Participants With Relapsed or Refractory Solid Malignancies (MK-7902-013/E7080)
Protocol Number:	965908998823247028
NCT Number:	NCT04447755
Sponsor Name:	Merck
Investigational Product:	lenvatinib
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Objective Response Rate (ORR) At Week 16 Per Response Evaluation Criteria In Solid Tumors Version 1.1 (RECIST 1.1) or Response Assessment in Neuro-Oncology (RANO) Criteria (for High Grade Glioma [HGG] Only), by Investigator Assessment: ORR at Week 16 was defined as the percentage of participants with a confirmed complete response (CR: disappearance of all target lesions) or partial response (PR: at least a 30% decrease in the sum of diameters \[SOD\] of target lesions, taking as reference the baseline SOD) as assessed by the investigator per RECIST 1.1 at 16 Weeks. For participants with HGG, response was assessed according to RANO criteria whereby overall response is based on both radiographic response (CR: disappearance of all target lesions, PR: sum of products of diameters \[SPD\] decreased by ? 50% from baseline value) and clinical performance status with steroid dose information.
Secondary Obj.:	1. ORR Per RECIST 1.1 or RANO Criteria (for HGG Only), by Investigator Assessment 2. Progression Free Survival (PFS) Per RECIST 1.1 or RANO (for HGG Only), by Investigator Assessment 3. Best Overall Response (BOR) Per RECIST 1.1 or RANO (for HGG Only), by Investigator Assessment

2.2 Background & Rationale

The main purpose of this study is to evaluate the antitumor activity and safety of lenvatinib (MK-7902/E7080) in children, adolescents, and young adults with relapsed or refractory solid malignancies after administration. Participants were enrolled into Ewing sarcoma (EWS), rhabdomyosarcoma (RMS), high-grade glioma (HGG), diffuse midline glioma, medulloblastoma, ependymoma, and Other Solid Tumors Excluding Osteosarcoma, diffuse midline glioma, medulloblastoma, and ependymoma cohorts.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, Status: COMPLETED

Target N: 127

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Has histologically or cytologically documented relapsed, or refractory pediatric solid malignancy excluding osteosarcoma
- * Has measurable disease as defined by Response Evaluation Criteria In Solid Tumors version 1.1 (RECIST 1.1) or Response Assessment in Neuro-Oncology (RANO) for High Grade Glioma (HGG)
- * Has a performance status defined as follows: 1) Lansky Play Score ≥ 50 for participants up to and including 16 years of age 2) Karnofsky performance status (KPS) ≥ 50 for participants >16 years of age 3) Neurologic deficits in participants with primary central nervous system (CNS) tumors must have been stable for at least 7 days prior to study enrollment
- * Demonstrate adequate organ function
- * No clinical evidence of nephrotic syndrome.
- * Has adequate blood pressure (BP) control with or without antihypertensive medications
- * Has adequate cardiac function
- * Has adequate neurologic function
- * Participant must have fully recovered to Common Terminology Criteria for Adverse Events, Version 5.0 (CTCAE v5.0) Grade ≤ 1 (except for alopecia, ototoxicity, and Grade ≤ 2 peripheral neuropathy) from the acute toxic effects of all prior anticancer therapy
- * Male participants must agree to use approved contraception during the treatment period and for at least 7 days after the last dose of study intervention and refrain from donating sperm during this period
- * Female participants are not pregnant and not breastfeeding, and are not a woman of childbearing potential (WOCBP) or are a WOCBP who agrees to follow contraceptive guidance during the treatment period and for at least 30 days after the last dose of study intervention

4.2 Exclusion Criteria

- * Has had major surgery within 3 weeks prior to Cycle 1 Day 1 (C1D1)
- * Has gastrointestinal (GI) bleeding or active hemoptysis (bright red blood of at least half teaspoon) within 21 days prior to enrollment
- * Has CNS tumors with a history of symptomatic tumor hemorrhage
- * Has evidence of new intracranial hemorrhage of more than punctate size on MRI assessment obtained within 28 days prior to study enrollment
- * Has radiographic evidence of encasement or invasion of a major blood vessel or of intratumoral cavitation
- * Has evidence of untreated CNS metastases (exception: participants with primary CNS tumors and leptomeningeal disease).
- * Has GI malabsorption, GI anastomosis, or any other condition that in the opinion of the investigator might affect the absorption of lenvatinib
- * Has preexisting $\geq$ Grade 3 GI or non-GI fistula

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Has any active infection requiring systemic therapy
- * Known to be Human immunodeficiency virus (HIV) positive
- * Known active viral hepatitis (B or C) as demonstrated by positive serology. Testing for hepatitis B or hepatitis C is required at screening only when mandated by local health authority
- * Is currently participating and receiving study therapy, or has participated in a study of an investigational agent and received study therapy or used an investigational device within 4 weeks of the date of allocation
- * Has a history or current evidence of any condition, therapy, or laboratory abnormality that might confound the results of the study, interfere with the participant's participation for the full duration of the study, or is not in the best interest of the participant to participate, in the opinion of the treating investigator
- * Has known hypersensitivity to any component of the investigational product (lenvatinib or ingredients)
- * Has known psychiatric or substance abuse disorders that would interfere with cooperation with the requirements of the study
- * Has clinically significant cardiovascular disease within 6 months from first dose of study intervention, including New York Heart Association Class III or IV congestive heart failure, unstable angina, myocardial infarction, cerebral vascular accident, or cardiac arrhythmia associated with hemodynamic instability
- * Has non-healing wound, tumor ulceration, unhealed or incompletely healed fracture, or a compound (open) bone fracture at the time of enrollment

11. Study Identifiers & Governance

Primary ID: 965908998823247028

Registry Number: NCT04447755

Sponsor: Merck

Study Title: A Study of Lenvatinib (MK-7902) in Pediatric Participants With Relapsed or Refractory Solid Malignancies (MK-7902-013/E7080)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____