

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Multicentre, Randomised, Double-blind, Active-Controlled, 2-arm Parallel-group Treatment, Phase II Study to Evaluate the Efficacy, Safety, and Tolerability of Zibotentan/Dapagliflozin Compared to Dapagliflozin Alone in Adult Participants With Chronic Kidney Disease and High Proteinuria
Short Title:	A Study to Investigate the Effects of Zibotentan/Dapagliflozin Combination Compared to Dapagliflozin Alone in Adult Participants With Chronic Kidney Disease and High Proteinuria
Protocol Number:	101047386724595813
NCT Number:	NCT06942910
Sponsor Name:	AstraZeneca
Investigational Product:	dapagliflozin
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Change in log-transformed Urinary Albumin to Creatinine Ratio (UACR) from baseline: To estimate the efficacy of zibotentan and dapagliflozin in FDC versus dapagliflozin alone in reducing albuminuria
Secondary Obj.:	1. Change in log-transformed Urinary Protein to Creatinine Ratio (UPCR) from baseline 2. Change in systolic and diastolic blood pressure (BP) from baseline 3. Number of participants experiencing adverse events

2.2 Background & Rationale

A Study to Investigate the Effects of Zibotentan/Dapagliflozin Combination Compared to Dapagliflozin Alone in Adult Participants with Chronic Kidney Disease and High Proteinuria (ZODIAC)

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: RECRUITING
Target N:	224

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * ? 18 years of age at the time of signing the informed consent.
- * Diagnosis of CKD with eGFR ? 20 and \< 90 mL/min/1.73m² AND UACR \> 700 mg/g (\> 79 mg/mmol) or UPCR \> 1000 mg/g (\> 113 mg/mmol).

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Body mass index (BMI) within the range ≥ 40 kg/m².
- * Female participants must be either - not of child-bearing potential or - women of childbearing potential (WOCBP) using at least one highly effective birth control method for at least 3 months prior to first dose of study intervention.
- * All WOCBP must have a negative serum pregnancy test result at screening.
- * Receiving RAASi therapy (ACEi or ARB), and for the patient maximum tolerated labelled daily dose, that has been stable for at least 4 weeks.

4.2 Exclusion Criteria

- * Clinically significant, unstable, or uncontrolled medical condition which in the Investigator's opinion makes it undesirable for the participant to participate in the study.
- * Known hypersensitivity to dapagliflozin or zibotentan or any of the excipients of the investigational product. History or ongoing allergy/hypersensitivity, as judged by the investigator, to SGLT2i therapy or ERAs.
- * NYHA class III or class IV HF.
- * Participants hospitalised for HF and/or who have not been stable on HF therapy during the last 6 months prior to screening.
- * HF due to cardiomyopathies that would primarily require other specific treatment.
- * High output HF (eg, due to hyperthyroidism or Paget's disease).
- * HF due to primary cardiac valvular disease/dysfunction, severe functional mitral or tricuspid valve insufficiency, or planned cardiac valve repair/replacement.
- * Evidence of rales or jugular venous distention on physical examination.
- * Type 1 diabetes mellitus.
- * History of any life-threatening ventricular dysrhythmia (continuous or paroxysmal).
- * Participants hospitalised for heart disease or cardiac procedures or for COVID-19 during the last 3 months prior to screening.
- * History of solid organ transplantation or bone marrow transplant.
- * Any condition with a life expectancy of less than 1 year based on investigator's clinical judgment.
- * Malignancy within the past 5 years. Exceptions to this criterion include non-melanoma skin cancer and curatively treated cervical carcinoma in situ.
- * Significant liver disease as judged by the investigator.
- * Renal replacement therapy or previous kidney transplant.
- * Known history of significant drug or alcohol abuse within 12 months of screening.
- * On treatment with strong or moderate CYP3A4 inducer.
- * On systemic immunosuppression therapy other than prespecified stable maintenance therapy.
- * Participants treated or expecting to be treated with tolvaptan (including as part of participation in a clinical trial), any other ERAs, or budesonide (where used to treat IBD or IgAN).
- * Systolic blood pressure above 160 mmHg and/or below 90 mmHg.
- * Significant impairment of liver function defined as AST or ALT ≥ 3 x upper limit of normal (ULN) or Total serum bilirubin ≥ 2 x ULN (an isolated increase in bilirubin in participants with known Gilbert's syndrome is not a reason for exclusion).
- * NT-proBNP ≥ 600 pg/mL (or NT-proBNP ≥ 1200 pg/mL, if associated with atrial fibrillation) measured by local laboratory at screening.
- * Any of the following results of echocardiography at screening:

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* left ventricular ejection fraction (LVEF) $\leq$ 50%

* significant ventricular wall motion abnormality or severe cardiac valve abnormalities

* isolated pulmonary arterial hypertension (as defined by local clinical practice) or right ventricular failure; in the absence of left-sided HF

* Women who are pregnant, breast-feeding, or women with intent of getting pregnant.

* Women who are not willing to use adequate contraception or cannot, in the opinion of the Investigator, understand and/or comply with the study requirements regarding contraception.

11. Study Identifiers & Governance

Primary ID: 101047386724595813

Registry Number: NCT06942910

Sponsor: AstraZeneca

Study Title: A Study to Investigate the Effects of Zibotentan/Dapagliflozin Combination Compared to Dapagliflozin Alone in Adult Participants With Chronic Kidney Disease and High Proteinuria

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____