

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	An Open-label, Multi-center, Phase I/II Study of ECI830 as a Single Agent and in Combination With Ribociclib and Endocrine Therapy in Patients With Advanced Hormone Receptor Positive, HER2-negative Breast Cancer and Advanced Solid Tumors
Short Title:	Study of ECI830 Single Agent or in Combination in Patients With Advanced HR+/HER2-Breast Cancer and Other Advanced Solid Tumors
Protocol Number:	112657688834418496
NCT Number:	NCT06726148
Sponsor Name:	Novartis
Investigational Product:	fulvestrant
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Phase I: Incidence of dose-limiting toxicities (DLTs): A DLT is defined as an adverse event or abnormal laboratory value of Common Terminology Criteria for Adverse Events (CTCAE) grade $\geq$ 3 assessed as unrelated to disease, disease progression, inter-current illness or concomitant medications that occurs within the first 28 days of treatment. Other clinically significant toxicities may be considered to be DLTs, even if not CTCAE grade 3 or higher. 2. Phase I: Incidence of adverse events (AEs) and serious adverse events (SAEs): Number of participants with AEs and SAEs by treatment group, including changes in vital signs, electrocardiograms (ECGs) and laboratory results qualifying and reported as AEs. 3. Phase I: Number of participants with dose interruptions, reductions and discontinuations: Assessment of tolerability. For patients who do not tolerate the protocol-specified dosing schedule, dose adjustments are permitted in order to allow patients to continue the study treatment.
Secondary Obj.:	1. Phase I and II: Area under the plasma concentration-time curve (AUC) of ECI830 and ribociclib 2. Phase I and II: Maximum observed plasma concentration (Cmax) of ECI830 and ribociclib 3. Phase I and II: Best overall response (BOR) per RECIST v1.1

2.2 Background & Rationale

Phase I: Characterize safety and tolerability of ECI830 as a single agent and in combination with ribociclib and fulvestrant. Identify dose range for optimization/recommended dose for future studies.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: RECRUITING
Target N:	280

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Age ≥ 18 years old.

Patients with one of the following indications:

Phase I:

HR+/HER2- aBC with disease progression on or following at least one line of hormone-based therapy in combination with a CDK4/6i and at least one additional line of systemic therapy for metastatic disease.

Histologically and/or cytologically confirmed diagnosis of locally advanced or metastatic cancer with a CCNE1 amplification. For dose expansion only: no more than 3 prior lines of therapy for advanced or metastatic disease.

Phase II:

HR+/HER2- aBC with disease progression on an aromatase inhibitor or tamoxifen in combination with a CDK4/6 inhibitor for unresectable/metastatic disease with no more than 2 lines of endocrine therapy.

Measurable disease as determined by RECIST v1.1.

BC only: If no measurable disease is present, then at least one predominantly lytic bone lesion must be present that can be accurately assessed at baseline and is suitable for repeated assessment.

4.2 Exclusion Criteria

Previous treatment with a CDK2 inhibitor at any time.

Patients with inadequate bone marrow and/or organ functions with out-of-range laboratory values.

Clinically significant, uncontrolled heart disease and/or cardiac repolarization abnormality including MI, CABG, long QT syndrome, or risk factors for TdP.

Presence of symptomatic CNS metastases or CNS metastases that require local therapy or increasing doses of corticosteroids within 2 weeks prior to study entry.

For the combination treatment:

Patients with symptomatic visceral disease or any disease burden that makes the patient ineligible for endocrine-based therapy.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Patients who could not tolerate the prescribed dose of ribociclib during a previous course of treatment, requiring dose reduction or permanent discontinuation due to adverse events.

For patients with BC: Patient is concurrently using hormone replacement therapy.

WOCBP who are unwilling to use highly effective contraception methods, pregnant or nursing women.

Other protocol-defined inclusion/exclusion criteria may apply.

11. Study Identifiers & Governance

Primary ID:	112657688834418496
Registry Number:	NCT06726148
Sponsor:	Novartis
Study Title:	Study of ECI830 Single Agent or in Combination in Patients With Advanced HR+/HER2- Breast Cancer and Other Advanced Solid Tumors

15. Sign-off & Approval

Principal Investigator: _____	Date: _____
Clinical Operations Lead: _____	Date: _____
Lead Statistician: _____	Date: _____