

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1/2 Open-Label Rolling-Arm Umbrella Platform Design of Investigational Agents With or Without Pembrolizumab or Pembrolizumab Alone in Participants With Melanoma (KEYMAKER-U02): Substudy 02C
Short Title:	Substudy 02C: Safety and Efficacy of Pembrolizumab in Combination With Investigational Agents or Pembrolizumab Alone in Participants With Stage III Melanoma Who Are Candidates for Neoadjuvant Therapy (MK-3475-02C/KEYMAKER-U02)
Protocol Number:	1129920083623432699
NCT Number:	NCT04303169
Sponsor Name:	Merck
Investigational Product:	tretinoin
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of participants who experience an adverse event (AE): An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. The percentage of participants who experience an AE will be reported. 2. Percentage of participants who discontinue study treatment due to an AE: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. The percentage of participants who discontinue study treatment due to an AE will be reported. 3. Pathological complete response (pCR) rate: pCR rate is defined as the proportion of participants with complete absence of viable tumor in the treated tumor bed. Assessments are according to Response Evaluation Criteria in Solid Tumors 1.1 (RECIST 1.1) by central review of the pathology results. RECIST 1.1 has been modified for this study to include a maximum of 10 target lesions and a maximum of 5 target lesions per organ.
Secondary Obj.:	1. Near pathological complete response (near pCR) rate 2. Pathological partial response (pPR) rate 3. Recurrence-free survival (RFS)

2.2 Background & Rationale

Substudy 02C is part of a larger research study that is testing experimental treatments for melanoma, a type of skin cancer. The larger study is the umbrella study.

3. Study Design & Methodology

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: COMPLETED
Target N: 146

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Has histologically or cytologically confirmed melanoma
- * Has clinically detectable and resectable Stage IIIB or IIIC or IIID melanoma amenable to surgery
- * Has been untreated for Stage IIIB, IIIC or IIID melanoma
 - * surgical resection of primary melanoma is allowed
 - * prior radiotherapy to the primary melanoma is allowed
- * Has provided a baseline tumor biopsy
- * Male participants who receive gebasaxturev are abstinent from heterosexual intercourse or agree to use contraception during the intervention period and for at least 120 days after the last dose of gebasaxturev
- * Male participants who receive ATRA are abstinent from heterosexual intercourse or agree to use contraception during the intervention period and for at least 7 days after the last dose of ATRA
- * Female participants are not pregnant or breastfeeding and are either not a woman of child-bearing potential (WOCBP) OR use a contraceptive method that is highly effective or are abstinent from heterosexual intercourse during the intervention period and for at least 120 days after the last dose of pembrolizumab, vibostolimab, gebasaxturev, or MK-4830, favezelimab + pembrolizumab, or 30 days after the last dose of ATRA, whichever occurs last
- * Has adequate organ function
- * Has resolution of toxic effect(s) of the most recent prior therapy to Grade 1 or less (except alopecia)

4.2 Exclusion Criteria

- * Has a diagnosis of immunodeficiency or is receiving immunosuppressive therapy within 7 days before the first dose of study intervention
- * Has a known additional malignancy that is progressing or requires active treatment within the past 2 years
- * Has known central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Has ocular or mucosal melanoma
- * Has known hypersensitivity including previous clinically significant hypersensitivity reaction to treatment with another monoclonal antibody (mAb)
- * Has an active autoimmune disease that has required systemic treatment in the past 2 years
- * Has an active infection requiring systemic therapy
- * Has known history of human immunodeficiency virus (HIV)
- * Has known history of hepatitis B
- * Has a history of (noninfectious) pneumonitis
- * Has a history of active tuberculosis (TB)
- * Has received prior systemic anticancer therapy within 4 weeks prior to randomization
- * Has received prior radiotherapy within 2 weeks of first dose of study intervention
- * Has had major surgery <3 weeks prior to first dose of study intervention

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Has received a live or live-attenuated vaccine within 30 days before the first dose of study intervention
- * Has participated in a study of an investigational agent within 4 weeks prior to the first dose of study intervention
- * Has had an allogeneic tissue/solid organ transplant
- * Has only mucosal lesions
- * Is not naïve to Talimogene laherparepvec (TVEC) and other oncolytic viruses

11. Study Identifiers & Governance

Primary ID: 1129920083623432699

Registry Number: NCT04303169

Sponsor: Merck

Study Title: Substudy 02C: Safety and Efficacy of Pembrolizumab in Combination With Investigational Agents or Pembrolizumab Alone in Participants With Stage III Melanoma Who Are Candidates for Neoadjuvant Therapy (MK-3475-02C/KEYMAKER-U02)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____