

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1/2 Open-Label, Umbrella Platform Design Study of Investigational Agents in Combination With Pembrolizumab (MK-3475) With or Without Chemotherapy in Participants With 1L Locally Advanced Unresectable/Metastatic Esophageal Cancer: KEYMAKER-U06 Substudy 06E
Short Title:	Substudy 06E: Umbrella Study of Combination Therapies in Esophageal Cancer (MK-3475-06E/KEYMAKER-U06)
Protocol Number:	440358557860943556
NCT Number:	NCT06780111
Sponsor Name:	Merck
Investigational Product:	fluorouracil
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of Participants who Experience Dose Limiting Toxicities (DLTs) During the Safety Lead-In Phase: A DLT is any of the following toxicities, if determined to be related to study treatment: * Nonhematologic adverse event (AE) ? Grade 3 (non laboratory) in severity, with exceptions * Grade ? 2 treatment-related interstitial lung disease * Grade 3 or 4 lab value, with exceptions * Potential drug-induced liver injury event, meeting criteria * Febrile neutropenia Grade 3 or 4 * Prolonged delay in initiating Cycle 2 due to study intervention-related toxicity * Study intervention-related toxicity causing participant to discontinue intervention during Cycle 1 * Missing \>25% of study intervention doses due to drug-related AEs during the first cycle * Grade 5 toxicity The number of participants who experience at least one DLT during the safety lead-in phase (approximately 21 days after the first dose of treatment) will be reported. 2. Percentage of Participants who Experience an Adverse Event (AE): An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. The percentage of participants who experience an AE will be reported. 3. Percentage of Participants Who Discontinue Study Intervention Due to an AE: An AE is
--------------------	---

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. The percentage of participants who discontinue study intervention due to an AE will be reported.

Secondary Obj.:

1. Duration of Response (DOR) 2. Progression-Free Survival (PFS) 3. Overall Survival (OS)

2.2 Background & Rationale

Researchers are looking for new ways to treat esophageal squamous cell carcinoma (ESCC). ESCC is a type of cancer that starts in certain cells that line the esophagus. The esophagus is the tube that connects the throat to the stomach. This study will look at ESCC that is either locally advanced unresectable, which means it has spread into tissue near where it started and cannot be completely removed by surgery, or metastatic, which means it has spread to other body parts.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: RECRUITING

Target N: 228

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Not Provided

4.2 Exclusion Criteria

Not Provided

11. Study Identifiers & Governance

Primary ID: 440358557860943556

Registry Number: NCT06780111

Sponsor: Merck

Study Title: Substudy 06E: Umbrella Study of Combination Therapies in Esophageal Cancer (MK-3475-06E/KEYMAKER-U06)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____