

# Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

## 1. Administrative & Study Identification

|                          |                                                                                                                                                                                                                                                  |
|--------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Full Study Title:        | A Phase 1/2 Randomized, Umbrella Study to Evaluate the Efficacy and Safety of MK-2870 Plus Enfortumab Vedotin (EV) With and Without Pembrolizumab, as Treatment for Participants With Advanced Urothelial Carcinoma (KEYMAKER-U04): Substudy 04C |
| Short Title:             | A Study of Efficacy and Safety of Sacituzumab Tirumotecan (MK-2870) Plus Enfortumab Vedotin (EV) With and Without Pembrolizumab in Advanced Urothelial Carcinoma (MK-3475-04C/KEYMAKER-U04)                                                      |
| Protocol Number:         | 516143286643921320                                                                                                                                                                                                                               |
| NCT Number:              | NCT06483334                                                                                                                                                                                                                                      |
| Sponsor Name:            | Merck                                                                                                                                                                                                                                            |
| Investigational Product: | pembrolizumab                                                                                                                                                                                                                                    |
| Phase:                   | PHASE1, PHASE2                                                                                                                                                                                                                                   |
| Medical Monitor:         | [Not Provided in Posting]                                                                                                                                                                                                                        |

## 2. Study Synopsis & Rationale

### 2.1 Study Objective

|                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|--------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Primary Objective: | 1. Part 1: Percentage of Participants with Dose-limiting toxicities (DLT): A DLT is defined as an event with toxicity including the type, severity, time of onset, time of resolution, and the probable association with study treatment that are not due to pre-existing conditions as defined by the National Cancer Institute (NCI) Common Terminology Criteria for Adverse Events, Version 5.0 (CTCAE 5.0). The number of participants who experience a DLT in Part 1 will be reported. 2. Part 1: Percentage of Participants Who Experienced At Least One Adverse Event (AE): An AE is defined as any untoward medical occurrence in a participant administered study drug, which does not necessarily have to have a causal relationship with the study drug. The number of participants experiencing an AE in Part 1 will be reported. 3. Part 1: Percentage of Participants Who Discontinued Study Treatment Due to an AE: An AE is defined as any untoward medical occurrence in a participant administered study drug, which does not necessarily have to have a causal relationship with the study drug. The number of participants who discontinue study treatment due to an AE in Part 1 will be reported. |
| Secondary Obj.:    | 1. Part 1: ORR 2. Part 2: Duration of Response (DOR) 3. Part 1: Maximum Serum Concentration (Cmax) of Sacituzumab Tirumotecan-Antibody-Drug Conjugate (ADC)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |

### 2.2 Background & Rationale

This study is a substudy being conducted under one pembrolizumab umbrella master study KEYMAKER-U04. The substudy will consist of 2 parts. Part 1 will evaluate the safety and preliminary efficacy of sacituzumab tirumotecan plus enfortumab vedotin (EV). Part 2 will be based on Part 1 results and will evaluate the efficacy, pharmacokinetics, and

# Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

safety of sacituzumab tirumotecan plus EV in combination with pembrolizumab in participants with advanced urothelial carcinoma.

## 3. Study Design & Methodology

**Design Framework:** Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: RECRUITING

**Target N:** 98

## 4. Subject Selection (Eligibility Criteria)

### 4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- \* Must have histologically documented, locally advanced/metastatic urothelial carcinoma (la/mUC).
- \* Must provide an archival tumor tissue sample or newly obtained core or excisional biopsy of a tumor lesion demonstrating UC, not previously irradiated, and adequate for biomarker evaluation. A newly obtained biopsy is strongly preferred, but not required if archival tissue is evaluable.
- \* Any AEs due to previous anticancer therapies must have recovered to ?Grade 1 or baseline. Endocrine-related AEs adequately treated with hormone replacement are eligible.
- \* PART 1 ONLY: Participants must have received platinum-based chemotherapy for treatment of la/mUC.
- \* PART 1 ONLY: Participants must not have received \>2 lines of therapy for la/mUC. Platinum-based chemotherapy followed by avelumab maintenance is considered 2 lines of therapy.
- \* PART 2 ONLY: Participants must not have received prior systemic therapy for la/mUC.

### 4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

- \* Known additional malignancy that is progressing or has required active treatment within the past 3 years.
- \* Known active central nervous system metastases and/or carcinomatous meningitis.
- \* Has Grade ?2 peripheral neuropathy.
- \* Has history of documented severe dry eye syndrome, severe Meibomian gland disease and/or blepharitis, or corneal disease that prevents/delays corneal healing.
- \* Has active inflammatory bowel disease requiring immunosuppressive medication or previous history of inflammatory bowel disease (eg, Crohn's disease, ulcerative colitis, or chronic diarrhea).
- \* Has uncontrolled, significant cardiovascular disease or cerebrovascular disease and/or serious cardiovascular and cerebrovascular diseases within the 6 months preceding study intervention.
- \* Has active keratitis or corneal ulcerations. Superficial punctate keratitis is allowed if the disorder is being adequately treated in the opinion of the investigator.
- \* Has a history of uncontrolled diabetes.
- \* Has received an investigational agent or has used an investigational device within 4 weeks prior to study intervention administration.
- \* Has received a live or live-attenuated vaccine within 30 days prior to the first dose of study intervention.

# Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- \* PART 2 ONLY: Has a diagnosis of immunodeficiency or is receiving chronic systemic steroid therapy or any other form of immunosuppressive therapy within 7 days before the first dose of study intervention. Inhaled or topical steroids are permitted in the absence of active autoimmune disease. Physiologic replacement doses of corticosteroids are permitted for participants with adrenal insufficiency.
- \* PART 2 ONLY: Has an active autoimmune disease that has required systemic treatment in past 2 years except replacement therapy. - Is human immunodeficiency virus (HIV)-infected and has a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease.
- \* Has active Hepatitis B or Hepatitis C virus infection.
- \* Has a history of (noninfectious) pneumonitis/interstitial lung disease that required steroids or has current pneumonitis/interstitial lung disease.
- \* Has an active infection requiring systemic therapy.
- \* PART 2 ONLY: History of allogeneic tissue/solid organ transplant.
- \* Has not adequately recovered from major surgery or has ongoing surgical complications.

## 11. Study Identifiers & Governance

**Primary ID:** 516143286643921320

**Registry Number:** NCT06483334

**Sponsor:** Merck

**Study Title:** A Study of Efficacy and Safety of Sacituzumab Tirumotecan (MK-2870) Plus Enfortumab Vedotin (EV) With and Without Pembrolizumab in Advanced Urothelial Carcinoma (MK-3475-04C/KEYMAKER-U04)

## 15. Sign-off & Approval

Principal Investigator: \_\_\_\_\_ Date: \_\_\_\_\_

Clinical Operations Lead: \_\_\_\_\_ Date: \_\_\_\_\_

Lead Statistician: \_\_\_\_\_ Date: \_\_\_\_\_