

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase II, Open-label, Multicentre, Randomised Study of Neoadjuvant and Adjuvant Treatment in Patients With Resectable, Early-stage (II to IIIB) Non-small Cell Lung Cancer (NeoCOAST-2)
Short Title:	Neoadjuvant and Adjuvant Treatment in Resectable Non-small Cell Lung Cancer
Protocol Number:	521499630002922265
NCT Number:	NCT05061550
Sponsor Name:	AstraZeneca
Investigational Product:	cisplatin
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of participants with pathological complete response (pCR) 2. Number of participants with adverse events (AEs) and serious adverse events (SAEs)
Secondary Obj.:	1. Number of participants experiencing an event-free survival (EFS) event 2. Number of participants experiencing a disease-free survival (DFS) event 3. Number of participants having surgical resection

2.2 Background & Rationale

The study is intended to assess the safety and efficacy of perioperative treatment with Durvalumab in combination with Oleclumab, Monalizumab, or AZD0171 and platinum doublet chemotherapy (CTX); or Volrustomig or Rilvegostomig in combination with CTX; or Datopotamab deruxtecan (Dato-DXd) in combination with Durvalumab or Rilvegostomig and single agent platinum chemotherapy in participants with resectable, early-stage non-small cell lung cancer.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: RECRUITING
Target N:	630

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Newly diagnosed NSCLC patients with resectable disease (Stage IIA to Stage IIIB).
- * WHO or Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1.
- * Adequate organ and bone marrow function.
- * Provision of tumour samples (newly acquired or archival tumour tissue \[? 6 months old\]) to confirm Programmed death-ligand 1 (PD-L1) status, epidermal growth factor receptor (EGFR), or anaplastic lymphoma kinase (ALK) status.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* Adequate pulmonary function.

4.2 Exclusion Criteria

- * Participants with sensitising EGFR mutations or ALK translocations.
- * Participants with baseline PD-L1 expression status $\leq 1\%$ (Arms 6 and 7 only).
- * Active or prior documented autoimmune or inflammatory disorders.
- * Uncontrolled intercurrent illness, uncontrolled hypertension, unstable angina pectoris, uncontrolled cardiac arrhythmia, active bleeding diseases, serious chronic gastrointestinal conditions associated with diarrhoea, or psychiatric illness/social situations that would limit compliance with study requirement.
- * History of another primary malignancy.
- * Participants with small-cell lung cancer or mixed small-cell lung cancer.
- * History of active primary immunodeficiency.
- * History of non-infectious interstitial lung disease (ILD) ILD/pneumonitis that required steroids, has current ILD/pneumonitis, or has suspected ILD/pneumonitis that cannot be ruled out by imaging at screening.
- * Participants who have preoperative radiotherapy treatment as part of their care plan.
- * Participants who require or may require pneumonectomy, segmentectomies, or wedge resections, as assessed by their surgeon at baseline, to obtain potentially curative resection of primary tumour.
- * QTcF (QT interval corrected by Fridericia's formula) interval ≥ 470 ms.
- * Any medical contraindication to treatment with chemotherapy as listed in the local labelling.
- * Participants with moderate or severe cardiovascular disease.
- * Any concurrent chemotherapy, investigational product, biologic, or hormonal therapy for cancer treatment.
- * Receipt of live attenuated vaccine within 30 days prior to the first dose of study interventions.
- * Prior exposure to approved or investigational immune-mediated therapy including, but not limited to, other anti-CTLA-4, anti-TIGIT (T cell immunoreceptor with Ig and ITIM domains), anti-PD-1, anti-PD-L1, and anti-PD-L2 antibodies. Participants who received agents targeting the adenosine pathway, anti-NKG2A, anti-HLA-E agents, and anti-LIF agents are also excluded. Participants who have received previous treatment with a TROP2 targeting ADC or with another ADC containing a chemotherapy agent that inhibits TOP1 activity are also excluded.
- * Current or prior use of immunosuppressive medication within 14 days before the first dose of study interventions.
- * Active or uncontrolled infections including HBA, HBV, HCV, and HIV.

11. Study Identifiers & Governance

Primary ID: 521499630002922265
Registry Number: NCT05061550
Sponsor: AstraZeneca
Study Title: Neoadjuvant and Adjuvant Treatment in Resectable Non-small Cell Lung Cancer

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____