

# Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

## 1. Administrative & Study Identification

|                          |                                                                                                                                                                                                                                              |
|--------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Full Study Title:        | A Phase IIb, Randomised, Multicentre, Double-Blind Study to Evaluate the Effect of Baxdrostat in Combination With Dapagliflozin Compared With Baxdrostat on Albuminuria in Participants With Chronic Kidney Disease and High Blood Pressure. |
| Short Title:             | A Phase IIb Study to Evaluate the Effect of Dapagliflozin in Combination With Baxdrostat Compared With Baxdrostat on Albuminuria in Participants With Chronic Kidney Disease and High Blood Pressure.                                        |
| Protocol Number:         | 568220845953007107                                                                                                                                                                                                                           |
| NCT Number:              | NCT07222917                                                                                                                                                                                                                                  |
| Sponsor Name:            | AstraZeneca                                                                                                                                                                                                                                  |
| Investigational Product: | dapagliflozin                                                                                                                                                                                                                                |
| Phase:                   | PHASE2                                                                                                                                                                                                                                       |
| Medical Monitor:         | [Not Provided in Posting]                                                                                                                                                                                                                    |

## 2. Study Synopsis & Rationale

### 2.1 Study Objective

|                    |                                                                                                                                                    |
|--------------------|----------------------------------------------------------------------------------------------------------------------------------------------------|
| Primary Objective: | 1. To determine whether baxdrostat/dapagliflozin is superior to baxdrostat/matching placebo at reducing albuminuria.: Change from baseline in UACR |
| Secondary Obj.:    | Not Provided                                                                                                                                       |

### 2.2 Background & Rationale

International, Multicenter and Double-Blind study. The purpose is to measure the effect of baxdrostat in combination with dapagliflozin compared with baxdrostat/placebo on albuminuria, as well as safety, in participants with chronic kidney disease and high blood pressure.

## 3. Study Design & Methodology

|                   |                                                                 |
|-------------------|-----------------------------------------------------------------|
| Design Framework: | Type: INTERVENTIONAL, Phase: PHASE2, Status: NOT_YET_RECRUITING |
| Target N:         | 218                                                             |

## 4. Subject Selection (Eligibility Criteria)

### 4.1 Inclusion Criteria

- Participants of any sex and gender must be  $\geq 18$  years of age at the time of signing the informed consent.
- Participants with eGFR  $\geq 30$  and  $< 90$  mL/min/1.73 m<sup>2</sup> at screening
- Participants with UACR  $> 200$  mg/g (22.6 mg/mmol) and  $< 5000$  mg/g (565 mg/mmol) at screening
- Participants with history of HTN and a SBP  $\geq 130$  mmHg at screening and  $\geq 120$  mmHg at the randomisation visit.
- Stable and maximum daily tolerated dose of either an ACE inhibitor or an ARB (not both) for at least 4 weeks prior to the screening visit, if not medically contraindicated.

# Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

## 6. Participants with:

1. Serum or plasma potassium  $\geq 3.0$  and  $\leq 4.8$  mmol/L if eGFR  $\geq 45$  mL/min/1.73 m<sup>2</sup>.
2. Serum or plasma potassium  $\geq 3.0$  and  $\leq 4.5$  mmol/L if eGFR  $< 45$  mL/min/1.73 m<sup>2</sup>.
7. Contraceptive use should be consistent with local regulations regarding the methods of contraception for those participating in clinical studies. Applicable to female participants.

## 4.2 Exclusion Criteria

1. Systolic blood pressure  $> 180$  mmHg, or diastolic blood pressure  $> 110$  mmHg at screening.
2. Known hyperkalaemia, defined as potassium of  $\geq 5.5$  mmol/L within 3 months before screening
3. Serum sodium  $< 135$  mmol/L at the Screening Visit (values obtained within 4 weeks prior to screening or at the Screening Visit).
4. Diabetes mellitus:
  1. T1DM at the screening visit
  2. Uncontrolled T2DM at screening: HbA1C  $> 10.5\%$  ( $> 91$  mmol/mol)
5. New York Heart Association functional HF class IV at screening
6. Any use of mineralocorticoid receptor antagonists (such as spironolactone, eplerenone, or finerenone), aldosterone synthase inhibitors, potassium-sparing diuretics (such as triamterene or amiloride), or potassium binders (such as sodium zirconium cyclosilicate, patiomer, or sodium polystyrene sulfonate) within 4 weeks prior to screening
7. Stroke, transient ischaemic cerebral attack, valve implantation or valve replacement, carotid surgery, or carotid angioplasty, acute coronary syndrome, or hospitalisation for worsening HF within previous 3 months prior to randomisation.
8. Known severe hepatic impairment, defined as Child-Pugh Class C, based on records that confirm documented medical history.
9. Documented history of adrenal insufficiency.
10. Any dialysis (including for acute kidney injury) within 3 months prior to the screening
11. Any acute kidney injury within 3 months prior to the screening visit.
12. Prohibited concomitant medications

## 11. Study Identifiers & Governance

|                  |                                                                                                                                                                                                       |
|------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Primary ID:      | 568220845953007107                                                                                                                                                                                    |
| Registry Number: | NCT07222917                                                                                                                                                                                           |
| Sponsor:         | AstraZeneca                                                                                                                                                                                           |
| Study Title:     | A Phase IIb Study to Evaluate the Effect of Dapagliflozin in Combination With Baxdrostat Compared With Baxdrostat on Albuminuria in Participants With Chronic Kidney Disease and High Blood Pressure. |

## 15. Sign-off & Approval

Principal Investigator: \_\_\_\_\_ Date: \_\_\_\_\_

# Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Clinical Operations Lead: \_\_\_\_\_ Date: \_\_\_\_\_

Lead Statistician: \_\_\_\_\_ Date: \_\_\_\_\_