

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	MK-5684-01A Substudy: A Phase 1/2 Umbrella Substudy of MK-5684-U01 Master Protocol to Evaluate the Safety and Efficacy of MK-5684-based Treatment Combinations or MK-5684 Alone in Participants With Metastatic Castration-resistant Prostate Cancer (mCRPC)
Short Title:	Substudy 01A: Safety and Efficacy of Opevesostat (MK-5684)-Based Treatment Combinations or Opevesostat Alone in Participants With Metastatic Castration-resistant Prostate Cancer (mCRPC) (MK-5684-01A)
Protocol Number:	871279987455803217
NCT Number:	NCT06353386
Sponsor Name:	Merck
Investigational Product:	cortisone
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of participants who experience one or more dose-limiting toxicities (DLTs): The following events, if considered drug related by the investigator, will be considered a DLT: Grade 4 nonhematologic toxicity (not laboratory value); Grade 4 hematologic toxicity lasting ≥ 7 days, except thrombocytopenia (Grade 4 thrombocytopenia of any duration, Grade 3 thrombocytopenia associated with clinically significant bleeding); Any nonhematologic adverse event (AE) $\geq$ Grade 3 in severity should be considered a DLT (with exceptions); Any Grade 3 or Grade 4 nonhematologic laboratory value (if certain criteria are met); Febrile neutropenia Grade 3 or Grade 4; Prolonged delay (≥ 2 weeks) in initiating treatment after the first 28 days due to study intervention-related toxicity; Missing $\geq 25\%$ of study intervention doses as a result of drug-related AE(s) during the first 28 days; Grade 5 toxicity. 2. Number of participants who experience one or more adverse events (AEs): An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. 3. Number of participants who discontinue study intervention due to an AE: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention.
Secondary Obj.:	1. Objective response rate (ORR) 2. Radiographic progression-free survival (rPFS) 3. Overall survival (OS)

2.2 Background & Rationale

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Substudy 01A is part of a larger research study that is testing experimental treatments for metastatic castration-resistant prostate cancer (mCRPC). The larger study is the umbrella study (U01).

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: RECRUITING

Target N: 220

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- * Histologically or cytologically confirmed diagnosis of adenocarcinoma of the prostate without small cell histology.
- * Prostate cancer progression and received androgen deprivation therapy (ADT) or post bilateral orchiectomy within 6 months before screening.
- * Evidence of disease progression from either, ≥ 4 weeks from last flutamide treatment, or ≥ 6 weeks from last bicalutamide or nilutamide treatment, if receiving first generation anti-androgen therapy as last treatment therapy.
- * Current evidence of metastatic disease.
- * Prior treatment with 1 to 2 novel hormonal agent(s) (NHA) for non-metastatic, or metastatic, hormone-sensitive prostate cancer or castration-resistant prostate cancer and have disease progression during or after treatment.
- * Treatment with bone resorptive therapy (including, but not limited to, bisphosphonate or denosumab) must have been on stable doses for ≥ 4 weeks before randomization.
- * Participants who experienced adverse events (AEs) due to previous anticancer therapies must have recovered to $\leq$ Grade 1 or baseline.
- * Human immunodeficiency virus (HIV)-infected participants must have well controlled HIV on antiretroviral therapy.
- * Participants who are Hepatitis B surface antigen (HBsAg) positive are eligible if they have received Hepatitis B Virus (HBV) antiviral therapy for at least 4 weeks, and have undetectable HBV viral load.
- * Participants with a history of Hepatitis C virus (HCV) infection are eligible if HCV viral load is undetectable.

4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

- * History of pituitary dysfunction.
- * Poorly controlled diabetes mellitus.
- * Active or unstable cardio/cerebro-vascular disease, including thromboembolic events and history of stroke or transient ischemic attack within 6 months before the first dose of study intervention, history of myocardial infarction within 6 months before the first dose of study intervention, New York Heart Association Class III or IV cardiac disease or congestive heart failure, coronary heart disease that is symptomatic, or unstable angina
- * History or family history of long corrected QT interval (QTc) syndrome.
- * Myelodysplastic syndrome (MDS)/acute myeloid leukemia (AML) or features suggestive of MDS/AML.
- * History or current condition of adrenal insufficiency.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * History of (noninfectious) pneumonitis requiring steroids, or current pneumonitis.
- * HIV-infected participants with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease.
- * Undergone major surgery, including local prostate intervention (except prostate biopsy) within 28 days before randomization, and has not recovered from the toxicities and/or complications.
- * Is on an unstable dose of thyroid hormone therapy within 6 months prior to first dose of study intervention.
- * Received a whole blood transfusion in the last 120 days before randomization (packed red blood cells and platelet transfusions are acceptable if not given within 28 days before randomization).
- * Received prior systemic anticancer therapy including investigational agents within 4 weeks before randomization.
- * Received prior radiotherapy within 2 weeks of start of study intervention or radiation-related toxicities, requiring corticosteroids.
- * Received a live or live-attenuated vaccine within 30 days before the first dose of study intervention. Administration of killed vaccines is allowed.
- * Diagnosis of immunodeficiency, or is receiving chronic systemic steroid therapy, or any other form of immunosuppressive therapy, within 7 days prior to the first dose of study intervention.
- * Known additional malignancy that is progressing or has required active treatment within the past 3 years.
- * Known active central nervous system (CNS) metastases and/or carcinomatous meningitis.
- * Active autoimmune disease that has required systemic treatment in the past 2 years.
- * Active infection requiring systemic therapy.
- * Concurrent active HBV or HCV infections.

11. Study Identifiers & Governance

Primary ID:	871279987455803217
Registry Number:	NCT06353386
Sponsor:	Merck
Study Title:	Substudy 01A: Safety and Efficacy of Opevesostat (MK-5684)-Based Treatment Combinations or Opevesostat Alone in Participants With Metastatic Castration-resistant Prostate Cancer (mCRPC) (MK-5684-01A)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____