

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase I/II, Open-Label, Multi-Cohort Study to Evaluate the Efficacy and Safety of Cevostamab in Prior B Cell Maturation Antigen-Exposed Patients With Relapsed/Refractory Multiple Myeloma
Short Title:	A Study Evaluating the Efficacy and Safety of Cevostamab in Prior B Cell Maturation Antigen (BCMA)-Exposed Participants With Relapsed/Refractory Multiple Myeloma
Protocol Number:	936350482842535436
NCT Number:	NCT05535244
Sponsor Name:	Roche
Investigational Product:	tocilizumab
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Objective Response Rate (ORR) as Determined by the Investigator 2. Percentage of Participants with Adverse Events
Secondary Obj.:	1. ORR as Determined by the Independent Review Committee (IRC) 2. Duration of Response (DOR) 3. Rate of Complete Response (CR) or Better

2.2 Background & Rationale

This study will evaluate the efficacy, safety, and pharmacokinetics of cevostamab in participants with relapsed or refractory multiple myeloma (R/R MM) via intravenous (IV) infusion.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: ACTIVE_NOT_RECRUITING
Target N:	90

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Documented diagnosis of MM based on standard International Myeloma Working Group (IMWG) criteria
- * Evidence of progressive disease based on investigators determination of response by IMWG criteria on or after their last dosing regimen
- * Prior BCMA ADC or CAR-T Cohort: participants who have received a BCMA-targeted CAR-T or ADC therapy and are triple-class relapsed or refractory
- * Prior BCMA Bispecific Cohort: participants who have received a BCMA-targeting T-cell-dependent bispecific (TDB)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

antibody and are triple-class relapsed or refractory

- * Eastern Cooperative Oncology Group (ECOG) Performance Status of 0 or 1
- * Life expectancy is at least 12 weeks
- * Agreement to protocol-specified assessments, including bone marrow biopsy and aspirate samples as detailed in the protocol
- * Resolution of AEs from prior anti-cancer therapy to Grade $\leq$ 1
- * For female participants of childbearing potential: agreement to remain abstinent (refrain from heterosexual intercourse) or use contraception during the treatment period and for at least 5 months after the final dose of cevostamab and for 3 months after the last dose of tocilizumab was administered
- * For male participants: agreement to remain abstinent (refrain from heterosexual intercourse) or use a condom, and agree to refrain from donating sperm during the treatment period and for at least 2 months after the final dose of tocilizumab (if applicable) to avoid exposing the embryo

4.2 Exclusion Criteria

- * Inability to comply with protocol-mandated hospitalization
- * Pregnancy or breastfeeding, or intention of becoming pregnant during the study or within 5 months after the final dose of cevostamab or tocilizumab or within 3 months after the last dose of tocilizumab (if applicable)
- * Prior treatment with cevostamab or another agent with the same target
- * Prior BCMA ADC or CAR-T Cohort: prior treatment with any T cell dependent bi-specific antibody (TDB) antibody including non BCMA targeting TDB
- * Prior use of any monoclonal antibody (mAb), radioimmunoconjugate, or ADC as anti-cancer therapy within 4 weeks before first study treatment, except for the use of non-myeloma therapy
- * Prior treatment with systemic immunotherapeutic agents
- * Prior treatment with CAR-T cell therapy within 12 weeks before first cevostamab infusion
- * Known treatment-related, immune-mediated adverse events associated with prior checkpoint inhibitors
- * Treatment with radiotherapy, any chemotherapeutic agent, or treatment with any other anti-cancer agent within 4 weeks or 5 half-lives of the drug, whichever is shorter, prior to first study treatment
- * Autologous stem cell transplantation (SCT) within 100 days prior to first study treatment
- * Prior allogeneic SCT
- * Circulating plasma cell count exceeding 500/ microliter (μ L) or 5% of the peripheral blood white cells
- * Prior solid organ transplantation
- * History of autoimmune disease
- * History of confirmed progressive multifocal leukoencephalopathy
- * History of severe allergic or anaphylactic reactions to mAb therapy
- * Known history of amyloidosis
- * Lesions in proximity of vital organs that may develop sudden decompensation/deterioration in the setting of a tumor flare
- * History of other malignancy within 2 years prior to screening, except those with negligible risk of metastasis or death, such as ductal carcinoma in situ not requiring chemotherapy, appropriately treated carcinoma in situ of the cervix, non-melanoma skin carcinoma, low-grade, localized prostate cancer not requiring treatment or appropriately treated Stage I uterine cancer

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Current or past history of central nervous system (CNS) disease, such as stroke, epilepsy, CNS vasculitis, neurodegenerative disease, or CNS involvement by MM
- * Significant cardiovascular disease that may limit a potential participant's ability to adequately respond to a cytokine release syndrome (CRS) event
- * Symptomatic active pulmonary disease or requiring supplemental oxygen
- * Known active bacterial, viral, fungal, mycobacterial, parasitic, or other infection at study enrollment, or any major episode of infection requiring treatment with IV (intravenous) antimicrobials where the last dose of IV antimicrobial was given within 14 days prior to first study treatment
- * Active symptomatic COVID-19 infection at study enrollment or requiring treatment with IV antiviral where the last dose of IV antiviral treatment was given within 14 days prior to first study treatment. Participants with active COVID-19 infection must have clinical recovery and two negative antigen tests at least 24 hours apart prior to first study treatment
- * Positive and quantifiable Epstein-Barr virus (EBV) polymerase chain reaction (PCR) or cytomegalovirus (CMV) PCR prior to first study treatment
- * Known or suspected chronic active EBV infection
- * Known history of Grade ≥ 3 CRS or immune effector cell-associated neurotoxicity syndrome (ICANS) with prior bispecific therapies
- * Known history of hemophagocytic lymphohistiocytosis (HLH) or macrophage activation syndrome (MAS)
- * Recent major surgery within 4 weeks prior to first study treatment
- * Positive serologic or PCR test results for acute or chronic hepatitis B virus (HBV) infection
- * Acute or chronic hepatitis C virus (HCV) infection
- * Known history of human immunodeficiency virus (HIV) seropositivity
- * Administration of a live, attenuated vaccine within 4 weeks before first study treatment or anticipation that such a live attenuated vaccine will be required during the study
- * Treatment with systemic immunosuppressive medications, with the exception of corticosteroid treatment ≤ 10 mg/day prednisone or equivalent, within 2 weeks prior to first study treatment
- * History of illicit drug or alcohol abuse within 12 months prior to screening, in the investigator's judgment
- * Any medical condition or abnormality in clinical laboratory tests that, in the investigator's judgment, precludes the participant's safe participation in and completion of the study, or which could affect compliance with the protocol or interpretation of results

11. Study Identifiers & Governance

Primary ID:	936350482842535436
Registry Number:	NCT05535244
Sponsor:	Roche
Study Title:	A Study Evaluating the Efficacy and Safety of Cevostamab in Prior B Cell Maturation Antigen (BCMA)-Exposed Participants With Relapsed/Refractory Multiple Myeloma

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____