

1. Administrative & Study Identification

Full Study Title: A Phase 2, Open-label, Randomized Study of MORAb-202 (Farletuzumab Ecteribulin), a Folate Receptor Alpha-targeting Antibody-Drug Conjugate, in Participants With Metastatic Non-Small Cell Lung Cancer (NSCLC) Adenocarcinoma (AC) After Progression on Prior Therapies **Short Title/Acronym:** A Study of MORAb-202 in Participants With Previously Treated Metastatic NSCLC AC **Protocol Number:** CA116-003 **NCT Number:** NCT05577715 **Sponsor Name:** Bristol-Myers Squibb **Investigational Product:** MORAb-202 (Farletuzumab Ecteribulin / BMS-986445) **Phase of Development:** Phase 2 **Medical Monitor:** Bristol-Myers Squibb Clinical Operations

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To characterize the safety and tolerability of MORAb-202, and to assess the objective response rate (ORR) in participants with previously treated, metastatic NSCLC Adenocarcinoma (AC). **Secondary Objectives:** To evaluate Progression-Free Survival (PFS), Duration of Response (DOR), and overall survival in treated participants.

2.2 Background & Rationale Patients with NSCLC adenocarcinoma who have experienced disease progression after prior approved therapies with platinum-doublet chemotherapy and immune checkpoint inhibitors, or targeted therapies for actionable mutations, face a high unmet medical need. Effective treatment options after progression on standard-of-care therapies are urgently needed. MORAb-202 is a novel folate receptor alpha (FR α)-targeting antibody-drug conjugate (ADC) designed to deliver an eribulin payload to FR α -expressing tumor cells, causing cell death by apoptosis.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Active evaluation (Single Group Assignment / Dose Evaluation) **Blinding:** Open-label **Randomization:** Randomized

3.2 Planned Enrollment & Duration Target \$N\$: 31 participants (Actual Enrollment) **Number of Sites:** 33 locations globally **Estimated Study Start:** 14-Nov-2022 (Actual) **Estimated Primary Completion:** 12-Aug-2024 (Actual)

4. Subject Selection (Eligibility Criteria)

- 4.1 Inclusion Criteria 1. Histologically or cytologically documented metastatic NSCLC AC (as defined by the 8th International Association for the Study of Lung Cancer Classification). 2. Measurable target disease assessed by the investigator according to RECIST 1.1. 3. Eastern Cooperative Oncology Group (ECOG) performance status (PS) of 0 or 1.
- 4.2 Exclusion Criteria 1. NSCLC histologies other than Adenocarcinoma (i.e., squamous cell carcinoma, large cell carcinoma). 2. Significant third-space fluid retention (e.g., ascites or pleural effusion) that requires repeated drainage. 3. Prior pneumonectomy. Prior lobectomy and segmentectomy are allowed if >12 months before treatment. Recent chest radiotherapy is excluded unless documented >6 months before starting study treatment.

5. Intervention & Treatment Plan

- 5.1 Investigational Regimen Dosage/Frequency: MORAb-202 administered every 3 weeks (21-day cycles). Cycle 1 includes additional visits on Days 8 and 15. Route of Administration: Intravenous infusion Duration of Treatment: Cycles repeat until documented disease progression, unacceptable toxicity, or study withdrawal.
- 5.2 Concomitant & Prohibited Therapy Permitted: Standard supportive care for advanced cancer symptoms as per institutional guidelines. Prohibited: Concurrent investigational agents, live vaccines, and other systemic anti-cancer therapies during the active treatment phase.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Screening	Day -14 to 0 Informed Consent, Medical History, Vital signs, Baseline Scans
Baseline	Cycle 1 Day 1	Randomization, First Dose, Safety Labs	Interim
Cycle 1 (Days 8, 15);	Subsequent Cycles (Day 1)	IV Infusion, AE Monitoring, Efficacy Scans (every ~6-8 weeks)	End of Study
30 Days Post-Last Dose	Final Safety Evaluation, Survival Follow-up up to 12 months		

7. Outcome Measures (Endpoints)

- 7.1 Primary Endpoint Definition: Objective Response Rate (ORR), defined as the proportion of participants achieving a Complete Response (CR) or Partial Response (PR). Measurement Method: Assessed by the investigator per RECIST 1.1 criteria.

7.2 Secondary & Exploratory Endpoints Progression-Free Survival (PFS), evaluated as the time from randomization to the date of documentation of disease progression or death due to any cause. Overall Incidence of Adverse Events (AEs) and Serious Adverse Events (SAEs).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: AEs will be monitored from the first dose and for 30 days after the last dose of study therapy, with survival/long-term follow-up extending to approximately 12 months. **Serious Adverse Events (SAEs):** Continuously tracked and reported per regulatory timelines. **Data Monitoring Committee (DMC):** A Data Monitoring Committee will not be used in this specific study.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: All randomized participants who receive at least one dose of the investigational product. **Sample Size Justification:** Enrolled target of 31 participants provided sufficient power to assess preliminary safety, tolerability, and objective response signals in a Phase 2 setting. **Handling of Missing Data:** Standard oncological censoring logic applies (e.g., participants lost to follow-up censored at the last evaluable tumor assessment for PFS).

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Standard clinical site monitoring and data verification conducted by Bristol-Myers Squibb and contracted CROs. **Audit Trail:** eCRF entry, laboratory data reconciliation, and investigator sign-offs on Response Criteria assessments.

11. Study Identifiers & Governance

Primary ID: CA116-003 **Registry Number:** NCT05577715 **Sponsor/Lead Organization:** Bristol-Myers Squibb **Study Title:** A Study of MORAb-202 in Participants With Previously Treated Metastatic Non-Small Cell Lung Cancer (NSCLC) Adenocarcinoma (AC) **Official Regulatory Title:** A Phase 2, Open-label, Randomized Study of MORAb-202 (Farletuzumab Ecteribulin), a Folate Receptor Alpha-targeting Antibody-Drug Conjugate, in Participants With Metastatic Non-Small Cell Lung Cancer (NSCLC) Adenocarcinoma (AC) After Progression on Prior Therapies

12. Clinical Framework (NSCLC Specific)

Primary Condition: Metastatic Non-Small Cell Lung Cancer (NSCLC) Adenocarcinoma **Diagnosis Threshold:** Progression on prior therapies (platinum-doublet chemotherapy, immune checkpoint inhibitors, or targeted therapies) **Medical Methodology:** Targeted Antibody-Drug Conjugate (ADC) therapy **Optimization Target:** Objective tumor response and disease control

13. Study Procedures & Methodology

Management Pathway: Standard oncology clinical pathway for advanced NSCLC progressing on first-line therapies. **Education Protocol (HCP):** Clinical site initiation, handling/infusion procedures for the ADC payload, and AE monitoring. **Education Protocol (Patient):** Informed consent, treatment expectations, and AE self-reporting guidance. **Quality Audit Mechanism:** Independent medical/radiological review of RECIST 1.1 criteria assessments as appropriate.

14. Observation & Follow-Up Schedule

Total Duration: Active treatment until progression + up to ~12 months survival follow-up. **Onsite Visit Cadence:** Cycle 1 (Days 1, 8, 15); Day 1 of all subsequent 21-day cycles. **Remote Monitoring Frequency:** Adverse event and concomitant medication checks between cycles. **Checklist Review Interval:** Pre-infusion clearance every 3 weeks.

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				