

1. Administrative & Study Identification

Posting ID: 1026907402041758790 **Registry Study Title:** A Study Evaluating the Safety and Efficacy of Multiple Treatments in Participants With Multiple Myeloma **Official Full Study Title:** A Platform Study Evaluating the Safety and Efficacy of Multiple Treatments in Patients With Multiple Myeloma **Short Title/Acronym:** PLYCOM **Protocol Number:** C043923 **NCT Number:** NCT05583617 **Sponsor/Company Name:** Roche (Hoffmann-La Roche) **Investigational Product:** Cevostamab, Lenalidomide, Iberdomide, Tocilizumab, Dexamethasone (*Dexamethasone Trade Names: Dexium, Baycadron, DexaJect, Neopolydex, Ozurdex, Eyetrol, Ciprodex, Robadex, Ocu-Dex, Ocu-Trol, Decadron, Maxidex, Dexasone, Hexadrol, Aeroseb-Dex, Decaspray, AK-Dex, AK-Neo-Dex, AK-Trol, Dalalone, De-Sone LA, Decadron-LA, Decadron with Xylocaine, Dexacidin, Dexacine, Dexacort Phosphate in Turbinaire, Dexacort Phosphate in Respihaler, Dexasporin, Dexacort-LA, Dexasol, Dexamethasone Intensol, Methadex, Neo-Dex, Neo-Decadron, Neo-Dexair, Poly-Dex, Tobradex, Maxitrol, Tresaderm, Hemady, Dextenza, Dexycu*) **Phase of Development:** Phase 1 / Phase 2 (PHASE1, PHASE2) **Criteria Type:** Phase Requirement **Trial Status:** RECRUITING (Currently recruiting participants) **Disease Area / Preferred Name:** Multiple Myeloma **Medical Monitor:** Hoffmann-La Roche Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective **Primary Objective:** To evaluate the safety, efficacy, and pharmacokinetics (PK) of multiple treatment combinations, as monotherapy or in combination, in participants with multiple myeloma (MM). **Secondary Objectives:** To determine the target dose of cevostamab in combination with iberdomide or lenalidomide, and to assess preliminary anti-tumor activity (including Objective Response Rate, Complete Response, and Progression-Free Survival) in specific MM subpopulations.

2.2 Background & Rationale To address the gap in effective therapeutic options for patients with Multiple Myeloma (MM) who have become refractory to standard lines of therapy (including proteasome inhibitors [PI], IMiDs, and anti-CD38 monoclonal antibodies), or who present with high-risk cytogenetic features post-transplant. This platform study provides a flexible framework to test novel treatments, including the bispecific antibody cevostamab, in combination with next-generation agents to improve depth and duration of response.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional (Platform Study)
Control Type: Single-arm (within respective substudies) **Blinding:** Open-label **Randomization:** N/A (Dose escalation and expansion cohorts)

3.2 Planned Enrollment & Duration Targeted Enrollment: 200 evaluable patients **Number of Sites:** Global, multi-center
Estimated Study Start: 14-Nov-2023 **Estimated Primary Completion:** 31-Jul-2028

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Age ≥ 18 years with a confirmed diagnosis of Multiple Myeloma (MM) per International Myeloma Working Group (IMWG) criteria. 2. Eastern Cooperative Oncology Group (ECOG) Performance Status of 0, 1, or 2. 3. Resolution of AEs from prior anti-cancer therapy to Grade ≤ 1 . 4. **Substudy 2 (DIRAC):** High-risk cytogenetic features at diagnosis, completion of planned induction therapy, experienced at least a partial response (PR), and Autologous Stem Cell Transplant (SCT) within 100 days prior to first study treatment. 5. **Substudy 4 (CHAWLA):** Relapsed/refractory MM (R/R MM), having received at least three prior lines of therapy including a PI, an IMiD, and an anti-CD38 monoclonal antibody for whom no suitable SOC therapy options are available.

4.2 Exclusion Criteria 1. Pregnant or breastfeeding, or intending to become pregnant during the study or within 5 months after the final dose. 2. Known active bacterial, viral, fungal, mycobacterial, parasitic, or other infection requiring IV antibiotics within 14 days prior to first study treatment. 3. Positive serologic or PCR test results for acute or chronic hepatitis B virus (HBV), hepatitis C virus (HCV), HIV, Epstein-Barr virus (EBV), or CMV. Active symptomatic COVID-19 infection. 4. Prior treatment with CAR T-cell therapy within 12 weeks before starting the pre-phase, or autologous SCT within 100 days (Substudy 4). 5. History of confirmed progressive multifocal leukoencephalopathy, current or past CNS disease, or significant cardiovascular disease that may limit ability to respond to a CRS event. 6. History of autoimmune disease, hemophagocytic lymphohistiocytosis (HLH), or macrophage activation syndrome (MAS).

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: - *Substudy 2:* Cevostamab pre-phase step-up doses on Day 1 and Day 4, target dose on Day 8; subsequently, target dose administered on Day 1 and Day 15 for cycles 1-6 and Day 1 of cycle 7 onwards.

Lenalidomide PO on a 28-day cycle. - *Substudy 4*: Cevostamab step-up doses on Day 1 and Day 4, target dose on Day 8; subsequently, target dose administered on Day 1 of each 21-day cycle.

Iberdomide PO on a 21-day cycle. **Route of Administration:** Intravenous (Cevostamab, Tocilizumab) and Oral (Lenalidomide, Iberdomide, Dexamethasone). **Duration of Treatment:** Up to 14 treatment cycles (Substudy 2) or 18 treatment cycles (Substudy 4), or until disease progression/unacceptable toxicity.

5.2 Concomitant & Prohibited Therapy Permitted: Tocilizumab for the prophylactic or symptomatic treatment of cytokine release syndrome (CRS), and standard MM supportive care. **Prohibited:** Systemic immunosuppressive medications, live attenuated vaccines within 4 weeks prior to initiation, and concurrent administration of strong CYP3A4/5 modulators or inhibitors.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, Cytogenetic profiling, Viral PCRs
Baseline (Pre-Phase)	Cycle 1	Pre-phase Cevostamab step-up dosing (Days 1, 4), CRS monitoring
Interim	Day 1 of each cycle	Safety Labs, Efficacy Assessment (IMWG criteria), IV/PO drug administration
End of Study	Follow-up	Up to 5 months safety follow-up after the final dose of study treatment

7. Outcome Measures (Endpoints)

7.1 Primary Endpoints 1. **Stage 1:** Percentage of Participants with Adverse Events (AEs). 2. **Stage 2:** Objective Response Rate (ORR). 3. **Stage 2:** Complete Response (CR) or Stringent Complete Response (sCR) Rate.

7.2 Secondary Endpoints 1. **Stage 1:** Conversion to a Better Response. 2. **Stage 1:** Progression-Free Survival (PFS). 3. **Stages 1 and 2:** Duration of Response (DOR).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring with specific protocols for Cytokine Release Syndrome (CRS) and Immune effector cell-associated neurotoxicity syndrome (ICANS). **Serious Adverse Events (SAEs):** Routine reporting; immediate hospitalization/monitoring required for suspected CRS events. **Data Monitoring Committee (DMC):** Internal Safety Review Committee evaluating step-up dosing and cohort dose escalations/expansions.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) for efficacy; Safety Set for all participants receiving at least one dose of study medication. **Sample Size Justification:** Platform adaptive design utilizing dose escalation rules, powering expansion cohorts to approximately 200 participants to demonstrate adequate precision for Objective Response Rate (ORR). **Handling of Missing Data:** Standard non-responder imputation (NRI) for categorical response endpoints; censoring for time-to-event endpoints.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** High-frequency onsite monitoring during Cycle 1 to ensure strict adherence to step-up dosing and CRS management protocols. **Audit Trail:** Detailed tracking of patient compliance with the Lenalidomide/Iberdomide global pregnancy prevention programs and eCRF locking.

11. Study Identifiers & Governance

Primary ID: C043923 **Registry Number:** NCT05583617 **Posting ID:** 1026907402041758790 **Sponsor/Lead Organization:** Roche **Study Title:** PLYCOM **Official Regulatory Title:** A Platform Study Evaluating the Safety and Efficacy of Multiple Treatments in Patients With Multiple Myeloma

12. Clinical Framework (Multiple Myeloma Specific)

Primary Condition: Multiple Myeloma (MM) **Diagnosis Threshold:** Relapsed/Refractory MM or High-Risk MM post-transplant **Medical Methodology:** Bispecific T-cell engager (Cevostamab) + IMiDs (Lenalidomide/Iberdomide) + Dexamethasone **Optimization Target:** Target dose determination for combination regimens

13. Study Procedures & Methodology

Management Pathway: Dose escalation followed by dose expansion **Education Protocol (HCP):** Comprehensive site training on Cytokine Release Syndrome (CRS) identification and Tocilizumab administration **Education Protocol (Patient):** Risk minimization plan including the global pregnancy prevention program for IMiDs **Quality Audit Mechanism:** Adherence monitoring for oral IMiD cycles and continuous safety/toxicity tracking

14. Observation & Follow-Up Schedule

Total Duration: Up to 14-18 treatment cycles plus 5 months safety follow-up **Onsite Visit Cadence:** Frequent during Cycle 1 (Days 1, 4, 8, 15); subsequently Day 1 of each 21-day or 28-day cycle **Remote Monitoring Frequency:** Continuous monitoring per platform protocol **Checklist Review Interval:** Every cycle (Q3W or Q4W depending on substudy arm)

15. Sign-off & Approval

Role	Name	Signature	Date	---	---	---	---
Principal Investigator	[Insert Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Insert Name]	_____	[DD-MMM-YYYY]				
Lead Statistician	[Insert Name]	_____	[DD-MMM-YYYY]				