

1. Administrative & Study Identification

Full Study Title: A PHASE 1, NON-RANDOMIZED, OPEN-LABEL, SINGLE-DOSE, PARALLEL GROUP STUDY TO COMPARE THE PHARMACOKINETICS OF VEPDEGESTRANT (PF-07850327) IN ADULT PARTICIPANTS WITH MODERATE AND SEVERE HEPATIC IMPAIRMENT RELATIVE TO HEALTHY PARTICIPANTS WITH NORMAL HEPATIC FUNCTION **Short Title/Acronym:** A Study to Learn How the Body Processes the Study Medicine Called Vepdegestrant in People With Loss of Liver Function **Protocol Number:** C4891004 **NCT Number:** NCT07231991 **Posting ID:** 1045308424203254342 **Trial Status:** NOT_YET_RECRUITING **Sponsor/Company Name:** Pfizer **Investigational Product:** Vepdegestrant (ARV-471, PF-07850327) **Phase of Development:** Phase 1 **Medical Monitor:** Pfizer Clinical Operations Lead / Pfizer CT.gov Call Center

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate and compare the pharmacokinetics (PK) of vepdegestrant in adult participants with moderate and severe hepatic impairment relative to healthy participants with normal hepatic function after a single oral dose. **Secondary Objectives:** To evaluate the overall safety and tolerability of a single dose of vepdegestrant in these participant populations.

2.2 Background & Rationale Vepdegestrant is an investigational, orally administered PROTAC (PROteolysis TArgeting Chimera) estrogen receptor (ER) degrader. Because the medicine is metabolized and eliminated by the body, this study is necessary to determine how a loss of liver function (hepatic impairment) alters the body's ability to process the drug. The findings will help establish appropriate dosing recommendations for patients who have varying degrees of liver impairment.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Parallel-group (Healthy participants with normal hepatic function vs. participants with hepatic impairment) **Blinding:** Open-label **Randomization:** Non-randomized

3.2 Planned Enrollment & Duration Targeted Enrollment (\$N\$): 24 (Phase 1 parallel cohort size: participants divided into Class B, Class C, and healthy matched control groups). **Number of Sites:** Conducted at dedicated Phase 1 clinical research facilities.

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Female participants of non-childbearing potential, or male participants. 2. Between 18 and 70 years of age (inclusive) at the time of screening. 3. Body Mass Index (BMI) of 17.5-38 kg/m², inclusive, and a total body weight ≥ 50 kg (110 lb). 4. **Normal hepatic function group:** Overtly healthy as determined by medical evaluations (history, physical exam, labs, ECGs). 5. **Hepatic impairment group:** Must satisfy the criteria for Class B (moderate) or Class C (severe) of the modified Child-Pugh Classification, with stable hepatic impairment (no clinically significant change in disease status within the last 28 days).

4.2 Exclusion Criteria 1. **Normal hepatic function group:** Evidence or history of clinically significant hematological, renal, endocrine, pulmonary, gastrointestinal, cardiovascular, hepatic, psychiatric, neurological, or allergic disease. 2. **Hepatic impairment group:** A diagnosis of hepatic dysfunction secondary to any acute ongoing hepatocellular process (documented by medical history, liver biopsy, ultrasound, CT, or MRI). 3. Any condition possibly affecting drug absorption.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: A single 200 mg dose of vepdegestrant. **Route of Administration:** Oral. **Duration of Treatment:** Single-dose administration, requiring an 11-day and 10-night continuous stay at the study clinic for observation and PK sampling.

5.2 Concomitant & Prohibited Therapy Permitted: Standard maintenance medications for stable hepatic impairment (as approved by the investigator). **Associated Medications (Data Registry Mapping):** Naphazoline (ATC Codes: S01GA01, R01AB02, R01AA08). Trade names associated in registry: Advanced Eye Relief Redness Instant Relief, Estivin, Degest, Nazil, Floril Redness Relief, Napha Forte, All Clear, Ocu-zoline, Privine, Naphcon, AK-Con, Albalon, Allerest Eye Drops, Allersol, Allersol A, Naphcon A, Nafazair, Ocuhist, Opcon-A, VasoClear A, VasoClear, Vasocon, AK-Con-A, Naphcon forte, Vasocon-A, Optivet Allergy Relief. **Prohibited:** Concomitant use of strong CYP3A4 inhibitors or inducers, which could interfere with the metabolism of vepdegestrant, as well as any other unapproved investigational agents.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to -2	Informed Consent, Medical History, Child-Pugh Assessment, Vital Signs, Labs
Baseline / Check-in	Day -1	Admission to the clinical facility, Baseline Safety Labs
Dosing & PK Period	Day 1 to 10	Single 200 mg Dose (Day 1), Serial PK Blood Sampling, Vital Signs, AE Monitoring
Discharge	Day 11	Final Inpatient Safety Assessments, Clinic Discharge
End of Study	Day 28 to 35	Safety Follow-up Contact

7. Outcome Measures (Endpoints)

7.1 Primary Endpoints 1. Maximum observed concentration (C_{max}) for vepdegestrant. 2. Area under the plasma concentration versus time curve from time zero to extrapolated infinite time [$AUC(0 - \infty)$] for vepdegestrant. It is obtained from $AUC(0 - t)$ plus $AUC(t - \infty)$.

7.2 Secondary & Exploratory Endpoints 1. Number of Participants With Treatment-Emergent Adverse Events. 2. Number of Participants With Clinically Significant Clinical Laboratory Abnormalities. 3. Number of Participants With Clinically Significant Change From Baseline in Vital Signs. 4. Additional pharmacokinetic parameters (e.g., T_{max} , half-life) for vepdegestrant and its metabolites.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring during the 11-day / 10-night inpatient clinic stay, and through the end of the follow-up period. **Serious Adverse Events (SAEs):** Evaluated from the signing of informed consent through the final follow-up contact at 28 to 35 days post-dose. **Data Monitoring Committee (DMC):** Internal safety monitoring managed by the Sponsor (Pfizer).

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Pharmacokinetic (PK) Analysis Set for all subjects who receive the study dose and have sufficient concentration data to calculate PK parameters. Safety Set for AE evaluations. **Sample Size Justification:** A standard Phase 1 matching cohort size powered to detect clinically meaningful differences in AUC and C_{max} between hepatically impaired classes (Child-Pugh B & C) and healthy matched controls. **Handling of Missing Data:** Standard non-compartmental PK analysis methodology (concentrations below the limit of quantification are

treated as zero for \$AUC\$ calculation; no imputation for missing samples).

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: High-frequency, continuous onsite monitoring during the inpatient confinement period to ensure precise PK sample timing and safety observation. **Audit Trail:** Precise tracking of dosing times, blood draw timestamps, and sample custody for central laboratory PK analysis.

11. Study Identifiers & Governance

Primary ID: C4891004 **Posting ID:** 1045308424203254342 **Registry Number:** NCT07231991 **Sponsor/Lead Organization:** Pfizer **Company:** Pfizer **Study Title:** A Study to Learn How the Body Processes the Study Medicine Called Vepdegestrant in People With Loss of Liver Function **Official Regulatory Title:** A PHASE 1, NON-RANDOMIZED, OPEN-LABEL, SINGLE-DOSE, PARALLEL GROUP STUDY TO COMPARE THE PHARMACOKINETICS OF VEPDEGESTRANT (PF-07850327) IN ADULT PARTICIPANTS WITH MODERATE AND SEVERE HEPATIC IMPAIRMENT RELATIVE TO HEALTHY PARTICIPANTS WITH NORMAL HEPATIC FUNCTION **Trial Status:** NOT_YET_RECRUITING

12. Clinical Framework (Hepatic Impairment Specific & Population)

Disease Areas: Healthy Participants / Hepatic Impairment **Preferred UMLS Name / Population Group:** Healthy Volunteers **Semantic Type:** Population Group **Population Definition:** An individual who is or becomes a participant in a research study and has no significant health-related issues. **Population Coding:** MeSH Code: D064368 | SNOMED ID: 106207008 | SNOMED Hierarchy: 440545006|Healthy volunteer|Healthy volunteer (person) **Diagnosis Threshold:** Modified Child-Pugh Classification Class B (moderate) or Class C (severe) **Medical Methodology:** Pharmacokinetic (PK) Evaluation **Optimization Target:** Defining exposure variances to inform future vepdegestrant dose adjustments

13. Study Procedures & Methodology

Management Pathway: Phase 1 confined inpatient pharmacokinetic clinical pathway **Education Protocol (HCP):** Site training on exact PK collection intervals and sample processing **Education Protocol (Patient):** Informed consent detailing the expectations of an 11-day/10-night clinic confinement and frequent blood draws **Quality**

Audit Mechanism: Independent PK data verification and standard Phase 1 clinical auditing

14. Observation & Follow-Up Schedule

Total Duration: Approximately 35 days post-dose **Onsite Visit Cadence:** Continuous confinement from Day -1 through Day 10 **Remote Monitoring Frequency:** Follow-up contact occurring 28 to 35 calendar days after study intervention **Checklist Review Interval:** N/A (Continuous inpatient review)

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				