

1. Administrative & Study Identification

Full Study Title: An Open Label, Multi-center Asciminib Roll-over Study to Assess Long-term Safety in Patients Who Have Completed a Novartis Sponsored Asciminib Study and Are Judged by the Investigator to Benefit From Continued Treatment **Short Title/ Acronym:** Asciminib Roll-over Study **Protocol Number:** CABL001A2001B **NCT Number:** NCT04877522 **Trial Status:** Recruiting **Sponsor/Company Name:** Novartis Pharmaceuticals **Investigational Product:** Asciminib (single agent, pediatric formulation, or in combination), Imatinib, Nilotinib, Bosutinib, Dasatinib **Phase of Development:** Phase IV **Medical Monitor:** Novartis Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To assess the long-term safety and provide continued treatment to participants who have previously participated in an asciminib Novartis-sponsored study. **Secondary Objectives:** To evaluate the ongoing efficacy, treatment compliance, and tolerability of asciminib and other continued TKIs in patients who completed a parent study. **2.2 Background & Rationale** To address the clinical necessity of providing continuous, uninterrupted therapy. This roll-over study is conducted to provide continued treatment to participants who have previously participated in an asciminib Novartis-sponsored study and who, in the opinion of the investigator, would benefit from continued treatment as in their parent study but are unable to access this treatment outside of the clinical study.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional (Roll-over study) **Control Type:** Single-arm **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target \$N\$: 347 evaluable patients **Number of Sites:** Multi-center, global (sites across Europe, North America, South America, Asia-Pacific, and Middle East) **Estimated Study Start:** 30-Aug-2022 **Estimated Primary Completion:** 29-Aug-2030

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Aged 7 years or older. 2. Participant with Philadelphia chromosome-positive (PH+) Chronic Myelogenous

Leukemia (CML) or PH+ Acute Lymphoblastic Leukemia (ALL) currently receiving treatment with asciminib (single agent or in combination with imatinib, nilotinib or dasatinib), imatinib, nilotinib, or bosutinib alone within a Novartis-sponsored study. 3. In the opinion of the Investigator, the participant would benefit from continued treatment. 4. Participant has demonstrated compliance on the parent study protocol and is willing and able to comply with scheduled visits, treatment plans and any other study procedures.

4.2 Exclusion Criteria 1. Participant has been discontinued from parent study treatment. 2. Participant currently has unresolved toxicities reported as possibly related to study treatment in the parent study. 3. Participant's ongoing treatment is currently approved and reimbursed at the country level. 4. Pregnant or nursing (lactating) women. 5. Women of child-bearing potential, unless they are using highly effective methods of contraception and willing to continue while taking study treatment. 6. Sexually active males receiving imatinib, nilotinib, bosutinib or dasatinib unwilling to follow the relevant contraception requirements in the local prescribing information. 7. Applicable for participants on bosutinib treatment at the end of the CABL001A2301 and on other TKIs for CABL001A2202 study that switch to asciminib treatment: * Asymptomatic (grade 2) pancreatitis if not resolved within 28 days. * QTcF >480msec or inability to determine QTc interval. * Any grade 3 or 4 toxicity not resolved to grade 2 or lower within 28 days before starting asciminib treatment.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Dosing schedule directly continues the regimen established in the participant's respective parent study. **Investigational Agents:** Asciminib, Imatinib (Trade name: Gleevec, ATC: L01EA01), Nilotinib, Dasatinib, or Bosutinib. **Route of Administration:** Oral **Duration of Treatment:** Long-term continued treatment until treatment discontinuation criteria are met or the drug becomes commercially accessible/reimbursed in the patient's country.

5.2 Concomitant & Prohibited Therapy Permitted: Standard supportive care and concomitant medications allowed during the parent study protocol. **Prohibited:** Any investigational agents outside of the specified parent trial protocol or medications with known severe drug-drug interactions with the specific TKIs administered.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Screening/Transition	Day -14 to 0 Informed

Consent, Medical History, Verification of Parent Study Compliance
| | **Baseline** | Day 0 | Roll-over Enrollment, Continued Dispensing
| | **Interim Visits** | Periodic (per parent protocol) | Long-term
Safety Labs, Efficacy Assessment, AE Monitoring, Drug
Dispensation | | **End of Study** | At Discontinuation | Final Vital
Signs, Exit Interview, IP Accountability, Transition to
commercial drug if applicable |

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Assessment of long-term safety, defined by the number of participants with adverse events (AEs) and serious adverse events (SAEs). All AEs and SAEs will be tabulated and listed for participants in the Safety Set by treatment group. **Timeframe:** From day of first administration of study treatment to 30 days after the last study treatment. **Measurement Method:** Clinical assessment and central/local laboratory evaluations graded by CTCAE criteria.

7.2 Secondary & Exploratory Endpoints Efficacy: Percentage of participants with clinical benefit as assessed by Investigator. **Maintenance:** Maintenance of clinical/molecular response (e.g., MMR or complete cytogenetic response) over time. **Survival:** Overall survival and progression-free survival in the extended follow-up phase.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous collection via patient interviews and safety labs at scheduled periodic visits. **Serious Adverse Events (SAEs):** Standard 24-hour reporting timeline to the sponsor safety desk. **Data Monitoring Committee (DMC):** Supervised by internal and external safety monitoring boards configured for Novartis oncology programs.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all enrolled patients who received at least one dose of the roll-over study drug). **Sample Size Justification:** $N = 347$ evaluable patients, determined organically by the number of eligible, consenting patients successfully completing their respective Novartis-sponsored parent studies. **Handling of Missing Data:** Standard handling for long-term safety extension trials (e.g., missing safety data are typically not imputed; time-to-event endpoints right-censored at last known contact).

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Frequency of onsite and remote monitoring visits adjusted according to the roll-over phase requirements and local regulations. **Audit Trail:** Standard electronic Case Report Form (eCRF) locking procedures and data clarification protocols managed by Novartis.

11. Study Identifiers & Governance

Primary ID: CABL001A2001B **Registry Number:** NCT04877522 **Posting ID:** 1086871291634538169 **Sponsor/Lead Organization:** Novartis **Study Title:** Asciminib Roll-over Study **Official Regulatory Title:** An Open Label, Multi-center Asciminib Roll-over Study to Assess Long-term Safety in Patients Who Have Completed a Novartis Sponsored Asciminib Study and Are Judged by the Investigator to Benefit From Continued Treatment

12. Clinical Framework (Leukemia Specific)

Primary Condition / Disease Areas: Chronic Myelogenous Leukemia (CML), Leukemia, Myelogenous, Chronic, BCR-ABL Positive / Acute Lymphoblastic Leukemia (ALL) **Diagnosis Threshold:** Philadelphia chromosome-positive (PH+) status **Coding Classifications:** MeSH Code: D007938 | SNOMED ID: 93143009 **Medical Methodology:** Continued Tyrosine Kinase Inhibitor (TKI) Therapy **Optimization Target:** Sustained disease control and long-term safety maintenance

13. Study Procedures & Methodology

Management Pathway: Long-term roll-over clinical pathway for maintenance therapy. **Education Protocol (HCP):** Protocol training, rigorous AE/SAE long-term reporting procedures. **Education Protocol (Patient):** Informed consent regarding long-term therapy and contraception adherence. **Quality Audit Mechanism:** Routine clinical site monitoring and centralized safety database tracking.

14. Observation & Follow-Up Schedule

Total Duration: Up to 8 years (Estimated 2022 to 2030) **Onsite Visit Cadence:** Aligned with the individualized schedule of the patient's specific parent study or local standard of care. **Remote Monitoring Frequency:** As indicated by site capability and parent protocol structure. **Checklist Review Interval:** Routine

verification of inclusion/exclusion and concomitant medications at each dispensing visit.

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				