

1. Administrative & Study Identification

Full Study Title A Phase 3, Multicenter, Long-Term, Open Label Study Evaluating the Safety and Efficacy of Abrocitinib, With or Without Topical Medications Administered to Pediatric Participants Aged 2 Years and Older With Moderate-to-Severe Atopic Dermatitis **Short Title/Acronym** A Long-term Study of the Medicine Called Abrocitinib in Children Aged 2 Years and Older With Moderate to Severe Eczema **Posting ID** 1123380521116202033 **Protocol Number** B7451031 **NCT Number** NCT06807281 **Trial Status** NOT_YET_RECRUITING (Not yet recruiting participants) **Sponsor/Company Name** Pfizer **Investigational Product** Abrocitinib (CIBINQO™) **Phase of Development** Phase III (Phase Requirement: PHASE3) **Medical Monitor** Pfizer Clinical Trial Contact Center

2. Study Synopsis & Rationale

2.1 Study Objective * **Primary Objective:** To evaluate the long-term safety and efficacy of abrocitinib oral liquid suspension or tablets in pediatric participants aged 2 to 11 years with moderate-to-severe atopic dermatitis. * **Secondary Objectives:** To evaluate long-term maintenance of clinical response (e.g., VIGA, EASI scores) and symptom relief, and to measure the tolerability of daily oral dosing over a prolonged period.

2.2 Background & Rationale Managing moderate-to-severe atopic dermatitis (AD) in young children is challenging, as topical treatments are not always effective, and other systemic medications may cause unwanted side effects or be difficult for children to tolerate. Abrocitinib (CIBINQO™), an oral JAK inhibitor, is currently approved for the treatment of moderate-to-severe eczema in people 12 years and older. This study is being conducted to evaluate it as a potential oral treatment option for children under 12 years of age.

3. Study Design & Methodology

3.1 Design Framework * **Study Type:** Interventional * **Control Type:** Single-arm (Open-label) * **Blinding:** Open-label * **Randomization:** N/A (Participants are assigned to an extension cohort from prior studies or a *de novo* cohort)

3.2 Planned Enrollment & Duration * **Target \$N\$:** 500 participants (up to 320 in the extension cohort; approx. 180 in the *de novo* cohort) * **Number of Sites:** Global multicenter study (including USA, Japan, Poland, Hungary) * **Estimated Study Start:** 02-Dec-2025 * **Estimated Primary Completion:** 22-Feb-2032 (Study Completion)

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Children 2 to 11 years of age, weighing at least 15 kg (33 lbs). 2. Documented diagnosis of chronic AD for at least 6 months prior to screening, with moderate-to-severe AD at baseline (BSA $\geq 10\%$, vIGA ≥ 3 , EASI ≥ 16 , and WI-NRS ≥ 4) for the *de novo* cohort. 3. Documented history of inadequate response to treatment with topical medical therapy for AD (e.g., TCS and TCI) for at least 4 weeks and are candidates for systemic therapy.

4.2 Exclusion Criteria 1. Prior treatment with a systemic JAK inhibitor for AD (for *de novo* participants). 2. Infection with HIV, hepatitis B, hepatitis C, active TB, or inadequately treated latent TB. 3. Live attenuated vaccination within 6 weeks prior to Day 1 or require vaccination with live attenuated vaccines during treatment or within 6 weeks after the last dose.

5. Intervention & Treatment Plan

5.1 Investigational Regimen * **Drug Name:** Abrocitinib * **Trade Name:** CIBINQO™ * **ATC Code:** D11AH08 * **EMA URL:** [CIBINQO EPAR Product Information](#) * **Dosage/Frequency:** Abrocitinib taken by mouth once daily (dose may be increased if AD does not improve within the first 3 months). * **Route of Administration:** Oral (liquid suspension or tablet). * **Duration of Treatment:** Up to 2 years (24 months) or commercial availability.

5.2 Concomitant & Prohibited Therapy * **Permitted:** Medicated and non-medicated topical therapies for AD as needed throughout the clinical trial. * **Prohibited:** Systemic JAK inhibitors and live attenuated vaccines.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, Confirmation of AD chronicity and severity (EASI, vIGA)
Baseline	Day 0	Cohort Assignment, First Dose, Safety Labs
Interim	Up to 2 Years	Safety Labs (LFTs, CBCs), Efficacy Assessment, Adjusting dose if no improvement in first 3 months
End of Study	Month 24	Final Vital Signs, Exit Interview, IP Accountability

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint * **Definition:** Long-term safety and tolerability of abrocitinib in pediatric patients, assessed by the incidence of treatment-emergent adverse events (TEAEs), serious adverse events (SAEs), adverse events leading to discontinuation, and clinical laboratory abnormalities over the

2-year period. * **Measurement Method:** Clinical evaluation, vital signs, and central laboratory analysis.

7.2 Secondary & Exploratory Endpoints * Laboratory Monitoring:

Number of participants with clinically significant laboratory abnormalities. * **Clinical Response:** Achievement of Validated Investigator's Global Assessment (vIGA) response of clear (0) or almost clear (1) and a 2-point reduction from baseline. * **Symptom Relief:** Percentage of response based on achieving a ≥ 4 point improvement from baseline in the Worst Itch Numerical Rating Scale (WI-NRS). * Change from baseline in the Eczema Area and Severity Index (EASI) score.

8. Safety & Adverse Event Reporting

- **Adverse Event (AE) Monitoring:** Standard collection via onsite clinical visits, monitoring for serious infections (e.g., herpes zoster, HSV) and lab abnormalities.
- **Serious Adverse Events (SAEs):** Reported to the sponsor within 24 hours of investigator awareness.
- **Data Monitoring Committee (DMC):** Internal and External review committees monitoring safety signals associated with systemic JAK inhibitors.

9. Statistical Analysis Plan (SAP) Summary

- **Analysis Populations:** Safety Set (all patients who receive ≥ 1 dose of abrocitinib); Efficacy Analysis Set.
- **Sample Size Justification:** $N = 500$ ensures adequate long-term exposure data to characterize the safety profile of abrocitinib in the 2-to-11-year-old population.
- **Handling of Missing Data:** Multiple Imputation or Last Observation Carried Forward (LOCF) for clinical efficacy endpoints (EASI, vIGA).

10. Quality Assurance & Regulatory Compliance

- **GCP Compliance:** This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.
- **Monitoring Plan:** Standard onsite and remote monitoring visits to ensure safety and protocol adherence.
- **Audit Trail:** Strict documentation in eCRFs and central laboratory records.

11. Study Identifiers & Governance

- **Primary ID:** B7451031
- **Posting ID:** 1123380521116202033
- **Registry Number:** NCT06807281
- **Trial Status:** NOT_YET_RECRUITING
- **Sponsor/Lead Organization:** Pfizer

- **Study Title:** A Long-term Study of the Medicine Called Abrocitinib in Children Aged 2 Years and Older With Moderate to Severe Eczema
- **Official Regulatory Title:** A Phase 3, Multicenter, Long-Term, Open Label Study Evaluating the Safety and Efficacy of Abrocitinib, With or Without Topical Medications Administered to Pediatric Participants Aged 2 Years and Older With Moderate-to-Severe Atopic Dermatitis

12. Clinical Framework (Atopic Dermatitis Specific)

- **Primary Disease Area:** Atopic Dermatitis
- **Preferred Name (UMLS):** Dermatitis, Atopic
- **Semantic Type:** Disease or Syndrome
- **Definition:** A chronic inflammatory genetically determined disease of the skin marked by increased ability to form reagin (IgE), with increased susceptibility to allergic rhinitis and asthma, and hereditary disposition to a lowered threshold for pruritus. It is manifested by lichenification, excoriation, and crusting, mainly on the flexural surfaces of the elbow and knee. In infants it is known as infantile eczema.
- **Clinical Terminology Codes:**
 - **MeSH Code:** D003876
 - **HPO Code:** HP:0001047
 - **SNOMED ID:** 200775004
- **Diagnosis Threshold:** BSA $\geq 10\%$, vIGA ≥ 3 , EASI ≥ 16 , and WI-NRS ≥ 4
- **Medical Methodology:** Targeted systemic therapy with oral JAK1 inhibitor (abrocitinib) combined with topical medical therapy
- **Optimization Target:** Achievement of clear or almost clear skin and improvement in pruritus over up to 2 years of treatment

13. Study Procedures & Methodology

- **Management Pathway:** Standard clinical pathway for moderate-to-severe AD unresponsive to topicals
- **Education Protocol (HCP):** Training on protocol guidelines, dose escalation steps, and JAK inhibitor safety monitoring
- **Education Protocol (Patient):** Parent/caregiver guidance on administering oral liquid suspension/tablets and applying medicated/non-medicated topical therapies
- **Quality Audit Mechanism:** Safety lab tracking & caregiver reporting of treatment adherence

14. Observation & Follow-Up Schedule

- **Total Duration:** Up to 24 months (104 weeks)
- **Onsite Visit Cadence:** Regular intervals throughout the 2-year duration (e.g., initial baseline, month 3 dose evaluation, followed by long-term maintenance visits)

- **Remote Monitoring Frequency:** As outlined per sponsor protocol
- **Checklist Review Interval:** Continuous reporting of concomitant topical use and adverse events

15. Sign-off & Approval

Role	Name	Signature	Date
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]
Statistician	[Name]	_____	[DD-MMM-YYYY]