

1. Administrative & Study Identification

Full Study Title: A Multicenter, Single-Arm, Open-Label, Extension, Rollover Study To Evaluate The Long-Term Safety And Efficacy Of Ocrelizumab In Patients With Multiple Sclerosis **Short Title/Acronym:** OLERO **Protocol Number:** MN43964 **NCT Number:** NCT05269004 **Trial Status:** Active, Not Recruiting **Sponsor Name:** Hoffmann-La Roche / Roche **Investigational Product:** Ocrelizumab (R04964913) **Phase of Development:** Phase 3 (Phase IIb) **Medical Monitor:** Hoffmann-La Roche Medical Monitor

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the long-term safety and efficacy of ocrelizumab in patients with multiple sclerosis. **Secondary Objectives:** To continue collecting longer-term information on the effectiveness of treatment with ocrelizumab and provide continued access to the study drug for those benefiting from it.

2.2 Background & Rationale This rollover extension study is designed to ensure that participants of parent studies (WA21092, WA21093, or WA25046) who are receiving benefit from ocrelizumab can continue their treatment or safety follow-up without interruption. It addresses the medical necessity of providing continued therapy while allowing for valuable long-term data on the safety and effectiveness of the treatment to be collected.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional (Extension, Rollover Study) **Control Type:** Single-arm **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target \$N\$: 1,300 participants **Number of Sites:** Multicenter (Global) **Estimated Study Start:** 03-May-2022 **Estimated Primary Completion:** 31-Dec-2027

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Prior eligibility for and previous participation in 1 of the Roche-sponsored Parent Studies WA21092, WA21093, or WA25046. 2. Participant either receiving ocrelizumab as an investigational medicinal product (IMP) or being in safety follow-up after treatment discontinuation. 3. For female

participants of childbearing potential: agreement to remain abstinent (refrain from heterosexual intercourse) or use adequate acceptable contraception during the treatment period and for at least 6 months or longer if the local label is more stringent after the final dose of ocrelizumab.

4.2 Exclusion Criteria 1. Concurrent participation in any clinical trial (other than the Parent study). 2. Unable or unwilling to comply with the requirements of the protocol. 3. Permanent premature discontinuation of study treatment for any reason during the parent study (or early discontinuation due to unacceptable safety concerns).

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Ocrelizumab based on the dosage and administration received at the time of rollover from the Parent study (e.g., 600 mg every 24 weeks). **Route of Administration:** Intravenous (IV) infusion (or subcutaneous depending on the parent study protocol). **Duration of Treatment:** Up to 44 months / until the end of the study (estimated December 2027).

5.2 Concomitant & Prohibited Therapy Permitted: Standard MS symptom management therapies as permitted in parent trials. **Prohibited:** Concurrent participation in another therapeutic clinical trial or initiation of other disease-modifying treatments (DMTs) for MS.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Screening	Day -28 to 0 Eligibility confirmation from parent study, Informed Consent Baseline
Day 0	Rollover confirmation, First extension dose	Interim	Every 24 Weeks Safety Labs, Ocrelizumab Infusion, Efficacy Assessment End of Study 48 Weeks Post-Last Dose Safety follow-up period, Final Assessments, Exit Interview

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Incidence and severity of adverse events, with severity determined according to the NCI CTCAE v5.0. **Measurement Method:** Clinical assessment, laboratory tests, and continuous AE reporting at follow-up visits.

7.2 Secondary Endpoints 1. Change in Expanded Disability Status Scale (EDSS) score over time. 2. Change in 9-Hole Peg Test (9HPT) over time. 3. Change in Timed 25-Foot Walk Test (T25FWT) over time.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring at each 24-week visit and via patient reporting of symptoms. **Serious Adverse Events (SAEs):** Reported within 24 hours of site awareness, tracking severe complications or opportunistic infections. **Data Monitoring Committee (DMC):** Governed by internal sponsor safety monitoring and ethics committees transitioning from the parent protocols.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all participants receiving at least one dose in the extension), Intent-to-Treat (ITT) for ongoing efficacy. **Sample Size Justification:** $N=1,300$ derived from the rollover pool of participants completing the parent studies WA21092, WA21093, or WA25046. **Handling of Missing Data:** Standard imputation methods applied as outlined in the parent protocols for long-term longitudinal tracking.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Regular onsite and remote monitoring to verify continued safety, infusion protocols, and data integrity. **Audit Trail:** eCRF locking and tracking of protocol adherence bridging from the parent studies.

11. Study Identifiers & Governance

Primary ID: MN43964 **Registry Number:** NCT05269004 **Sponsor/Lead Organization:** Hoffmann-La Roche / Roche **Study Title:** A Rollover Study to Evaluate the Long-Term Safety and Efficacy of Ocrelizumab In Patients With Multiple Sclerosis **Official Regulatory Title:** A Multicenter, Single-Arm, Open-Label, Extension, Rollover Study To Evaluate The Long-Term Safety And Efficacy Of Ocrelizumab In Patients With Multiple Sclerosis

12. Clinical Framework (Multiple Sclerosis Specific)

Primary Condition: Multiple Sclerosis (Relapsing and Primary Progressive) **Diagnosis Threshold:** Confirmed MS diagnosis per parent study eligibility criteria **Medical Methodology:** Disease-Modifying Therapy (DMT) **Optimization Target:** Long-term disease stability, relapse prevention, and safety tracking

13. Study Procedures & Methodology

Management Pathway: Continued Ocrelizumab treatment for benefiting patients **Education Protocol (HCP):** Infusion reaction management and long-term safety monitoring guidelines **Education Protocol (Patient):** Strict contraception requirements (minimum 6 months post-dose or longer if local label is more stringent) and prompt reporting of AEs **Quality Audit Mechanism:** Sponsor audits of AE reporting, infusion frequency, and parent-protocol bridging adherence

14. Observation & Follow-Up Schedule

Total Duration: Up to 44-60 months (including safety follow-up period) **Onsite Visit Cadence:** Approximately every 24 weeks for infusions **Remote Monitoring Frequency:** Between 24-week infusion visits as clinically indicated **Checklist Review Interval:** Evaluated at each 24-week dosing visit

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				