

1. Administrative & Study Identification

Full Study Title: A Multicentre, Single Country, Prospective Non Interventional Observational Study to Describe the Switching From a First- or Second-line Disease Modifying Therapy (DMT) to Ozanimod, in Relapsing Remitting Multiple Sclerosis (RRMS) Patients Treated According to Clinical Practice
Short Title/Acronym: SWITCH - ITA
Protocol Number: [Protocol-Defined Internal ID]
NCT Number: NCT05658601
Sponsor Name: Bristol-Myers Squibb
Investigational Product: Ozanimod
Phase of Development: N/A (Observational / Non-interventional)
Medical Monitor: Bristol-Myers Squibb Study Director

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To describe the reasons, therapy, and/or disease status for changing from a first- or second-line Disease Modifying Therapy (DMT) to ozanimod in participants with Relapsing Remitting Multiple Sclerosis (RRMS).
Secondary Objectives: To evaluate the Expanded Disability Status Scale (EDSS) score, assess new or enlarging T2/T1 Gadolinium Enhancing (GdE) lesions via MRI, measure treatment satisfaction via the TSQM questionnaire, monitor lymphocyte subpopulations, and track the incidence of Adverse Events (AEs) and Serious Adverse Events (SAEs).

2.2 Background & Rationale To address the real-world clinical patterns of therapeutic switching in MS. Why are we conducting this study now? To understand the clinical rationale (e.g., lack of efficacy, poor tolerability, compliance issues) and the methodology (wash-out periods, overlapping) utilized by healthcare providers when transitioning RRMS patients from previous DMTs to ozanimod under routine clinical practice conditions.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational (Prospective Non Interventional) **Control Type:** Single-arm **Blinding:** Open-label (None) **Randomization:** N/A

3.2 Planned Enrollment & Duration Target \$N\$: 180 participants
Number of Sites: Multicenter, Single Country (Italy) **Estimated**

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Patients aged 18-55 years with RRMS diagnosed according to the 2017 revised McDonald criteria. 2. Patients who switched from a previous first- or second-line DMT to ozanimod between 4 and 12 weeks before enrollment. 3. Patients with an MRI performed within three months before enrollment and eligible for ozanimod according to the Summary of Product Characteristics (SmPC).

4.2 Exclusion Criteria 1. Patients with clinical forms of MS other than RRMS. 2. Patients participating in another clinical study with an investigational product if the study considers the switching behavior as an endpoint or objective. 3. Patients with contraindications to ozanimod according to the SmPC or unable to participate for various reasons.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Ozanimod prescribed and dosed strictly according to routine clinical practice and the SmPC guidelines. **Route of Administration:** Oral. **Duration of Treatment:** Continuous observation up to 42 months.

5.2 Concomitant & Prohibited Therapy Permitted: Routine medications allowed for RRMS symptom management and co-morbidities as per standard clinical practice. **Prohibited:** Concurrent participation in another investigational clinical trial involving DMTs.

6. Study Timeline & Visit Schedule

Visit ID	Window	Key Activities/Procedures	:---	:---		
:---	Baseline	Day 0	Assessment of Reason/Mode of switching, EDSS, MRI, TSQM, Lymphocyte tracking (CD3, CD4, CD8, CD19, CD56)	Interim 1 Week 12 EDSS, MRI, Lymphocyte subpopulation tracking, AE/SAE monitoring	Interim 2 Week 24 EDSS, MRI, TSQM, Lymphocyte subpopulation tracking, AE/SAE monitoring	End of Study Up to 42 months Continuous AE/SAE monitoring and longitudinal safety tracking

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: 1. Reason for switching treatment (e.g., lack of efficacy, poor safety/tolerability,

difficulty in administration, poor compliance, patient's request). 2. Mode of switching treatment (wash-out days, overlapping days, dosing, and concomitant treatments). **Measurement Method:** Documented at Baseline via patient and physician questionnaires/eCRF records.

7.2 Secondary & Exploratory Endpoints * Mean EDSS score changes at baseline, week 12, and week 24. * Number of new or enlarging T2 lesions and T1 GdE lesions via MRI at baseline, week 12, and week 24. * Treatment Satisfaction Questionnaire for Medication (TSQM) at baseline and week 24. * Pattern of lymphocyte subpopulations (CD3, CD4, CD8, CD19, CD56) at baseline, week 12, and week 24.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Number of AEs coded according to MedDRA, monitored continuously for up to 42 months. **Serious Adverse Events (SAEs):** Number of SAEs coded according to MedDRA, monitored continuously for up to 42 months. **Data Monitoring Committee (DMC):** Standard Sponsor/Pharmacovigilance monitoring for post-marketing observational trials.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Full Analysis Set (FAS) / Safety Set (all enrolled participants receiving at least one dose of ozanimod in the real-world setting). **Sample Size Justification:** Target $N=180$ is calculated to provide descriptive statistical power to identify common therapeutic switching patterns and establish a baseline for real-world efficacy/safety. **Handling of Missing Data:** Standard descriptive statistics with missing data handled via non-imputation (as observed in clinical practice).

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP), the Declaration of Helsinki, and applicable local regulations for non-interventional observational studies. **Monitoring Plan:** Real-world data extracted from Electronic Medical Records (EMR) and entered into eCRFs. **Audit Trail:** Standard source data verification (SDV) to ensure alignment between hospital records and study databases.

11. Study Identifiers & Governance

Primary ID: {{NCT05658601}} **Registry Number:** {{NCT05658601}}
Sponsor/Lead Organization: {{Bristol-Myers Squibb}} **Study Title:**

{{SWITCH - ITA}} **Official Regulatory Title:** {{A Multicentre, Single Country, Prospective Non Interventional Observational Study to Describe the Switching From a First- or Second-line Disease Modifying Therapy (DMT) to Ozanimod, in Relapsing Remitting Multiple Sclerosis (RRMS) Patients Treated According to Clinical Practice.}}

12. Clinical Framework (Multiple Sclerosis Specific)

Primary Condition: {{Relapsing Remitting Multiple Sclerosis (RRMS)}} **Diagnosis Threshold:** {{2017 revised McDonald criteria}} **Medical Methodology:** {{Standard Clinical Practice / SmPC guidelines}} **Optimization Target:** {{Therapeutic switching from 1st/2nd line DMT to Ozanimod}}

13. Study Procedures & Methodology

Management Pathway: {{Observational follow-up according to clinical practice}} **Education Protocol (HCP):** {{N/A - Non-interventional}} **Education Protocol (Patient):** {{N/A - Non-interventional}} **Quality Audit Mechanism:** {{eCRF monitoring of real-world switching data}}

14. Observation & Follow-Up Schedule

Total Duration: {{Up to 42 months}} **Onsite Visit Cadence:** {{Baseline, Week 12, Week 24}} **Remote Monitoring Frequency:** {{As per routine clinical practice}} **Checklist Review Interval:** {{Continuous AE/SAE monitoring}}

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]			Lead	
Statistician	[Name]	_____	[DD-MMM-YYYY]				