

1. Administrative & Study Identification

Full Study Title: A PHASE 1, RANDOMIZED STUDY WITH DOUBLE-BLIND AND SPONSOR-OPEN, PLACEBO-CONTROLLED SINGLE- AND MULTIPLE-DOSE ESCALATION TO EVALUATE THE SAFETY, TOLERABILITY, PHARMACOKINETICS, AND PHARMACODYNAMICS OF PF-08049820 IN HEALTHY ADULT PARTICIPANTS **Short Title/Acronym:** A Study to Learn How Different Amounts of the Study Medicine Called PF-08049820 Are Tolerated and Act in the Body in Healthy Adults **Protocol Number:** C6231001 **NCT Number:** NCT06686797 **Sponsor Name:** Pfizer **Investigational Product:** PF-08049820 **Phase of Development:** Phase 1 **Medical Monitor:** Pfizer Clinical Operations Lead / Medical Monitor

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the safety and tolerability of single and multiple escalating doses of PF-08049820 in healthy adult participants. **Secondary Objectives:** To evaluate the pharmacokinetics (how the body processes the medicine) and pharmacodynamics of PF-08049820, and to determine if food affects the amount of the study medicine in the blood.

2.2 Background & Rationale The study medicine, PF-08049820, is being developed for the treatment of moderate to severe atopic dermatitis, also known as eczema, which causes severe itching and rashes on the skin. Before proceeding to advanced trials in patients, this Phase 1 study is necessary to learn how different amounts of the drug are tolerated, how it acts in the body, and the impact of food on drug absorption in healthy adults.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Placebo-controlled **Blinding:** Double-blind, Sponsor-open **Randomization:** Randomized

3.2 Planned Enrollment & Duration Target \$N\$: 71 participants **Number of Sites:** Single/Multicenter (Pfizer Clinical Research Unit) **Estimated Study Start:** 27-Nov-2024 **Estimated Primary Completion:** 23-Feb-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Healthy males and females who can no longer have children. 2. Age 18 to 65 years old. 3. Body mass index (BMI) of 16 to 32 kg/m² and a total body weight of >50 kg (110 lbs). (Note: Cohort 3 is specific to Japanese participants with 4 biological Japanese grandparents and a total body weight >45 kg).

4.2 Exclusion Criteria 1. Evidence or history of clinically significant medical conditions. 2. History of human immunodeficiency virus (HIV) infection, hepatitis B, or hepatitis C. 3. History of alcohol abuse, binge drinking, and/or any illicit drug use or dependence within 6 months of Screening.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Escalating single and multiple doses divided into 3 parts: * **Part A (Cohorts 1-3):** Up to 4 dosing periods of a single dose. * **Part B (Cohorts 4-7):** Multiple doses in a single dosing period. * **Part C (Cohorts 8-9):** Single dose in a single dosing period.

Route of Administration: Oral (Liquid solution/suspension for Part A; Tablets for Parts B and C). **Duration of Treatment:** Up to 4 dosing periods (with approximately 8-day in-clinic stays per period).

5.2 Concomitant & Prohibited Therapy Permitted: Standard clinical meals (fed or fasted status assigned by cohort rules).

Prohibited: Participation in studies of other investigational products (drug or vaccine) within 30 days, alcohol, and illicit drugs.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
1	-28 to 0	Informed Consent, Medical History, Eligibility Check
2	Day 1 to ~Day 8	In-clinic stay: Dosing (fasted/fed), PK/PD blood and urine sampling, Safety monitoring
3	Day 14 to ~Day 22	Washout (Part A)
4	Day 28 to ~Day 36	Minimum 2-week gap between dosing periods (Cohorts 1 & 2)
5	~1 Month	Telephone follow-up about 1 month after the last dose

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Incidence of Treatment-Emergent Adverse Events (TEAEs) and Serious Adverse Events (SAEs) to evaluate safety and tolerability. **Measurement Method:** In-clinic

observation, physical examinations, vital signs, and standard laboratory tests.

7.2 Secondary & Exploratory Endpoints * Pharmacokinetic parameters (e.g., maximum concentration (\$C_{max}\$), area under the curve (\$AUC\$)) of PF-08049820. * Effect of food on the pharmacokinetics of the study medicine.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring during the ~8-day in-clinic stay and via follow-up telephone call ~1 month post-dose. **Serious Adverse Events (SAEs):** Standard Phase 1 reporting timeline, typically within 24 hours of investigator awareness. **Data Monitoring Committee (DMC):** Internal Dose Escalation Committee to review safety data before advancing to higher dose cohorts.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all participants who received at least one dose of the investigational product), PK Analysis Set. **Sample Size Justification:** Typical for Phase 1 dose-escalation studies (e.g., 6-8 per cohort) to provide initial safety and pharmacokinetic profiling without being powered for formal statistical hypothesis testing. **Handling of Missing Data:** Summarized as observed for safety; PK profiles with missing values handled according to standard non-compartmental analysis rules.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Continuous onsite monitoring during inpatient clinic stays and routine sponsor monitoring visits. **Audit Trail:** Standard eCRF data entry, query resolution workflows, and database locking procedures.

11. Study Identifiers & Governance

Primary ID: C6231001 **Registry Number:** NCT06686797 **Sponsor/Lead Organization:** Pfizer **Study Title:** A Study to Learn How Different Amounts of the Study Medicine Called PF-08049820 Are Tolerated and Act in the Body in Healthy Adults **Official Regulatory Title:** A PHASE 1, RANDOMIZED STUDY WITH DOUBLE-BLIND AND SPONSOR-OPEN, PLACEBO-CONTROLLED SINGLE- AND MULTIPLE-DOSE ESCALATION TO

12. Clinical Framework (Phase 1 / Atopic Dermatitis Specific)

Primary Condition: Healthy Volunteers (Therapeutic Target: Moderate to Severe Atopic Dermatitis / Eczema) **Diagnosis Threshold:** Healthy verification (BMI 16-32 kg/m², Weight >50 kg) **Medical Methodology:** Single- and Multiple-Dose Escalation **Optimization Target:** Pharmacokinetic profiling and Safety/Tolerability assessment

13. Study Procedures & Methodology

Management Pathway: In-patient Phase 1 clinical monitoring **Education Protocol (HCP):** Investigator brochure and protocol-specific training **Education Protocol (Patient):** Informed consent process and clinic rules orientation **Quality Audit Mechanism:** Source data verification and clinical site monitoring by sponsor

14. Observation & Follow-Up Schedule

Total Duration: Up to several months per participant (includes washout periods and a 1-month follow-up) **Onsite Visit Cadence:** In-clinic stays of approximately 8 days per dosing period **Remote Monitoring Frequency:** Telephone follow-up approximately 1 month after the last dose **Checklist Review Interval:** Daily during in-clinic confinement

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				