

1. Administrative & Study Identification

Full Study Title: A Phase 3, Randomized, Double-blind, Placebo-controlled Study to Evaluate the Efficacy and Safety of Rimegepant in Migraine Prevention in Children and Adolescents ≥ 6 to <18 years of age **Short Title/Acronym:** Efficacy and Safety Study of Rimegepant for the Preventative Treatment of Migraine in Pediatric Subjects **Protocol Number:** BHV3000-315 / C4951009 **NCT Number:** NCT05156398 **Posting ID:** 292342933567513069 **Sponsor Name:** Pfizer (Biohaven Pharmaceuticals Holding Company Limited) **Investigational Product:** Rimegepant (Trade Name: Nurtec, ATC Code: N02CD06) Orally Disintegrating Tablet (ODT) **Phase of Development:** Phase III **Medical Monitor:** Pfizer Clinical Trial Contact Center

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To compare the efficacy and safety of rimegepant to placebo as a preventative treatment for migraine in children and adolescents ≥ 6 to <18 years with episodic migraine. **Secondary Objectives:** To evaluate the improvement in pediatric quality of life (PedsQL), the proportion of subjects achieving a $\geq 50\%$ reduction in mean monthly migraine days, and the reduction in migraine days in the first 4 weeks.

2.2 Background & Rationale Migraine is a debilitating disorder that affects about 7.7% to 9.1% of children and increases to about 15% in adolescents. While rimegepant (an oral calcitonin gene-related peptide [CGRP] receptor antagonist) is approved for both the acute and preventive treatment of migraine in adults, there is a critical medical necessity to evaluate its safety, tolerability, and preventative efficacy in the pediatric population (aged 6 to 17), providing young patients with targeted, innovative therapeutic options.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Placebo-controlled **Blinding:** Double-blind **Randomization:** 1:1 Randomized

3.2 Planned Enrollment & Duration Target N : 640 participants (expected to take the study drug out of ~1,100 screened) **Number of Sites:** Approximately 100 study sites (including US, Canada,

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Male and female subjects ≥ 6 to <18 years of age (subjects must be less than 18 at the time of signing assent/consent). 2. Subject has at least a 6-month history of migraine (with or without aura), including: - 14 or less headache days per month during the 3-month period prior to the Screening Visit. - 6 or more migraine days during the Observation Period. - 14 or less headache days during the Observation Period. 3. Pediatric Migraine Disability Assessment Scale (PedMIDAS) Disability Score of >10 to ≤ 50 , indicating mild (score of 11 to 30) or moderate (score of 31 to 50) disruption in daily activities, as assessed at the Baseline (Randomization) Visit. 4. Ability to verbally distinguish migraine attacks from tension/cluster or other types of headaches. 5. Migraine attacks, on average, lasting 4 - 72 hours if untreated. 6. Subjects must have a weight of ≥ 40 kg (child cohort requirement ≥ 15 kg) at the Screening Visit. 7. Subjects on prophylactic migraine medication are permitted to remain on therapy if the dose has been stable for at least 3 months (12 weeks) prior to the Screening Phase, and the dose is not expected to change during the course of the study.

4.2 Exclusion Criteria 1. Subjects with a history of basilar migraine, cluster headaches, or hemiplegic migraine. 2. Subject has a continuous migraine (defined as an unrelenting headache) within 1 month prior to Screening Visit. 3. Subject has a history or diagnosis of complications of migraine. 4. Subject has a confounding and clinically significant pain syndrome that may interfere with the subject's ability to participate in this study. 5. Subject has any current psychiatric condition that is uncontrolled and/or untreated for a minimum of 6 months prior to the Screening Visit. Subjects with a lifetime history of psychosis and/or mania are excluded. 6. History of suicidal behavior or the subject is at risk of self-harm or harm to others. 7. History of major psychiatric disorder. 8. Subject has a current diagnosis or history of substance abuse. 9. Subject has a history of moderate or severe head trauma or other neurological disorder (including seizure disorder) or systemic medical disease that is, in the investigator's opinion, likely to affect central nervous system functioning.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Rimegepant or placebo administered every other day. **Route of Administration:** Oral (Orally Disintegrating Tablet - ODT). **Duration of Treatment:**

12-week Double-blind Study Treatment Phase, followed by a 52-week Open-label Extension Phase.

5.2 Concomitant & Prohibited Therapy Permitted: Subjects on prophylactic migraine medication are permitted to remain on therapy if the dose has been stable for at least 3 months (12 weeks) prior to screening and is not expected to change.

Prohibited: Acute medications or substances that confound the assessment of the pain syndrome or strongly interfere with central nervous system functioning as outlined by the protocol.

6. Study Timeline & Visit Schedule

Visit ID	Window	Key Activities/Procedures
1	4 Weeks	Informed Consent/ Assent, Medical History, Migraine baseline observation (2 on-site visits)
2	12 Weeks	Randomization, Efficacy/Safety tracking via diary (5 on-site visits)
3	52 Weeks	Active Rimegepant treatment for all participants (1 telephone visit, 14 on-site visits)
4	8 Weeks	Final Assessments, Study Exit (2 on-site visits)

(Note: Study participation spans a total of 76 weeks)

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint 1. Change from baseline in the mean number of migraine days per month as measured over the 12-week double-blind phase of the study in adolescents with episodic migraine.

7.2 Secondary & Exploratory Endpoints 1. Achievement of at least a 50% reduction from baseline in mean monthly migraine days over the course of the double-blind phase in adolescents with episodic migraine. 2. Change from baseline in the mean number of migraine days per month in the first 4 weeks (Weeks 1 through 4) of the double-blind treatment phase in adolescents with episodic migraine. 3. The mean change from baseline in the Pediatric Quality of Life (PedsQL™) total score at Week 12 of the double-blind treatment phase in adolescents with episodic migraine.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Close monitoring by a trial doctor of the child's migraine attacks, side effects, and overall health via recurring onsite and telephone visits. **Serious Adverse Events (SAEs):** Continuously tracked, managed, and reported as per standard Phase III pediatric safety protocols. **Data Monitoring Committee (DMC):** Utilized to monitor patient safety continuously throughout the 76-week study lifecycle.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) population for primary efficacy; Safety Set for all subjects receiving at least one dose of study drug. **Sample Size Justification:** Target enrollment of $N=640$ receiving study drug to provide adequate statistical power ($\alpha = 0.05$, $Power = 80\%$) to detect differences in preventative efficacy between the rimegepant and placebo groups. **Handling of Missing Data:** Standard multiple imputation/mixed-effects models for repeated measures (MMRM) for diary and questionnaire endpoints.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Scheduled cross-verification of digital diaries and clinic measurements (23 total on-site visits across the entire study). **Audit Trail:** E-diary locking, standard electronic case report form (eCRF) auditing, and strict subject data protection.

11. Study Identifiers & Governance

Primary ID: BHV3000-315 **Registry Number:** NCT05156398 **Sponsor/Lead Organization:** Pfizer **Study Title:** Efficacy and Safety Study of Rimegepant for the Preventative Treatment of Migraine in Pediatric Subjects **Official Regulatory Title:** A Phase 3, Randomized, Double-blind, Placebo-controlled Study to Evaluate the Efficacy and Safety of Rimegepant in Migraine Prevention in Children and Adolescents ≥ 6 to < 18 years of age

12. Clinical Framework (Migraine Specific)

Primary Condition: Episodic Migraine in Pediatric Subjects (MeSH: D008881 | SNOMED: 37796009 | HPO: HP:0002076) **Diagnosis Threshold:** ≥ 6 migraine days per month and ≤ 14 headache days per month **Medical Methodology:** Calcitonin gene-related peptide (CGRP) receptor antagonist **Optimization Target:** Migraine day reduction and improvement in quality of life (PedsQL)

13. Study Procedures & Methodology

Management Pathway: Preventative treatment pathway for episodic migraine **Education Protocol (HCP):** Standard investigator and site-level training for pediatric patient administration **Education Protocol (Patient):** Daily headache diary adherence,

differentiation of migraine vs. tension/cluster headaches **Quality Audit Mechanism:** Tracking of patient symptom tracking compliance, study drug adherence

14. Observation & Follow-Up Schedule

Total Duration: 76 weeks (inclusive of all phases) **Onsite Visit Cadence:** 2 during Screening, 5 during Double-blind, 14 during Open-label, and 2 during Follow-up **Remote Monitoring Frequency:** Intermittent telephone visits and ongoing diary review **Checklist Review Interval:** Regular safety and efficacy endpoint assessments every 4 weeks during the double-blind phase

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				