

1. Administrative & Study Identification

Brief Study Title: Efficacy and Safety of Deucravacitinib Versus Placebo in Participants With Moderate-to-severe Scalp Psoriasis
Official Title: A Phase 3B/4, Multicenter, Randomized, Double-blind Placebo-controlled Study to Evaluate the Efficacy and Safety of Deucravacitinib in Participants With Moderate-to-severe Scalp Psoriasis
Short Title/Acronym: PSORIATYK SCALP **Protocol Number:** IM011-220 **NCT Number:** NCT05478499 **Posting ID:** 331079207309219613 **Trial Status:** COMPLETED **Sponsor/Company Name:** Bristol Myers Squibb **Investigational Product:** Deucravacitinib (Trade Name: Sotyktu) **ATC Code:** L04AA56 **Phase of Development:** Phase 3b/4 (Phase 4) **Disease Area:** Psoriasis (MeSH: D011565 | SNOMED ID: 9014002 | HPO: HP:0003765 | Semantic Type: Disease or Syndrome) **Medical Monitor:** Head of the Global Submission Management - Clinical Trials (BMS)

2. Study Synopsis & Rationale

2.1 Study Objective **Primary Objective:** To evaluate the efficacy and safety of deucravacitinib compared with placebo at Week 16 in adult participants with moderate-to-severe scalp psoriasis.
Secondary Objectives: To evaluate the achievement of Psoriasis Scalp Severity Index (PSSI) 90 response, improvement in scalp-specific itch numerical rating scale (NRS) score, and overall static Physician Global Assessment (sPGA) 0/1 response at Week 16.

2.2 Background & Rationale Psoriasis is a chronic inflammatory skin disorder, and up to 80% of people living with psoriasis may have scalp involvement. Deucravacitinib is an oral, selective, allosteric tyrosine kinase 2 (TYK2) inhibitor. This study generates dedicated clinical data characterizing its efficacy and safety specifically for moderate-to-severe scalp psoriasis, including patients with less extensive overall body surface area (BSA) involvement (3-10%) who are candidates for systemic therapy.

3. Study Design & Methodology

3.1 Design Framework **Study Type:** Interventional **Control Type:** Placebo-controlled **Blinding:** Double-blind **Randomization:** 1:2 (Placebo to Deucravacitinib)

3.2 Planned Enrollment & Duration **Targeted Enrollment:** 154 participants **Number of Sites:** Multicenter (Approx. 29 locations)

Estimated Study Start: 2022 **Trial Status:** COMPLETED (Primary endpoint analysis at Week 16, followed by 52-week treatment completion).

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Men and women diagnosed with stable plaque psoriasis with scalp involvement for 6 months or more. Stable psoriasis is defined as no morphology changes or significant flares of disease activity in the opinion of the Investigator. 2. Deemed by the Investigator to be a candidate for phototherapy or systemic therapy. 3. Moderate-to-severe scalp psoriasis as defined by scalp-specific Physician's Global Assessment (ss-PGA) ≥ 3 ; $\geq 20\%$ scalp surface area (SSA); Psoriasis Scalp Severity Index (PSSI) ≥ 12 at the Screening visit and Day 1. 4. $\geq 3\%$ of Body Surface Area (BSA) involvement at the Screening visit and Day 1. 5. Evidence of plaque psoriasis in a non-scalp area. 6. Failed to respond to, or intolerant of ≥ 1 topical therapy for scalp psoriasis.

4.2 Exclusion Criteria 1. Target Disease Exceptions: Has nonplaque psoriasis (ie, guttate, inverse, pustular, erythrodermic or drug-induced psoriasis) at Screening or Day 1. 2. History of hypersensitivity reaction to deucravacitinib or any of its excipients. 3. Requirement of prohibited combination therapies, such as other potent immunosuppressants.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Deucravacitinib 6 mg once daily (QD) OR matched oral Placebo. At Week 16, all participants receive deucravacitinib 6 mg QD for the remainder of the study. **Route of Administration:** Oral **Duration of Treatment:** 52 weeks of treatment.

5.2 Concomitant & Prohibited Therapy Permitted: Standard supportive care permitted per protocol. **Prohibited:** Use in combination with other potent immunosuppressants or biological systemic therapies.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Up to 4 weeks	Informed Consent, Medical History, Baseline BSA and PGA assessments
Baseline	Day 0	Randomization, First Dose, Quality of Life Questionnaires
Interim	Week 16	Primary Efficacy Assessment (ss-PGA), Safety Labs, Cross-over to active treatment for placebo group
End of Treatment	Week 52	Final efficacy assessments, exit

evaluations | | **Follow-up** | Week 56 | 4-week post-treatment
safety follow-up |

(Note: Participants complete approximately 15 visits to the study doctor during the 60-week study duration)

7. Outcome Measures (Endpoints)

7.1 Primary Outcome * **Percentage of Participants With a Scalp-specific Physician Global Assessment Score of 0 or 1 (ss-PGA 0/1) at Week 16:** ss-PGA 0/1 response as a percentage of participants with an ss-PGA score of 0 (clear) or 1 (almost clear) with at least a 2-point reduction from baseline at Week 16. Scalp lesions are evaluated in terms of clinical signs of redness, thickness, and scaliness and scored on the following 5-point ss-PGA scale: 0 = absence of disease, 1 = very mild disease, 2 = mild disease, 3 = moderate disease, 4 = severe disease.

7.2 Secondary Outcomes * **Percentage of Participants With a Psoriasis Scalp Severity Index 90 (PSSI 90) at Week 16.** * **Change From Baseline in Scalp-specific Itch Numerical Rating Scale (NRS) Score at Week 16.** * **Percentage of Participants With a Static Physician Global Assessment Score of 0 or 1 (s-PGA 0/1) at Week 16.**

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring of AEs graded 1 to 4 (mild to severe/hospitalization). Common monitored AEs include nasopharyngitis, upper respiratory tract infections, acne, and headache. **Serious Adverse Events (SAEs):** Routine monitoring and rapid reporting, including tracking laboratory abnormalities and vital signs (blood pressure, heart rate). **Data Monitoring Committee (DMC):** Overseen by the sponsor (BMS) for safety signal detection, with safety findings demonstrating a profile consistent with known profiles of TYK2 inhibitors.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) for primary efficacy; Safety Set for all AEs. **Sample Size Justification:** Powered to detect superiority of deucravacitinib versus placebo in achieving ss-PGA 0/1 at Week 16, utilizing 154 randomized subjects (~103 active, ~51 placebo). **Handling of Missing Data:** Defined by standard SAP models (e.g., non-responder imputation [NRI] for binary efficacy endpoints at Week 16).

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Regular onsite and remote monitoring throughout the 60-week study period. **Audit Trail:** eCRF locking and tracking of investigator clinical assessments (ss-PGA, PSSI) to ensure valid data output.

11. Study Identifiers & Governance

Primary ID: IM011-220 **Registry Number:** NCT05478499 **Posting ID:** 331079207309219613 **Trial Status:** COMPLETED **Sponsor/Lead**

Organization: Bristol Myers Squibb **Study Title:** PSORIATYK SCALP **Official Regulatory Title:** A Phase 3B/4, Multicenter, Randomized, Double-blind Placebo-controlled Study to Evaluate the Efficacy and Safety of Deucravacitinib in Participants With Moderate-to-severe Scalp Psoriasis

12. Clinical Framework (Study Specific)

Primary Condition: Moderate-to-Severe Scalp Psoriasis (SNOMED ID: 9014002) **Diagnosis Threshold:** ss-PGA ≥ 3 , PSSI ≥ 12 , SSA $\geq 20\%$, and BSA $\geq 3\%$ **Medical Methodology:** Selective TYK2 Inhibitor Systemic Therapy **Optimization Target:** Deucravacitinib 6 mg QD oral administration

13. Study Procedures & Methodology

Management Pathway: Standard systemic therapy pathway for immune-mediated psoriasis **Education Protocol (HCP):** Protocol-specific site initiation and ss-PGA/PSSI grading calibration **Education Protocol (Patient):** Dosing compliance and patient-reported outcomes (Itch NRS) self-reporting instructions **Quality Audit Mechanism:** Clinical monitoring and centralized tracking of PROs and clinician-reported outcomes

14. Observation & Follow-Up Schedule

Total Duration: 60 Weeks (up to 4-week screening, 52-week treatment, 4-week follow-up) **Onsite Visit Cadence:** Approx. 15 clinic visits spanning screening, Week 16 crossover, and Week 52 EOT **Remote Monitoring Frequency:** Intermittent per protocol **Checklist Review Interval:** Routine safety lab and vital sign reviews aligned with protocol milestones

15. Sign-off & Approval

Role	Name	Signature	Date		:---	:---	:---	:---
Principal Investigator	[Name]	_____	07-Apr-2026					
Clinical Operations Lead	[Name]	_____	07-Apr-2026					Lead
Statistician	[Name]	_____	07-Apr-2026					