

1. Administrative & Study Identification

Full Study Title: Special Investigation of TALZENNA Capsules - Study on BRCA Mutation-Positive Metastatic Castration-Resistant Prostate Cancer **Short Title/Acronym:** A Study to Learn About the Safety of TALZENNA in People With Prostate Cancer **Protocol Number:** Pfizer Internal Study ID **NCT Number:** NCT06733337 **Sponsor Name:** Pfizer Inc. **Investigational Product:** TALZENNA (talazoparib) **Phase of Development:** Observational / Post-Marketing Surveillance (Special Investigation) **Medical Monitor:** Pfizer Medical Monitor

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the safety of TALZENNA in patients with BRCA mutation-positive metastatic castration-resistant prostate cancer over a 12-month observation period. **Secondary Objectives:** To observe treatment patterns, dose modifications, and the real-world incidence of adverse events of special interest.

2.2 Background & Rationale Metastatic castration-resistant prostate cancer (mCRPC) has a poor prognosis, especially when harboring DNA damage repair gene mutations. This study aims to evaluate the safety of TALZENNA in patients with this specific type of prostate cancer, which continues to grow despite low levels of male hormones and has spread to other parts of the body. As a targeted PARP inhibitor therapy, it is crucial to observe the natural progression, safety profile, and outcomes associated with TALZENNA use in routine clinical practice without altering standard treatment.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational **Control Type:** Single-arm cohort **Blinding:** Open-label (No masking) **Randomization:** N/A

3.2 Planned Enrollment & Duration Target \$N\$: Target enrollment based on post-marketing cohort requirements in the participating region **Number of Sites:** Multiple centers (focusing on the Japanese region) **Estimated Study Start:** 14-Jul-2025 **Estimated Primary Completion:** 22-Jun-2029

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Adult male patients with a confirmed diagnosis of BRCA mutation-positive metastatic castration-resistant prostate cancer (mCRPC). 2. Patients for whom the clinical decision to initiate treatment with TALZENNA has been made by the attending physician prior to study enrollment. 3. Willingness to participate and allow data collection from routine medical records.

4.2 Exclusion Criteria 1. Known hypersensitivity to TALZENNA or any of its excipients. 2. Concurrent participation in an interventional clinical trial that would confound the observational real-world data collection. 3. Patients strictly contraindicated for TALZENNA per the locally approved prescribing information.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: TALZENNA capsules, dosed orally according to the locally approved prescribing information and the treating physician's discretion. **Route of Administration:** Oral. **Duration of Treatment:** Observation for up to 52 weeks (12 months) following treatment initiation for each participant.

5.2 Concomitant & Prohibited Therapy Permitted: Standard concomitant medications for mCRPC and comorbidities as prescribed by the Healthcare Provider in a real-world setting. **Prohibited:** Concurrent investigational therapies that interfere with observational safety tracking.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Baseline	Day 0	Informed Consent, Medical History, BRCA Mutation confirmation, Treatment Initiation
Interim	Months 3, 6, 9	Routine Clinical Visits: Safety Assessments, AE Recording, Dose Modification Tracking
End of Observation	Month 12	Final Safety Evaluation, Exit Documentation, Treatment Status

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Incidence and severity of Adverse Events (AEs) and Serious Adverse Events (SAEs) over a 12-month period from the start of TALZENNA treatment. **Measurement Method:** Medical record review and routine clinical laboratory assessments abstracted into the electronic data capture system.

7.2 Secondary & Exploratory Endpoints Incidence of specific adverse events of special interest (e.g., hematological toxicities). Treatment discontinuation or dose reduction rates due to adverse events.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard collection via electronic case report forms (eCRF) during routine real-world clinical visits. **Serious Adverse Events (SAEs):** Reported to the Sponsor within standard pharmacovigilance timelines (typically 24 hours of investigator awareness). **Data Monitoring Committee (DMC):** N/A (Standard safety management for observational post-marketing studies).

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Analysis Set (all enrolled patients who received at least one dose of TALZENNA). **Sample Size Justification:** N/A; driven by real-world prescription events during the planned enrollment window. **Handling of Missing Data:** Missing data will not be imputed; analyses will be based on observed data as is standard for real-world cohort studies.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Pharmacoepidemiology Practices (GPP) and local regulatory requirements for post-marketing surveillance. **Monitoring Plan:** Centralized data monitoring with selective onsite source data verification as dictated by the surveillance protocol. **Audit Trail:** Validated electronic data capture (EDC) system maintaining a full audit trail of patient clinical data entry.

11. Study Identifiers & Governance

Primary ID: Pfizer Internal Protocol ID **Registry Number:** NCT06733337 **Sponsor/Lead Organization:** Pfizer Inc. **Study Title:** A Study to Learn About the Safety of TALZENNA in People With Prostate Cancer **Official Regulatory Title:** Special Investigation of TALZENNA Capsules - Study on BRCA Mutation-Positive Metastatic Castration-Resistant Prostate Cancer

12. Clinical Framework (Prostate Cancer Specific)

Primary Condition: BRCA Mutation-Positive Metastatic Castration-Resistant Prostate Cancer (mCRPC) **Diagnosis Threshold:**

Histologically or cytologically confirmed mCRPC with documented BRCA mutations. **Medical Methodology:** Targeted PARP inhibition in a real-world setting. **Optimization Target:** Real-world safety tracking and treatment pattern observation.

13. Study Procedures & Methodology

Management Pathway: Standard routine clinical pathway for mCRPC. **Education Protocol (HCP):** Protocol training on prospective real-world data collection and pharmacovigilance AE reporting. **Education Protocol (Patient):** Standard clinical counseling on PARP inhibitor usage and potential side effects. **Quality Audit Mechanism:** Centralized review of eCRF completeness and safety event reconciliation.

14. Observation & Follow-Up Schedule

Total Duration: Up to 52 weeks (12 months) of follow-up per patient. **Onsite Visit Cadence:** Dictated by routine clinical care practice. **Remote Monitoring Frequency:** Continuous EDC data review by the Sponsor. **Checklist Review Interval:** Periodically corresponding with standard oncology follow-up visits.

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				