

1. Administrative & Study Identification

Full Study Title: A Study to Learn About the Safety of TALZENNA for the Treatment of Breast Cancer **Short Title/Acronym:** TALZENNA Breast Cancer Safety Study **Protocol Number:** [Pfizer Internal ID / N/A] **NCT Number:** NCT06735742 **Sponsor Name:** Pfizer **Investigational Product:** TALZENNA (talazoparib) **Phase of Development:** Phase N/A (Observational / Post-marketing safety study) **Medical Monitor:** Pfizer Pharmacovigilance/Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the safety of TALZENNA in patients with breast cancer that cannot be removed by surgery and has come back. Specifically, to evaluate the incidence, time of onset, grade, and outcome of specific adverse drug reactions (ADRs). **Secondary Objectives:** To monitor general patient outcomes and track treatment continuation/discontinuation rates over a 52-week (12-month) period.

2.2 Background & Rationale To address the need for real-world safety tracking of TALZENNA (talazoparib) in patients with BRCA mutation-positive and HER2-negative unresectable or recurrent breast cancer. The study aims to gather robust safety data on the incidence of critical adverse events (such as myelosuppression and interstitial lung disease) in patients who have a history of prior cancer chemotherapy and are receiving TALZENNA as a standard of care.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational **Control Type:** Single-arm **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target \$N\$: [Observational Cohort Size - dependent on real-world prescriptions] **Number of Sites:** 3 (Japan) **Estimated Study Start:** 31-Jan-2025 **Estimated Primary Completion:** 25-Jul-2031

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Patients with BRCA mutation-positive and HER2-negative unresectable or recurrent breast cancer. 2. History of prior cancer chemotherapy. 3. No history of previous use of TALZENNA (talazoparib).

4.2 Exclusion Criteria 1. Prior treatment with TALZENNA (talazoparib). 2. Breast cancer that is amenable to curative surgery. 3. HER2-positive breast cancer or lack of confirmed BRCA mutation.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: TALZENNA administered orally according to the local prescribing information and physician's clinical judgment. **Route of Administration:** Oral **Duration of Treatment:** Patients will be followed up to 52 weeks (12 months) from the start of treatment.

5.2 Concomitant & Prohibited Therapy Permitted: Standard concomitant medications for symptom management and co-morbidities as per standard of care. **Prohibited:** Concomitant use of other investigational anti-cancer agents or PARP inhibitors.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
1	Day -14 to 0	Informed Consent, Medical History, BRCA/HER2 verification
2	Day 0	First Dose of TALZENNA, Baseline Safety Labs
3	Weeks 12, 24, 36, 48	Safety Labs, ADR Assessment (Myelosuppression, ILD, Thromboembolism tracking)
4	Week 52	Final Safety Review, Discontinuation Status, ADR outcome tabulation

(Note: Participants completing or stopping treatment earlier than 52 weeks will be tracked until the date of completion or discontinuation)

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Incidence, time of onset, grade, and outcome of targeted adverse drug reactions (ADRs), specifically: myelosuppression, interstitial lung disease (ILD), thromboembolism, and secondary malignant tumour. **Measurement Method:** Clinical evaluation, laboratory monitoring, and pharmacovigilance tabulation over the 52-week follow-up.

7.2 Secondary & Exploratory Endpoints - Overall safety and tolerability profile of TALZENNA up to 12 months. - Rate of treatment discontinuation due to adverse events.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Method of collection via routine clinical care visits and patient reporting. Special focus on ILD

and myelosuppression. **Serious Adverse Events (SAEs):** Reported within 24 hours of site awareness per standard pharmacovigilance protocols. **Data Monitoring Committee (DMC):** Internal safety monitoring mechanisms managed by Pfizer.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (All patients receiving at least one dose of TALZENNA). **Sample Size Justification:** Descriptive safety study; powered observationally based on the real-world adoption of TALZENNA in the target demographic. **Handling of Missing Data:** Last Observation Carried Forward (LOCF) or descriptive tabulation without imputation for safety events.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and local regulatory guidelines (e.g., Japan's special drug use investigation standards). **Monitoring Plan:** Standard pharmacovigilance safety data monitoring and site data verification. **Audit Trail:** eCRF locking and query resolution via standard Pfizer clinical database systems.

11. Study Identifiers & Governance

Primary ID: N/A (Observational Registry) **Registry Number:** NCT06735742 **Sponsor/Lead Organization:** Pfizer **Study Title:** A Study to Learn About the Safety of TALZENNA for the Treatment of Breast Cancer **Official Regulatory Title:** A Study to Learn About the Safety of TALZENNA for the Treatment of Breast Cancer

12. Clinical Framework (Breast Cancer Specific)

Primary Condition: Breast Cancer (Unresectable or recurrent) **Diagnosis Threshold:** BRCA mutation-positive and HER2-negative **Medical Methodology:** PARP Inhibitor Therapy **Optimization Target:** Drug safety and toxicity management (Myelosuppression / ILD)

13. Study Procedures & Methodology

Management Pathway: Standard clinical pathway for Metastatic Breast Cancer **Education Protocol (HCP):** Safety reporting training for targeted ADRs **Education Protocol (Patient):** Counseling on signs of myelosuppression and interstitial lung disease **Quality Audit Mechanism:** Ongoing pharmacovigilance tracking and AE reconciliation

14. Observation & Follow-Up Schedule

Total Duration: 52 weeks (12 months) **Onsite Visit Cadence:** Standard clinical oncology intervals (e.g., Weeks 4, 12, 24, 36, 48, 52) **Remote Monitoring Frequency:** Continuous pharmacovigilance reporting **Checklist Review Interval:** Every visit for targeted AEs

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				