

1. Administrative & Study Identification

Full Study Title: A Study to Learn About Ritlecitinib for the Potential Treatment of Chronic Spontaneous Urticaria in Adults
Short Title/Acronym: Ritlecitinib in CSU **Protocol Number:** PF-06651600 **NCT Number:** NCT07219615 **Sponsor Name:** Pfizer
Investigational Product: Ritlecitinib (Litfulo / PF-06651600)
Phase of Development: Phase 2 (Phase 2b) **Trial Status:** Recruiting
Medical Monitor: Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the efficacy of ritlecitinib in reducing the Urticaria Activity Score 7 (UAS7) at Week 12, and to assess safety through the incidence of Treatment Emergent Adverse Events, Serious Adverse Events, and adverse events leading to discontinuation. **Secondary Objectives:** To evaluate the changes from baseline in UAS7 at all other timepoints, Itch Severity Score 7 (ISS7) at Week 12, and Hives Severity Score 7 (HSS7) at Week 12.

2.2 Background & Rationale To address the medical necessity of providing effective disease control for adult patients with CSU inadequately controlled by second-generation H1-antihistamines (sgAH). CSU is an inflammatory skin disease characterized by recurrent, unpredictable, and itchy hives and angioedema. Ritlecitinib, an oral Janus kinase 3 (JAK3) and TEC family kinase inhibitor, offers a targeted approach to address the underlying inflammatory pathways driving the overactive immune response.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Placebo-controlled **Blinding:** Quadruple-blind (Participant, Care Provider, Investigator, Outcomes Assessor) **Randomization:** Randomized, Parallel Assignment

3.2 Planned Enrollment & Duration Target \$N\$: 200 participants
Number of Sites: Global multicenter trial **Estimated Study Start:** 30-Oct-2025 **Estimated Primary Completion:** 25-Dec-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Age ≥ 18 years. 2. Confirmed diagnosis of chronic spontaneous urticaria (CSU) for ≥ 3

months at the time of screening. 3. CSU inadequately controlled by second-generation antihistamine (sgAH) defined as the presence of itch and hives for ≥ 6 consecutive weeks prior to screening, with an Urticaria Activity Score 7 (UAS7) ≥ 16 and an Itch Severity Score 7 (ISS7) ≥ 8 during the 7 days prior to randomization.

4.2 Exclusion Criteria 1. Urticaria solely due to inducible urticaria. 2. Active dermatologic diseases (or conditions) other than chronic urticaria with urticaria wheals or angioedema symptoms (e.g., urticarial vasculitis, erythema multiforme, cutaneous mastocytosis). 3. Use of prohibited alternative CSU treatments; patients must be willing to stop all other treatments for CSU aside from the permitted sgAH.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Ritlecitinib 50 mg, 100 mg, or matching placebo taken orally once daily. **Route of Administration:** Oral **Duration of Treatment:** 24 weeks total. Divided into Period A (12 weeks of 50 mg, 100 mg, or placebo) and Period B (12 weeks where participants on 50 mg and 100 mg continue their doses, while the placebo group is switched to 100 mg).

5.2 Concomitant & Prohibited Therapy Permitted: Second-generation antihistamine (sgAH) at locally approved doses or higher. **Prohibited:** All other systemic therapies for CSU (e.g., omalizumab, cyclosporine, systemic corticosteroids outside allowed windows).

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, Baseline UAS7/ISS7/HSS7 Assessment, ECG, Blood/Hearing tests
Baseline	Day 1	Randomization, First Dose, Electronic Diary (eDiary) initialization
Interim 1	Week 12	End of Period A: Primary Efficacy Assessment, Safety Labs, Treatment Switch for Placebo Group
End of Study	Week 24	Final Efficacy & Safety Evaluation, Exit Interview, IP Accountability (followed by a 4-week off-treatment follow-up)

7. Outcome Measures (Endpoints)

7.1 Primary Endpoints 1. Change from baseline in Urticaria Activity Score 7 (UAS7) at Week 12. 2. Incidence of Treatment Emergent Adverse Events, Serious Adverse Events, and Adverse Events leading to discontinuation.

7.2 Secondary & Exploratory Endpoints 1. Change from baseline in UAS7 at all timepoints except Week 12. 2. Change from baseline in Itch Severity Score 7 (ISS7) at Week 12. 3. Change from baseline in Hives Severity Score 7 (HSS7) at Week 12.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard collection via physical exams, laboratory tests, hearing tests, and patient interviews during clinic visits up to Week 24 and the post-treatment follow-up. **Serious Adverse Events (SAEs):** Evaluated continuously; must be reported within 24 hours of site awareness. **Data Monitoring Committee (DMC):** Internal safety monitoring mechanisms managed by Pfizer.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) population for primary efficacy outcomes; Safety Set for all AE/SAE evaluations. **Sample Size Justification:** $N=200$ powered appropriately ($\alpha = 0.05$) to detect statistically meaningful reductions in UAS7, ISS7, and HSS7 scores between the ritlecitinib active arms and placebo. **Handling of Missing Data:** Mixed Model for Repeated Measures (MMRM) or Multiple Imputation applied for missing continuous eDiary endpoints.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Routine onsite and remote monitoring visits to ensure clinical pathway adherence. **Audit Trail:** eCRF locking, automated audit trails for the patient eDiaries, and data clarification forms via standard quality audit mechanisms.

11. Study Identifiers & Governance

Primary ID: {{PF-06651600}} **Posting ID:** {{388920154108427755}}
Registry Number: {{NCT07219615}} **Sponsor/Lead Organization:** {{Pfizer}} **Study Title:** {{A Study to Learn About Ritlecitinib for the Potential Treatment of Chronic Spontaneous Urticaria in Adults}} **Official Regulatory Title:** {{A PHASE 2B RANDOMIZED, DOUBLE-BLIND, 12-WEEK PLACEBO-CONTROLLED STUDY WITH A 12-WEEK DOUBLE-BLIND EXTENSION PERIOD TO EVALUATE THE EFFICACY, SAFETY AND TOLERABILITY OF RITLECITINIB (PF-06651600) IN ADULT PARTICIPANTS WITH CHRONIC SPONTANEOUS URTICARIA}}

12. Clinical Framework (Chronic Spontaneous Urticaria Specific)

Primary Condition: {{Chronic Spontaneous Urticaria (CSU)}}
Diagnosis Threshold: {{UAS7 >= 16 and ISS7 >= 8 with >= 3 months history}}
Medical Methodology: {{JAK3/TEC family kinase inhibition}}
Optimization Target: {{Symptom reduction and disease control (UAS7 <= 6)}}

13. Study Procedures & Methodology

Management Pathway: {{Standard CSU assessment with daily eDiary tracking}}
Education Protocol (HCP): {{Site initiation, study protocol, and AE reporting training}}
Education Protocol (Patient): {{Instruction on accurate daily eDiary completion for UAS7/ISS7/HSS7}}
Quality Audit Mechanism: {{Regular eDiary compliance reviews and clinical site monitoring}}

14. Observation & Follow-Up Schedule

Total Duration: {{28 weeks (24 weeks treatment + 4 weeks follow-up)}}
Onsite Visit Cadence: {{Up to 9 in-person visits throughout the screening, treatment, and follow-up periods}}
Remote Monitoring Frequency: {{Daily eDiary entry for hives and itch severity}}
Checklist Review Interval: {{At every scheduled clinic visit}}

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				