

1. Administrative & Study Identification

**Full Study Title** A Multicenter, Randomized, Double-blind, Placebo-controlled Phase 3 Study to Evaluate the Efficacy, Safety, and Pharmacokinetics of Deucravacitinib in Adolescent Participants (12 Years to Less Than 18 Years) With Moderate to Severe Plaque Psoriasis **Brief Study Title** A Study to Evaluate the Efficacy, Safety, and Drug Levels of Deucravacitinib (BMS-986165) in Adolescent Participants With Moderate to Severe Plaque Psoriasis **Short Title/Acronym** POETYK-PsO-Ped **Posting ID** 397439511254758128 **Protocol Number** IM011-1128 **NCT Number** NCT06979453 **Trial Status** Recruiting **Sponsor Name / Company** Bristol Myers Squibb **Investigational Product (Drug Name)** Deucravacitinib (BMS-986165) **Trade Name** Sotyktu **ATC Code** L04AA56 **EMA URL** [https://www.ema.europa.eu/en/documents/product-information/sotyktu-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/sotyktu-epar-product-information_en.pdf) **Phase of Development** Phase 3 **Medical Monitor** Bristol Myers Squibb Clinical Operations Lead

---

2. Study Synopsis & Rationale

**2.1 Study Objective** **Primary Objective:** To evaluate the efficacy of deucravacitinib compared to placebo in adolescent participants (12 to <18 years) with moderate to severe plaque psoriasis. **Secondary Objectives:** To evaluate the safety, tolerability, pharmacokinetics, and long-term efficacy (up to 5 years) of deucravacitinib in the study population.

**2.2 Background & Rationale** To address the medical necessity for a well-tolerated, oral targeted therapy (TYK2 inhibitor) for adolescent patients with moderate to severe plaque psoriasis. This study aims to provide a safe and effective oral alternative to biologic injectables and traditional immunosuppressants, assessing its ability to achieve clear or almost clear skin over a long-term treatment period.

---

3. Study Design & Methodology

**3.1 Design Framework** **Study Type:** Interventional **Control Type:** Placebo-controlled **Blinding:** Double-blind **Randomization:** Randomized

**3.2 Planned Enrollment & Duration** **Targeted Enrollment:** 366 **Number of Sites:** Multicenter (Global) **Estimated Study Start:** 03-Dec-2025 **Estimated Primary Completion:** 12-Oct-2029

---

## 4. Subject Selection (Eligibility Criteria)

**4.1 Inclusion Criteria** 1. Adolescent participants 12 to < 18 years of age. 2. Stable plaque psoriasis for 6 months or more prior to Screening. 3. Moderate to severe psoriasis defined by: Psoriasis Area and Severity Index (PASI)  $\geq$  12, Static Physician's Global Assessment (sPGA)  $\geq$  3, and Body Surface Area (BSA)  $\geq$  10% involvement at screening visit and Day 1. 4. A female (as assigned at birth) participant is eligible to participate if she is not pregnant or breastfeeding, and at least 1 of the following conditions applies: Is not an individual of childbearing potential (IOCBP), or is an IOCBP and using a contraceptive method that is highly effective (with a failure rate of < 1% per year) during the study intervention period and for at least 3 days after discontinuation of the study intervention and agrees not to donate eggs for the purpose of reproduction for the same period.

**4.2 Exclusion Criteria** 1. Participants weighing < 30.0 kg at Screening and Day 1. 2. Non-plaque psoriasis (e.g., guttate, inverse, pustular, erythrodermic, or drug-induced) or any skin condition other than plaque psoriasis that could interfere with assessments. 3. History of serious bacterial, fungal, or viral infection requiring hospitalization and intravenous (IV) antimicrobial treatment within 60 days prior to Day 1, or receipt of live vaccines/BCG within 60 days prior to Day 1, or plans to receive a live vaccine during the study, or within 60 days after completing study intervention. 4. Any untreated bacterial infection within 60 days prior to Day 1, or any ongoing evidence of chronic bacterial infection (e.g., chronic pyelonephritis, chronic osteomyelitis, chronic bronchiectasis). 5. Herpes simplex/zoster, active tuberculosis (TB), hepatitis C virus (HCV), hepatitis B virus (HBV), or human immunodeficiency virus (HIV) infection. 6. Any major illness/condition or evidence of an unstable clinical condition (e.g., renal, hepatic, hematologic, gastrointestinal, endocrine, pulmonary, psychiatric, neurologic, immunologic, or local active infection/infectious illness) that will substantially increase risk. 7. Cancer or history of cancer (solid organ or hematologic including myelodysplastic syndrome) or lymphoproliferative disease within the previous 5 years. 8. Prior exposure to deucravacitinib, or receipt of any medication that is specifically prohibited.

---

## 5. Intervention & Treatment Plan

**5.1 Investigational Regimen Dosage/Frequency:** Deucravacitinib (investigational doses) or Placebo, administered daily. **Route of Administration:** Oral **Duration of Treatment:** Up to 5 years (including long-term efficacy and safety follow-up).

**5.2 Concomitant & Prohibited Therapy Permitted:** Standard non-medicated emollients. **Prohibited:** Prior exposure to

deucravacitinib; live vaccines or BCG during the study and within 60 days after completing the study intervention.

---

## 6. Study Timeline & Visit Schedule

| Visit ID     | Window (Days)   | Key Activities/Procedures                                                    |
|--------------|-----------------|------------------------------------------------------------------------------|
| Screening    | Day -14 to 0    | Informed Consent/<br>Assent, Medical History, PASI/sPGA/BSA baseline scoring |
| Baseline     | Day 0           | Randomization, First Dose, Safety Labs                                       |
| Interim 1    | Week 16 $\pm$ 7 | Primary Efficacy Assessment (PASI/sPGA), Safety evaluations                  |
| End of Study | Up to Year 5    | Final Vital Signs, Exit Interview, IP Accountability, Long-term safety       |

---

## 7. Outcome Measures (Endpoints)

**7.1 Primary Endpoints** 1. Number of participants achieving at least 75% improvement in Psoriasis Area and Severity Index (PASI 75). 2. Number of participants achieving a static Physician's Global Assessment (sPGA) score of 0 (clear) or 1 (almost clear) with at least a 2-point reduction from baseline.

**7.2 Secondary & Exploratory Endpoints** 1. Number of participants achieving at least 90% improvement in PASI (PASI 90). 2. Change from baseline in PASI. 3. Change from baseline in body surface area (BSA) involvement. 4. Long-term safety profile and maintenance of clinical response (up to 5 years).

---

## 8. Safety & Adverse Event Reporting

**Adverse Event (AE) Monitoring:** Investigator assessments, laboratory evaluations, and patient reporting at each visit.

**Serious Adverse Events (SAEs):** To be reported within 24 hours of site awareness. **Data Monitoring Committee (DMC):** External independent Data Monitoring Committee for safety oversight.

---

## 9. Statistical Analysis Plan (SAP) Summary

**Analysis Populations:** Intent-to-Treat (ITT) for efficacy, Safety Set for adverse events. **Sample Size Justification:**  $N=366$  to provide sufficient power ( $\alpha = 0.05$ ,  $Power = 80\%$ ) to detect superiority of deucravacitinib over placebo. **Handling of Missing Data:** Non-Responder Imputation (NRI) for binary primary efficacy endpoints.

---

## 10. Quality Assurance & Regulatory Compliance

**GCP Compliance:** This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

**Monitoring Plan:** Routine onsite and remote monitoring visits by the sponsor. **Audit Trail:** Validated electronic data capture (EDC) system with full audit trails and eCRF locking.

---

## 11. Study Identifiers & Governance

**Primary ID:** IM011-1128 **Registry Number:** NCT06979453 **Sponsor/Lead Organization:** Bristol Myers Squibb **Study Title:** POETYK-PsO-Ped **Official Regulatory Title:** A Multicenter, Randomized, Double-blind, Placebo-controlled Phase 3 Study to Evaluate the Efficacy, Safety, and Pharmacokinetics of Deucravacitinib in Adolescent Participants (12 Years to Less Than 18 Years) With Moderate to Severe Plaque Psoriasis

---

## 12. Clinical Framework (Psoriasis Specific)

**Primary Condition / Disease Area:** Moderate to Severe Plaque Psoriasis

**Preferred UMLS Name:** Plaque Psoriasis **Diagnosis**

**Threshold:** PASI  $\geq$  12, sPGA  $\geq$  3, and BSA  $\geq$  10% **Medical**

**Methodology:** Oral Targeted Therapy (TYK2 Inhibitor) **Optimization**

**Target:** PASI 75 and sPGA 0/1

---

## 13. Study Procedures & Methodology

**Management Pathway:** Plaque Psoriasis systemic treatment pathway

**Education Protocol (HCP):** Site initiation visits, protocol training, and PASI/sPGA standardized grading training

**Education Protocol (Patient):** Informed assent (adolescent) and consent (parent/guardian), dosing diary training **Quality Audit Mechanism:** Centralized data monitoring and independent clinical site audits

---

## 14. Observation & Follow-Up Schedule

**Total Duration:** 5 years (long-term follow-up) **Onsite Visit**

**Cadence:** Screening, Baseline, Week 16, and scheduled long-term maintenance visits **Remote Monitoring Frequency:** Interim safety

checks as defined by the protocol **Checklist Review Interval:** Every clinical visit

---

## 15. Sign-off & Approval

|                        |        |           |               |     |     |     |     |
|------------------------|--------|-----------|---------------|-----|-----|-----|-----|
| Role                   | Name   | Signature | Date          | :-- | :-- | :-- | :-- |
| Principal Investigator | [Name] | ____      | [DD-MMM-YYYY] |     |     |     |     |

Clinical Operations Lead | [Name] | \_\_\_\_ | [DD-MMM-YYYY] | | Lead  
Statistician | [Name] | \_\_\_\_\_ | [DD-MMM-YYYY] |