

1. Administrative & Study Identification

Study Title: Real World Evidence in China: Faricimab Use in Diabetic Macular Edema, Retinal Vein Occlusion, and Neovascular Age-Related Macular Degeneration (The Farseeing Study) **Official Title:** China Faricimab Real World Evidence: Evaluation of Faricimab Effectiveness, Safety and Treatment Pattern, in Diabetic Macular Edema, Retinal Vein Occlusion and Neovascular Age-Related Macular Degeneration: The Farseeing Study **Short Title/Acronym:** The Farseeing Study **Protocol Number:** ML45401 **NCT Number:** NCT06439576 **Posting ID:** 415886411132371914 **Sponsor Name:** Roche (Hoffmann-La Roche / Shanghai Roche Pharmaceutical Co., Ltd.) **Investigational Product:** Faricimab (VABYSMO®) | **ATC Code:** S01LA09 | **EMA URL:** [Product Information](#) **Phase of Development:** Not Applicable (Observational / Real World Evidence) **Medical Monitor:** Hoffmann-La Roche Clinical Operations (Reference Study ID: ML45401)

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the changes in visual acuity (converted to ETDRS letters) from baseline after 12 months of treatment with faricimab in individuals with Diabetic Macular Edema (DME), Retinal Vein Occlusion (RVO), and Neovascular Age-Related Macular Degeneration (nAMD). **Secondary Objectives:** To explore long-term effectiveness, safety, and treatment patterns among patients receiving faricimab in real-world, routine clinical practice in China.

2.2 Background & Rationale To address the need for real-world, long-term data to gain clinical evidence on faricimab by observing patient cohorts in routine clinical practice. The data collected will contribute to a better understanding of the long-term safety and effectiveness of faricimab in the Chinese population, potentially benefiting future patients and optimizing real-world treatment strategies for serious retinal conditions.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational (Prospective and Retrospective, Primary data collection) **Control Type:** Single-arm (Real-world cohort) **Blinding:** Open-label / Non-blinded **Randomization:** N/A

3.2 Planned Enrollment, Duration & Status Target **\$N\$:** 1,000 evaluable patients **Number of Sites:** Multi-center across China

Study Status: Active but not currently recruiting
(ACTIVE_NOT_RECRUITING) **Estimated Primary Completion:** ~12 months post-enrollment

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Have signed the informed consent. 2. Female and male Chinese patients, who had been diagnosed with nAMD, DME, or RVO by CFP, OCT, FFA, ICGA, or OCTA. 3. ≥50 years old for patients with nAMD, ≥18 years old both for patients with DME and RVO, at the time of signing informed consent (or the first administration of faricimab, whichever occurs first). 4. Patients for whom the decision to receive treatment with faricimab is made prior to and independent from study participation. 5. Patients have received at least one faricimab treatment (the first dose) in the study eye.

4.2 Exclusion Criteria 1. Patients not receiving treatment for nAMD/DME/RVO with faricimab according to the standard of care and in line with the current summary of product characteristics (SPC) / labeling in China. 2. Active ocular inflammation or suspected/active ocular infection in either eye. 3. Received any other anti-VEGF treatment after faricimab. 4. Received any steroid treatment within 6 months (180 days) before the first faricimab treatment in the study eye. 5. Any participation in any other clinical trials currently. 6. Patients could not provide the clinical data (visual acuity and OCT images) within 2 weeks (14 days) before receiving the initial faricimab injection.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Intravitreal injections of faricimab administered according to standard clinical practice and local guidelines. The interval between maintenance injections is personalized based on patient response and visual acuity metrics. **Route of Administration:** Intravitreal Injection **Duration of Treatment:** At least 12 months for primary endpoint observation; long-term follow-up as per standard of care.

5.2 Concomitant & Prohibited Therapy Permitted: Standard concomitant therapies for underlying conditions (e.g., diabetes management, hypertension). **Prohibited:** Other anti-VEGF therapies after faricimab initiation, and steroid treatments within 6 months prior to the first faricimab dose.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Screening	Day -14 to 0 Diagnosis

confirmation (nAMD/DME/RVO), prescription of faricimab, visual acuity and OCT imaging, informed consent for real-world data collection. | | **Baseline** | Day 0 | First faricimab intravitreal injection, baseline visual acuity assessment. | | **Interim** | Real-world intervals | Routine clinical visits (e.g., Q4W, Q8W) tailored to individual response: visual acuity checks, continued injections. | | **End of Study** | Month 12 | Final visual acuity measurement (ETDRS), safety profile review, treatment pattern extraction. |

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint 1. Change in Visual Acuity from Baseline at Month 12: Visual acuity will be collected as measured per local practice and then will be converted to the approximate ETDRS letter score.

7.2 Secondary Outcomes 1. Visual Acuity Over Time. 2. Change in Visual Acuity from Baseline Over Time. 3. Central Subfield Thickness Over Time.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Extracted from routine clinical practice records; focused on incidence of intraocular inflammation, intraocular pressure spikes, and other AEs. **Serious Adverse Events (SAEs):** Standard safety reporting to the Sponsor and regulatory bodies within typical real-world evidence frameworks. **Data Monitoring Committee (DMC):** Internal safety governance teams managed by Roche.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Real-world observational cohort (all enrolled patients fulfilling eligibility and receiving at least one dose of faricimab). **Sample Size Justification:** Target $N=1,000$ to adequately capture real-world treatment patterns and power sub-group analyses across the three indications (nAMD, DME, RVO) within the Chinese population. **Handling of Missing Data:** Standard epidemiological imputation mechanisms for observational, real-world data tracking.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Pharmacoepidemiology Practices (GPP) and local regulations concerning non-interventional, real-world evidence collection. **Monitoring Plan:** Data abstraction and validation from

participating centers' electronic medical records. **Audit Trail:** Standardized electronic data capture (EDC) systems used to maintain a secure audit trail for all abstracted patient data.

11. Study Identifiers & Governance

Primary ID: {{ML45401}} **Registry Number:** {{NCT06439576}} **Posting ID:** {{415886411132371914}} **Sponsor/Lead Organization:** {{Roche}} **Study Title:** {{Real World Evidence in China: Faricimab Use in Diabetic Macular Edema, Retinal Vein Occlusion, and Neovascular Age-Related Macular Degeneration (The Farseeing Study)}} **Official Regulatory Title:** {{China Faricimab Real World Evidence: Evaluation of Faricimab Effectiveness, Safety and Treatment Pattern, in Diabetic Macular Edema, Retinal Vein Occlusion and Neovascular Age-Related Macular Degeneration: The Farseeing Study}}

12. Clinical Framework (Ophthalmology Specific)

Primary Condition: {{Macular Degeneration, Diabetic Macular Edema, Neovascular Age-related Macular Degeneration, Retinal Vein Occlusion}} **Preferred UMLS Name:** {{Retinal Vascular Diseases}} **Diagnosis Threshold:** {{Clinical diagnosis requiring anti-VEGF therapy, confirmed by CFP, OCT, FFA, ICGA, or OCTA}} **Medical Methodology:** {{Routine Real-World Clinical Practice / Standard of Care}} **Optimization Target:** {{Improvement and maintenance of Visual Acuity (ETDRS letters) and Central Subfield Thickness}}

13. Study Procedures & Methodology

Management Pathway: {{Standard clinical pathway for intravitreal anti-VEGF therapy}} **Education Protocol (HCP):** {{Routine local product labeling (SPC) compliance}} **Education Protocol (Patient):** {{Standard of care counseling for intravitreal injections}} **Quality Audit Mechanism:** {{Real-World Data (RWD) extraction and source-data verification}}

14. Observation & Follow-Up Schedule

Total Duration: {{At least 12 months follow-up per patient}} **Onsite Visit Cadence:** {{Determined by routine clinical practice / Personalized Treatment Interval}} **Remote Monitoring Frequency:** {{N/A - Non-interventional observational study}} **Checklist Review Interval:** {{Periodic electronic medical record abstraction}}

15. Sign-off & Approval

	Role		Name		Signature		Date			:---		:---		:---		:---	
	Principal Investigator		[Name]		_____		02-Apr-2026										
	Clinical Operations Lead		[Name]		_____		02-Apr-2026										
	Statistician		[Name]		_____		02-Apr-2026										