

1. Administrative & Study Identification

Full Study Title: A Single Arm, Open Label Multicentre Extension Study To Evaluate The Effectiveness And Safety Of Ocrelizumab In Patients With Multiple Sclerosis Previously Enrolled In A F. Hoffmann-La Roche Sponsored Ocrelizumab Phase IIb/IV Clinical Trials **Brief Title:** This is an Extension Study of the Roche P-trials to Investigate Safety and Effectiveness of Ocrelizumab in Participants With Multiple Sclerosis (MS) **Short Title/Acronym:** LIBERTO Study **Posting ID:** 504857113401145913 **Protocol Number:** MN39158 (Other IDs: 2023-506543-41-00, 2017-004886-29) **NCT Number:** NCT03599245 **Sponsor/Company:** F. Hoffmann-La Roche Ltd (Roche) **Investigational Product:** Ocrelizumab (Trade Name: Ocrevus; ATC Code: L04AA36; Anti-CD20 Monoclonal Antibody) **Phase of Development:** Phase 3 **Medical Monitor:** F. Hoffmann-La Roche Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the long-term effectiveness and safety of ocrelizumab in participants with multiple sclerosis (MS) who previously completed the treatment period in F. Hoffmann-La Roche sponsored Phase IIb/IV parent trials (e.g., CASTING, ENSEMBLE). **Secondary Objectives:** To evaluate No Evidence of Disease Activity (NEDA), which includes the absence of protocol-defined relapses, confirmed disability progression, and absence of new/enlarging T2-weighted or gadolinium-enhancing T1 lesions on MRI. Additional objectives include tracking annualized relapse rates and adverse events (AEs).

2.2 Background & Rationale Multiple sclerosis is a chronic autoimmune disease of the central nervous system featuring demyelination and axonal degeneration. While ocrelizumab has demonstrated a favorable benefit-risk ratio in parent trials, there remains a medical necessity to evaluate the long-term durability of its effectiveness and its long-term safety profile. This extension study offers patients continued disease-modifying therapy to reduce the frequency of relapses, prevent new brain lesions, and delay disability accumulation.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional (Extension Study) **Control Type:** Single-arm **Blinding:** Open-label **Randomization:** N/A (Rollover Extension)

3.2 Planned Enrollment & Duration Target Enrollment: 1055 participants (Rollover from parent trials) **Number of Sites:** Global trial across 25 countries **Estimated Study Start:** 2018 **Trial Status:** Completed

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Signed Informed Consent Form. 2. Able to comply with the study protocol, in the investigator's judgment. 3. Completed the treatment period of Roche sponsored ocrelizumab P-trials. 4. Age ≥ 18 to ≤ 65 years with a confirmed diagnosis of Relapsing-Remitting Multiple Sclerosis (RRMS) or Primary Progressive Multiple Sclerosis (PPMS). 5. For women of childbearing potential: agreement to use a reliable method of birth control during treatment and for at least 6 months after the last dose.

4.2 Exclusion Criteria 1. Hypersensitivity to ocrelizumab or any of its excipients. 2. Participants in a severely immunocompromised state until the condition resolves. 3. Evidence of any adverse event potentially attributable to ocrelizumab, for which the local label recommends permanent discontinuation. 4. Existence of a contra-indication as per SmPC. 5. Prohibited concomitant medication as specified in protocol. 6. Participants intending to become pregnant during the study or within 6 months after the last dose of the study drug in the parent study.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Ocrelizumab 600 mg every 24 weeks (6 months). **Route of Administration:** Intravenous (IV) Infusion. **Duration of Treatment:** Up to 144 weeks (approximately 2 to 2.5 years total including follow-up).

5.2 Concomitant & Prohibited Therapy Permitted: Standard symptomatic therapies for MS symptoms as prescribed by the treating physician. **Prohibited:** Concomitant use of other disease-modifying therapies (DMTs) or treatments that strongly suppress the immune system.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening/Rollover	Day -14 to 0	Informed Consent, Eligibility Confirmation, MRI Re-baselining
Baseline	Day 0	Continued IV Infusion of Ocrelizumab, Vital Signs, EDSS Assessment
Interim	Every 24 Weeks	Safety Labs, Ocrelizumab Infusion, Clinical Effectiveness Assessment (Relapses, EDSS), Annual MRI
End of Study	Follow-up	Final

Safety Checks, Exit Interview, Evaluation of Disability Progression (1-2 visits up to 1 year post-last dose) |

7. Outcome Measures (Endpoints)

7.1 Primary Endpoints 1. Time to onset of Confirmed Disability Progression (CDP) sustained for at least 24 weeks and for at least 48 weeks. 2. Percentage of participants who have confirmed disability improvement (CDI), CDP for at least 24 weeks and for at least 48 weeks yearly and over the duration of the treatment. 3. Percentage of participants who have improved, stable or worsened disability compared with baseline: Improved, stable or worsened disability is measured by Expanded Disability Status Scale (EDSS) (annually/by epoch and over duration of the study). Stable EDSS is defined as EDSS change +/- 0.5. Worsening is > 0.5 increase of EDSS, Improvement is > 0.5 decrease of EDSS.

7.2 Secondary Endpoints 1. Time to first protocol-defined event of disease activity. 2. Time to first relapse. 3. Annualized relapse rate.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous evaluation of infusion-related reactions, infection rates, and changes in concurrent medications. Participants are monitored closely during and after infusions. **Serious Adverse Events (SAEs):** Safety assessments performed regularly; standard regulatory reporting timelines (e.g., 24 hours) apply for SAEs. **Data Monitoring Committee (DMC):** Internal safety tracking conducted by F. Hoffmann-La Roche Ltd.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all patients receiving at least one dose of study medication in the extension) and Effectiveness Set (rollover patients with available post-baseline efficacy data). **Sample Size Justification:** Target defined dynamically by the number of eligible patients successfully completing parent trials (CASTING, ENSEMBLE, etc.) opting for extension. **Handling of Missing Data:** Standard survival analysis algorithms (e.g., Kaplan-Meier) for time-to-event endpoints (like confirmed disability progression).

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Routine onsite and remote monitoring per

Sponsor's standard extension protocol guidelines to ensure safety and protocol adherence. **Audit Trail:** Detailed electronic Case Report Form (eCRF) entries maintaining linkage to the parent trial history.

11. Study Identifiers & Governance

Primary ID: MN39158 **Registry Number:** NCT03599245 **Sponsor/Lead Organization:** F. Hoffmann-La Roche Ltd **Study Title:** LIBERTO Study **Official Regulatory Title:** A Single Arm, Open Label Multicentre Extension Study To Evaluate The Effectiveness And Safety Of Ocrelizumab In Patients With Multiple Sclerosis Previously Enrolled In A F. Hoffmann-La Roche Sponsored Ocrelizumab Phase IIIB/IV Clinical Trials

12. Clinical Framework (Multiple Sclerosis Specific)

Primary Condition / Disease Area: Multiple Sclerosis (Relapsing-Remitting / Primary Progressive) **Diagnosis Threshold:** Clinically Definite MS; Rollover criteria established by successful completion of parent trial **Medical Methodology:** Anti-CD20 B-cell depletion therapy **Optimization Target:** Maintenance of NEDA (No Evidence of Disease Activity) and suppression of long-term disability accumulation

13. Study Procedures & Methodology

Management Pathway: Long-term open-label extension therapeutic pathway **Education Protocol (HCP):** Standard MS infusion protocol and safety-monitoring training **Education Protocol (Patient):** Education on infusion reactions, reporting new neurological symptoms (relapses), and contraception requirements **Quality Audit Mechanism:** Ongoing tracking of EDSS scores, MRI outputs, and comprehensive AE reporting

14. Observation & Follow-Up Schedule

Total Duration: Approximately 144 weeks (up to 2.5 years including safety follow-up) **Onsite Visit Cadence:** Every 24 weeks (coinciding with infusion schedule) **Remote Monitoring Frequency:** Ongoing safety check-ins as clinically indicated **Checklist Review Interval:** Comprehensive review every 24 weeks prior to each consecutive ocrelizumab infusion

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---	
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]					
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]			Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]					