

1. Administrative & Study Identification

Full Study Title: Effectiveness of Treatment With Tofacitinib in Patients With Psoriatic Arthritis in Routine Clinical Practice
Short Title/Acronym: TOPSATI
Protocol Number: A3921332
NCT Number: NCT04517669
Sponsor Name: Pfizer
Investigational Product: Tofacitinib (Xeljanz®)
Phase of Development: Observational / Phase IV (Post-marketing)
Medical Monitor: Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the effectiveness of treatment with tofacitinib on disease activity (specifically, remission defined by PASDAS score ≤ 1.9) in a real-world setting over a 12-month observation period. **Secondary Objectives:** To evaluate the effectiveness of tofacitinib on achieving Low Disease Activity (LDA), DAPSA remission (score ≤ 4.0), and improvement in patient Quality of Life (measured by HAQ-DI) in routine clinical practice.

2.2 Background & Rationale To address the need for real-world evidence regarding the use of tofacitinib in patients with moderate to severe Psoriatic Arthritis (PsA). While randomized controlled trials demonstrate efficacy, understanding how tofacitinib performs in routine clinical practice, including its impact on composite disease activity measures and quality of life over 12 months, is critical for optimizing patient care.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational, Prospective Cohort **Control Type:** Single-arm (Non-comparative) **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target N: 116 evaluable patients (Initially planned for up to 500 across multiple countries) **Number of Sites:** Multinational (Including centers in Belgium, Czech Republic, Denmark, Finland, France, Israel, Netherlands, Spain, Sweden, and Switzerland) **Estimated Study Start:** 17-May-2021 **Estimated Primary Completion:** 30-Jul-2024

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Patients aged ≥ 18 years. 2. Moderate to severe PsA disease activity diagnosed. 3. Patients for whom the physician's decision has been made to initiate treatment with tofacitinib, in usual clinical practice conditions and in compliance with the local label. 4. Patients are treatment naïve to tofacitinib on the date of providing informed consent.

4.2 Exclusion Criteria 1. Contraindications according to the Xeljanz® (tofacitinib) Prescribing Information. 2. Receipt of any investigational drug within 3 months before study inclusion. 3. Patient is pregnant or breastfeeding. 4. Recent herpes zoster infection (within past 6 months) or history of severe disseminated herpes zoster infection. 5. Concomitant treatment with a biological disease-modifying antirheumatic drug (bDMARD).

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Tofacitinib administered as per standard routine clinical practice (e.g., 5 mg BID orally). **Route of Administration:** Oral. **Duration of Treatment:** Up to 12 months of observational follow-up.

5.2 Concomitant & Prohibited Therapy Permitted: Conventional synthetic DMARDs (csDMARDs), provided patients have not had a treatment change in the past 3 months. **Prohibited:** Concomitant treatment with bDMARDs or receipt of any investigational drug within 3 months prior to inclusion.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
1	Screening / Baseline	Day 0 Informed Consent, Medical History, baseline PASDAS/DAPSA evaluation, HAQ-DI assessment
2	Interim	Month 6 (or routine care) Disease Activity Assessment (PASDAS, DAPSA), Quality of Life (HAQ-DI), Safety assessment
3	End of Study	Month 12 Final PASDAS/DAPSA scoring, HAQ-DI, Exit Interview, Safety/AE recording

(Note: Exact visit schedules follow routine clinical practice standards at the investigator's discretion)

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Remission rate at 12 months, defined as a Psoriatic Arthritis Disease Activity Score (PASDAS) of ≤ 1.9 . **Measurement Method:** Clinical evaluation and patient-reported outcomes collected during standard care.

7.2 Secondary & Exploratory Endpoints * Disease Activity in Psoriatic Arthritis (DAPSA) remission (score ≤ 4.0). * Low Disease Activity (LDA) rate defined by a PASDAS score of ≤ 3.2 . * Change from baseline in the Health Assessment Questionnaire-Disability Index (HAQ-DI) score.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard observational collection via routine clinical practice visits and patient reports over the 12-month period. **Serious Adverse Events (SAEs):** Pharmacovigilance reporting timelines strictly enforced (typically within 24 hours of site awareness). **Data Monitoring Committee (DMC):** Managed by the Sponsor (Pfizer) to oversee global real-world safety data.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Full Analysis Set (FAS) comprising all enrolled patients who received at least one dose of tofacitinib and had at least one post-baseline effectiveness evaluation. **Sample Size Justification:** A sample size of ~116 patients is deemed adequate to provide descriptive statistics for real-world PASDAS/DAPSA remission rates with reasonable confidence intervals in the observed cohort. **Handling of Missing Data:** As per observational standards, analyses are generally performed on observed data, with methods like Last Observation Carried Forward (LOCF) or Multiple Imputation utilized for longitudinal continuous endpoints (e.g., HAQ-DI) where appropriate.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Pharmacoepidemiology Practices (GPP), applicable Good Clinical Practice (GCP) guidelines, and the Declaration of Helsinki. **Monitoring Plan:** Periodic source data verification and risk-based monitoring tailored for non-interventional studies to ensure data validity from electronic health records (EHR). **Audit Trail:** Secure eCRF systems equipped with full audit trails and data clarification forms to track clinical pathway documentation.

11. Study Identifiers & Governance

Primary ID: A3921332 **Registry Number:** NCT04517669 **Sponsor/Lead Organization:** Pfizer **Study Title:** TOPSATI **Official Regulatory Title:** A Non-Interventional Multinational Study of Tofacitinib in Patients Treated for Psoriatic Arthritis

12. Clinical Framework (Psoriatic Arthritis Specific)

Primary Condition: Psoriatic Arthritis (PsA) **Diagnosis Threshold:** Moderate to severe PsA disease activity **Medical Methodology:** Routine Clinical Practice (Non-interventional) **Optimization Target:** Remission (PASDAS ≤ 1.9 or DAPSA ≤ 4.0)

13. Study Procedures & Methodology

Management Pathway: Standard clinical treatment pathway for PsA **Education Protocol (HCP):** N/A (Observational study; treatment per local label) **Education Protocol (Patient):** N/A (Observational study; routine clinical care counseling) **Quality Audit Mechanism:** Periodic chart review & EHR extraction monitoring

14. Observation & Follow-Up Schedule

Total Duration: 12 months follow-up per patient **Onsite Visit Cadence:** Dictated by standard clinical practice (typically Baseline, Month 6, Month 12) **Remote Monitoring Frequency:** As applicable per institutional and sponsor guidelines for non-interventional data **Checklist Review Interval:** Data entered retrospectively or concurrently post-routine visits

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				