

1. Administrative & Study Identification

Full Study Title: ATGAM INTRAVENOUS INFUSION 250mg GENERAL INVESTIGATION **Short Title/Acronym:** ATGAM General Investigation **Protocol Number:** Pfizer Internal Study ID **NCT Number:** NCT06039020 **Sponsor Name:** Pfizer **Investigational Product:** ATGAM (Equine Thymocyte Immune Globulin / anti-human thymocyte immunoglobulin, equine) Intravenous Infusion 250mg **Phase of Development:** Phase IV / Observational (Post-Marketing) **Medical Monitor:** Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective **Primary Objective:** To confirm the safety of ATGAM in patients with moderate to severe aplastic anemia under actual clinical use in Japan. **Secondary Objectives:** To evaluate the real-world effectiveness and long-term tolerability of ATGAM in the target population over a 24-week observation period.

2.2 Background & Rationale To monitor and document the real-world safety profile of ATGAM (equine anti-human thymocyte immunoglobulin) when treating Japanese patients with moderate to severe aplastic anemia. This general investigation fulfills post-marketing surveillance requirements to ensure the continued safe and effective use of the drug in routine clinical practice in Japan.

3. Study Design & Methodology

3.1 Design Framework **Study Type:** Observational (General Investigation / Post-Marketing Surveillance) **Control Type:** Single-arm (No control group) **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration **Target N\$:** Unspecified (based on the actual number of patients who receive the treatment in the designated sites during the surveillance period) **Number of Sites:** Multiple clinical sites across Japan **Estimated Study Start:** 08-Sep-2023 **Estimated Primary Completion:** 24 weeks (6 months) after the last patient initiates administration

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Confirmed clinical diagnosis of moderate to severe aplastic anemia. 2. Patients who are scheduled to receive or are actively receiving ATGAM therapy.

4.2 Exclusion Criteria 1. Patients without a confirmed diagnosis of moderate to severe aplastic anemia. 2. Patients with a known hypersensitivity to equine (horse) gamma globulin preparations or any of the excipients.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Usually, 40 mg of anti-human thymocyte immunoglobulin, equine per kilogram of body weight, administered once daily. **Route of Administration:** Slow intravenous (IV) drip infusion. **Duration of Treatment:** Treatment is administered for a duration of 4 days. The total observation period is 24 weeks (6 months) from the start of administration (Day 1).

5.2 Concomitant & Prohibited Therapy Permitted: Standard concomitant therapies for aplastic anemia (e.g., cyclosporine, hematologic support, transfusions) as prescribed by the treating physician under standard Japanese clinical practice. **Prohibited:** Concurrent administration of other unapproved experimental treatments that might confound safety evaluations.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -14 to 0	Diagnosis of moderate to severe aplastic anemia, Treatment decision
Baseline	Day 0 to Day 4	Start of ATGAM IV infusion (Day 1), Daily Dosing for 4 days, Baseline Safety Assessments
Interim	Weeks 1 to 23	Routine clinical follow-up, Hematologic Response Monitoring, Safety/AE tracking
End of Study	Week 24	Final Observation Check, Safety Event Reconciliation (Observation continues to 24 weeks even if treatment was discontinued earlier)

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Incidence and severity of Adverse Events (AEs) and Serious Adverse Events (SAEs) occurring over the 24-week observation period. **Measurement Method:** Clinical safety tracking, laboratory monitoring, and spontaneous reports by the treating HCPs per post-marketing surveillance guidelines.

7.2 Secondary & Exploratory Endpoints Real-world clinical effectiveness, including hematologic recovery and survival rates at 24 weeks following ATGAM administration.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard collection via HCP reporting and routine post-marketing pharmacovigilance mechanisms. **Serious Adverse Events (SAEs):** Expedited reporting required as per Japanese regulatory guidelines for post-marketing safety. **Data Monitoring Committee (DMC):** Pharmacovigilance and Safety Management Teams managed by Pfizer.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all patients who receive at least one dose of ATGAM and have post-treatment safety data). **Sample Size Justification:** Descriptive observational study; powered based on the cumulative real-world enrollment in Japan rather than strict hypothesis testing. **Handling of Missing Data:** Last Observation Carried Forward (LOCF) for continuous variables; safety events tabulated as observed.

10. Quality Assurance & Regulatory Compliance

GCP/GPSP Compliance: This study will be conducted in accordance with Good Post-marketing Study Practice (GPSP) and local Japanese regulatory guidelines. **Monitoring Plan:** Centralized remote data monitoring of Case Report Forms (CRFs) completed by the treating institutions. **Audit Trail:** Electronic safety database locking and data clarification logs tracking real-world clinical reports.

11. Study Identifiers & Governance

Primary ID: Pfizer Internal Registry **Registry Number:** NCT06039020
Sponsor/Lead Organization: Pfizer **Study Title:** ATGAM General Investigation **Official Regulatory Title:** ATGAM INTRAVENOUS INFUSION 250mg GENERAL INVESTIGATION

12. Clinical Framework (Aplastic Anemia Specific)

Primary Condition: Moderate to severe aplastic anemia **Diagnosis Threshold:** Clinically confirmed marrow failure matching moderate or severe aplastic anemia criteria **Medical Methodology:** Immunosuppressive Therapy (IST) **Optimization Target:** Immune modulation, hematologic recovery, and prevention of severe complications

13. Study Procedures & Methodology

Management Pathway: Standard clinical pathway for aplastic anemia under actual use in Japan **Education Protocol (HCP):** Safety reporting training and GPSP documentation guidance **Education Protocol (Patient):** Routine clinical informed consent and counseling on IST side effects **Quality Audit Mechanism:** Post-marketing pharmacovigilance tracking and query resolution by the sponsor

14. Observation & Follow-Up Schedule

Total Duration: 24 weeks (6 months) from the start of administration **Onsite Visit Cadence:** Dictated by the patient's individual routine care schedule **Remote Monitoring Frequency:** Continuous safety monitoring via HCP adverse event submissions **Checklist Review Interval:** Final clinical and safety outcomes collected at Week 24

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				