

1. Administrative & Study Identification

Full Study Title: An Open-label Extension Study to Evaluate the Long-term Safety of Inclacumab Administered to Participants With Sick Cell Disease Who Have Participated in an Inclacumab Clinical Trial **Short Title/Acronym:** THRIVE-133 / Long-term Safety of Inclacumab in SCD **Protocol Number:** GBT2104-133 (Other IDs: C5361003, 2020-005289-32) **NCT Number:** NCT05348915 **Sponsor Name:** Pfizer **Investigational Product:** Inclacumab (PF-07940370) **Phase of Development:** Phase 3 **Medical Monitor:** Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the safety of long-term administration of inclacumab in adult and adolescent participants with sickle cell disease (SCD) who have completed a prior study of inclacumab. **Secondary Objectives:** To evaluate the long-term efficacy, including the frequency of vaso-occlusive crises (VOCs) and overall healthcare utilization, over an extended treatment period.

2.2 Background & Rationale In SCD, sickled red blood cells and abnormal cell adhesion lead to painful and potentially life-threatening vaso-occlusive crises (VOCs). Inclacumab is a monoclonal antibody that selectively targets P-selectin, a cell adhesion molecule that mediates the interactions causing these blockages. This open-label extension study provides continued access to treatment for patients who completed parent trials (e.g., THRIVE-131, THRIVE-132) and generates long-term real-world safety and efficacy data regarding inclacumab administration.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Single-arm (Open-label extension) **Blinding:** Open-label **Randomization:** N/A (Rollover participants)

3.2 Planned Enrollment & Duration Target N\$: 242 (Actual enrollment; initially targeted ~520) **Number of Sites:** Global Multicenter **Estimated Study Start:** 29-Mar-2022 **Estimated Primary Completion:** 07-Nov-2025 (Note: Study was terminated based on the results of the parent study GBT2104-131)

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Male or female participant ≥ 12 years of age with SCD who participated and received study drug in a GBT/Pfizer-sponsored inclacumab clinical study. 2. Participant has completed the originating inclacumab study within 30 calendar days of the Day 1 Visit. 3. Participant (and/or legal guardian) has provided written informed consent/assent.

4.2 Exclusion Criteria 1. Participant was lost to follow-up from the originating inclacumab clinical study. 2. Participant has any medical, psychological, safety, or behavioral conditions that, in the opinion of the Investigator, may confound safety interpretation, interfere with compliance, or preclude informed consent. 3. Female participants of childbearing potential or male participants unwilling to use highly effective methods of contraception consistently throughout the study and for at least 165 days after the last dose.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Inclacumab 30 mg/kg body weight administered every 12 weeks. **Route of Administration:** Intravenous (IV) infusion. **Duration of Treatment:** Long-term administration (initially planned for up to 5 years, subject to early termination).

5.2 Concomitant & Prohibited Therapy Permitted: Standard of care for Sickle Cell Disease, including management of VOCs and pain. **Prohibited:** Use of other investigational medical products during the study period.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening/Transition	Day -30 to 0	Transition from parent trial, Informed Consent, Pregnancy Test
Baseline	Day 1	First open-label IV dose of inclacumab
Interim	Every 12 Weeks	IV Infusion, Safety Labs, VOC Assessment, AE monitoring
End of Study	Variable	Final Safety Assessment, Exit Interview, Contraception counseling (165 days post-dose)

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Long-term safety and tolerability profile, assessed by the incidence and severity of Adverse Events (AEs), Serious Adverse Events (SAEs), and laboratory abnormalities over the study duration. **Measurement Method:** Clinical evaluation, laboratory tests, and patient reporting at regular intervals.

7.2 Secondary & Exploratory Endpoints * Incidence and frequency of Vaso-occlusive Crises (VOCs) per year. * Rates of 90-day hospital readmissions due to VOCs.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring via regular 12-week site visits for IV infusions and interval check-ins.

Serious Adverse Events (SAEs): Routine expedited reporting timeline (typically 24 hours to the sponsor). **Data Monitoring**

Committee (DMC): External/Internal safety review committee assigned by the sponsor to oversee long-term cumulative safety.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all participants receiving at least one dose of open-label inclacumab). **Sample Size**

Justification: Based on the number of rollover participants from originating pivotal trials (THRIVE-131, THRIVE-132); actual \$N = 242\$. **Handling of Missing Data:** No formal imputation for safety data; descriptive statistics summarizing cumulative exposure and adverse event rates.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Routine onsite and remote monitoring to ensure protocol adherence, proper IV administration, and safety reporting. **Audit Trail:** Electronic Case Report Form (eCRF) tracking for VOC events and hospitalizations.

11. Study Identifiers & Governance

Primary ID: GBT2104-133 **Registry Number:** NCT05348915 **Sponsor/Lead Organization:** Pfizer **Study Title:** A Study to Evaluate the Long-term Safety of Inclacumab Administered to Participants With Sickle Cell Disease **Official Regulatory Title:** An Open-label Extension Study to Evaluate the Long-term Safety of Inclacumab Administered to Participants With Sickle Cell Disease Who Have Participated in an Inclacumab Clinical Trial

12. Clinical Framework (Sickle Cell Disease Specific)

Primary Condition: Sickle Cell Disease (SCD) **Diagnosis Threshold:** Confirmed SCD diagnosis with a history of Vaso-occlusive Crises

Medical Methodology: Anti-P-selectin monoclonal antibody therapy

Optimization Target: Reduction in the frequency of VOCs and VOC-related hospitalizations

13. Study Procedures & Methodology

Management Pathway: Open-label long-term extension clinical pathway **Education Protocol (HCP):** Training on IV infusion protocols, P-selectin inhibitor safety profiles, and VOC recording **Education Protocol (Patient):** Contraception adherence (165 days post-dose), AE reporting, and continuous study participation requirements **Quality Audit Mechanism:** Tracking of timely IV administration every 12 weeks and accurate VOC event logging.

14. Observation & Follow-Up Schedule

Total Duration: Originally planned for up to 5 years (terminated early) **Onsite Visit Cadence:** Every 12 weeks for IV dosing **Remote Monitoring Frequency:** As needed between 12-week dosing intervals **Checklist Review Interval:** At every 12-week infusion visit

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---	
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]					
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]			Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]	[[]				