

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 02-Apr-2026

1. Administrative & Study Identification

Brief Study Title: A Study to Investigate Effects of Ocrelizumab Treatment on Neurofilament Light Chain (NfL) Levels and Participant Satisfaction in Participants With Multiple Sclerosis (MS)

Full Study Title: SurfSubQ - A Prospective Longitudinal Multicenter Observational Study to Investigate Neurofilament Light Chain Levels and Patient Satisfaction After Subcutaneous Ocrelizumab Administration in Persons With Multiple Sclerosis

Short Title/Acronym: SurfSubQ **Protocol Number:** ML45551 **NCT**

Number: NCT06780150 **Sponsor Name:** Roche (Hoffmann-La Roche)

Investigational Product: Ocrelizumab Subcutaneous (Ocrevus® / R04964913) **Phase of Development:** Not Applicable **Trial Status:**

RECRUITING **Medical Monitor:** Hoffmann-La Roche Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate participant satisfaction after administration of ocrelizumab subcutaneous (SC) after 12 months using the therapy administration satisfaction questionnaire subcutaneous (TASQ-SC). **Secondary Objectives:** To evaluate the Treatment Satisfaction Questionnaire for Medication (TSQM), changes in serum neurofilament light chain (sNfL) and serum glial fibrillary acidic protein (sGFAP) levels, changes in NEDA-3 (No Evidence of Disease Activity) status, and correlations with MRI lesion counts.

2.2 Background & Rationale Ocrelizumab SC may improve patient satisfaction and convenience for people with relapsing or primary progressive multiple sclerosis (pwRMS/pwPPMS). Patient-reported outcomes (PROs) are crucial for long-term therapy adherence, while serum neurofilament light chain (sNfL) acts as a proven biomarker for MS disease activity. This study collects real-world evidence (RWE) to explore the relationship between SC administration satisfaction and corresponding biomarker trends.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational (Prospective, non-interventional real-world study) **Control Type:** Single-arm (Cohort) **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target \$N\$: 349 participants **Number of Sites:** Multicenter (Germany and Switzerland) **Estimated**

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Diagnosis of MS; RMS and PPMS participants diagnosed according to the McDonald criteria of 2017. 2. First treatment during the course of MS therapy with ocrelizumab SC according to the local prescribing information, regardless of the reason for starting treatment with ocrelizumab. 3. Age ≥ 18 years.

4.2 Exclusion Criteria 1. Participation in interventional studies investigating disease-modifying therapies (DMTs) for MS. 2. Prior or simultaneous participation in the CONFIDENCE or MoOzaRt (ISRCTN55332718) non-interventional studies at the same study site. 3. Prior treatment with rituximab (MabThera®) or ublituximab (Briumvi®). 4. Severe psychiatric disability. 5. Pregnant and/or breastfeeding women.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: 920 mg every 6 months (initial treatment or switch). **Route of Administration:** Subcutaneous (SC) injection. **Duration of Treatment:** 12 months observation.

5.2 Concomitant & Prohibited Therapy Permitted: Routine MS therapies at the treating physician's discretion in accordance with local clinical practice and local labeling. **Prohibited:** Investigational DMTs for MS; prior use of specific anti-CD20 therapies (rituximab or ublituximab).

6. Study Timeline & Visit Schedule

Visit ID	Window	Key Activities/Procedures	:---	:---
:---	Screening/Baseline	Day 0	Inclusion/Exclusion verification, baseline PROs (TASQ-SC, TSQM), baseline sNfL/sGFAP blood draws, MRI evaluation (if available in routine care).	
Interim	Month 6 $\pm$ [Y]	Subcutaneous Ocrelizumab administration, routine clinical assessments.	End of Study	
Month 12 $\pm$ [Y]	Final PROs (TASQ-SC, TSQM), final sNfL and sGFAP lab draws, safety evaluation, and NEDA-3 status check.			

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Number of Participants by Their Level of Satisfaction With Ocrelizumab SC After 12 Months

Assessed Using TASQ-SC. **Measurement Method:** Therapy Administration Satisfaction Questionnaire Subcutaneous (TASQ-SC) is a 12-item, participant-reported questionnaire measuring the impact of treatment administration on five domains: Physical Impact, Psychological Impact, Impact on Activities of Daily Living, Convenience, and Satisfaction. Each of the domain/scale scores is scored on a scale of 1-100, where higher scores indicate higher overall satisfaction with the therapy.

7.2 Secondary & Exploratory Endpoints 1. Number of Participants by Their Level of Satisfaction With Ocrelizumab SC After the First Injection and After 6 Months Assessed Using TASQ-SC. 2. Change From Baseline in NfL Protein Levels in Serum. 3. Concentration of NfL Protein Levels in Serum. 4. Treatment Satisfaction Questionnaire for Medication (TSQM v1.4) score evaluated on 4 domains (Effectiveness, Side Effects, Convenience, Global Satisfaction). 5. Changes in serum glial fibrillary acidic protein (sGFAP) levels. 6. Changes in NEDA-3 (No Evidence of Disease Activity) defined by no confirmed MS relapse, no new/enlarging T2 lesions, no Gadolinium-positive T1 lesions, and no 6-month confirmed disability worsening.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard real-world pharmacovigilance per routine clinical practice alongside observational data collection. **Serious Adverse Events (SAEs):** Routine regulatory reporting timeline. **Data Monitoring Committee (DMC):** N/A (Standard non-interventional primary data collection).

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: All enrolled participants meeting criteria and receiving at least one dose of SC ocrelizumab in routine practice. **Sample Size Justification:** Targeted 349 participants to provide robust real-world data collection across Germany and Switzerland. **Handling of Missing Data:** The correlations between NfL/GFAP levels, MRI lesion counts, and MS relapses/NEDA-3 will be mapped using Pearson or Spearman correlation coefficients, handling missing data per standard observational guidelines.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This Non-Interventional Study (NIS) is conducted in accordance with Good Clinical Practice (GCP) and regional observational guidelines in Germany and Switzerland. **Monitoring Plan:** Real-world data extracted directly from routine MS clinic visits. **Audit Trail:** eCRF logging consistent with registry requirements for Roche Protocol ML45551.

11. Study Identifiers & Governance

Primary ID: ML45551 **Registry Number (NCT):** NCT06780150 **Posting ID:** 634208512431539219 **Sponsor/Lead Organization:** Roche **Study Title:** SurfSubQ **Official Regulatory Title:** SurfSubQ - A Prospective Longitudinal Multicenter Observational Study to Investigate Neurofilament Light Chain Levels and Patient Satisfaction After Subcutaneous Ocrelizumab Administration in Persons With Multiple Sclerosis

12. Clinical Framework (Multiple Sclerosis Specific)

Disease Area / Primary Condition: Multiple Sclerosis (Relapsing Multiple Sclerosis (RMS) and Primary Progressive Multiple Sclerosis (PPMS)) **Diagnosis Threshold:** McDonald criteria of 2017 **Medical Methodology:** Disease-Modifying Therapy (DMT) for MS **Optimization Target:** Subcutaneous administration transition and sustained patient adherence

13. Study Procedures & Methodology

Management Pathway: Real-world routine clinical practice pathway for MS **Education Protocol (HCP):** N/A (Non-interventional NIS) **Education Protocol (Patient):** N/A (Non-interventional NIS) **Quality Audit Mechanism:** TASQ-SC and TSQM standardized patient-reported outcome (PRO) questionnaires

14. Observation & Follow-Up Schedule

Total Duration: 12 months observational follow-up **Onsite Visit Cadence:** Baseline, Month 6, Month 12 (aligning with typical SC dosing intervals) **Remote Monitoring Frequency:** N/A **Checklist Review Interval:** N/A

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				