

1. Administrative & Study Identification

Full Official Title: A PHASE 1b/2 STUDY OF PF-07901801, A CD47 BLOCKING AGENT, WITH TAFASITAMAB AND LENALIDOMIDE FOR PARTICIPANTS WITH RELAPSED/REFRACTORY DIFFUSE LARGE B CELL LYMPHOMA NOT ELIGIBLE FOR STEM CELL TRANSPLANTATION **Short/Study Title:** Effects of Maplirpacept (PF-07901801), Tafasitamab, and Lenalidomide in People With Relapsed or Refractory Diffuse Large B-cell Lymphoma **Trial Status:** TERMINATED **Protocol Number:** C4971003 **NCT Number:** NCT05626322 **Sponsor/Company:** Pfizer **Investigational Product:** Maplirpacept (PF-07901801), Tafasitamab, and Lenalidomide **Phase of Development:** PHASE1, PHASE2 (Phase 1b/2) **Medical Monitor:** Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: * Phase 1b: To assess dose-limiting toxicities (DLTs) up to 28 days following the first dose of maplirpacept to determine the maximum tolerated dose or recommended Phase 2 doses. * Phase 2: To evaluate the Objective Response Rate (ORR) as per the Lugano Response Classification Criteria 2014. **Secondary Objectives:** To evaluate the safety, tolerability, and potential clinical benefits of maplirpacept in combination with tafasitamab and lenalidomide.

2.2 Background & Rationale Diffuse large B-cell lymphoma (DLBCL) is an aggressive type of non-Hodgkin lymphoma (NHL). An estimated 35-40% of patients relapse after their initial treatment. This study addresses an unmet medical need for patients with Relapsed/Refractory (R/R) DLBCL who are unable or unwilling to undergo intensive therapies such as high-dose chemotherapy with autologous stem cell transplantation (ASCT) or CAR-T cell therapy. Maplirpacept (PF-07901801) is a novel anti-CD47 molecule designed to enhance macrophage phagocytosis and antitumor activity by preventing CD47-mediated inhibitory signaling, which is hypothesized to work synergistically with tafasitamab and lenalidomide.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Active Comparator / Dose Optimization **Blinding:** Open-label **Randomization:** Randomized in Phase 2 (participants are randomized to high or low doses of maplirpacept)

3.2 Planned Enrollment & Duration Targeted Enrollment: 6 participants **Number of Sites:** Multicenter **Estimated Study Start:** 04-Aug-2023 **Estimated Primary Completion:** 12-Aug-2024

4. Subject Selection (Eligibility Criteria)

4.1 Key Inclusion Criteria 1. Histologically confirmed diagnosis of Diffuse Large B-Cell Lymphoma (DLBCL). 2. Relapsed or refractory disease. 3. Participant is not a candidate for or is unwilling to undergo high dose chemotherapy and subsequent stem cell transplant and/or is unable to receive chimeric antigen receptor (CAR) T-cell therapy. 4. Previous treatment with at least one prior line of systemic therapy (for phase 2, at least 1 and no more than 2 prior lines of systemic therapy). Prior therapy must include an anti-CD20 antibody. 5. Adequate bone marrow, hepatic and renal function. 6. Eastern Cooperative Oncology Group (ECOG) ≤ 2 . 7. Must provide a tumor tissue sample (fresh or archival, collected prior to start of treatment) for biomarker analysis.

4.2 Key Exclusion Criteria 1. Prior treatment with an anti-CD47 or anti-CD19 (other than CAR T) or immunomodulatory agents. 2. Prior allogeneic stem cell transplantation or autologous stem cell transplantation within 12 weeks prior to enrolment. 3. Participants with active, uncontrolled bacterial, fungal or viral infection.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Administered in 28-day cycles. * **Maplirpcept (PF-07901801):** Weekly for the first 3 cycles and then every 2 weeks. * **Tafasitamab:** Days 1, 4, 8, 15, and 22 in Cycle 1; weekly in Cycles 2 and 3; then every 2 weeks in Cycle 4 and beyond. * **Lenalidomide:** Taken every day for Days 1 to 21 of each 28-day cycle. **Route of Administration:** Intravenous (IV) infusion for Maplirpcept and Tafasitamab; Oral for Lenalidomide. **Duration of Treatment:** Lenalidomide is discontinued after 12 cycles. Maplirpcept and Tafasitamab can be continued until the lymphoma is no longer responding (disease progression) or unacceptable toxicity occurs.

5.2 Concomitant & Prohibited Therapy Permitted: Standard supportive care measures as per institutional guidelines. **Prohibited:** Other concurrent investigational agents or systemic anti-cancer therapies outside of the study protocol.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Screening	Day -28 to 0 Informed Consent, Medical History, Disease Assessment
Baseline	Cycle 1 Day 1	Randomization (for Phase 2), First Dose of Study Interventions	
Interim	Every 28 Days	Treatment Administration, Safety Labs, Efficacy Assessment (Scans)	End of Study
Treatment	Final Vital Signs, Disease Progression Follow-up, Survival Follow-up		

7. Outcome Measures (Endpoints)

7.1 Primary Outcomes * **Phase 1b:** Number of Participants With Dose Limiting Toxicities (DLTs). * *DLT details:* Hematological: Grade (G) 4 thrombocytopenia lasting ≥ 72 hours or $\leq 10,000/\text{mcL}$ anytime; $\geq \text{G3}$ thrombocytopenia with $\geq \text{G2}$ bleeding; G4 anemia; G4 neutropenia lasting ≥ 7 days; G3 febrile neutropenia lasting ≥ 7 days; G4 febrile neutropenia. Non-hematological: $\geq \text{G3}$ treatment-related non-hematologic toxicity; other $\geq \text{G2}$ PF-07901801-related non-hematologic toxicities requiring dose reduction/discontinuation.

7.2 Secondary Outcomes * **Phase 1b:** Number of Participants With Treatment Emergent Adverse Events (TEAEs). * **Phase 1b:** Number of Participants With Serious Treatment Emergent Adverse Events. * **Phase 1b:** Number of Participants With Treatment-Related AEs. * *Exploratory/Efficacy (Phase 2):* Complete Response (CR) rate, Progression-Free Survival (PFS), Overall Survival (OS), Objective Response Rate (ORR).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring from informed consent through the end of treatment via clinical assessment and laboratory tests. **Serious Adverse Events (SAEs):** Reported continuously within standard 24-hour timelines. **Data Monitoring Committee (DMC):** Utilized for reviewing safety data to dictate dose-escalation decisions and optimization during Phase 1b.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) for efficacy endpoints; Safety Analysis Set for all AE/DTL evaluations. **Sample Size Justification:** Targeted enrollment of $N=6$ participants to evaluate safety and determine optimal dosing. **Handling of Missing Data:** Standard survival analysis methodologies (e.g., censoring for time-to-event endpoints) will be applied.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Regular onsite and remote monitoring to ensure protocol adherence and data integrity. **Audit Trail:** Clinical data captured via secure electronic Case Report Forms (eCRF) utilizing automated data clarification rules.

11. Study Identifiers & Governance

Primary ID: {{C4971003}} **Registry Number:** {{NCT05626322}} **Posting ID:** {{651144921965066195}} **Sponsor/Lead Organization:** {{Pfizer}}

Study Title: {{Effects of Maplirpacept (PF-07901801),Tafasitamab, and Lenalidomide in People With Relapsed or Refractory Diffuse Large B-cell Lymphoma}} **Official Regulatory Title:** {{A PHASE 1b/2 STUDY OF PF-07901801, A CD47 BLOCKING AGENT, WITH TAFASITAMAB AND LENALIDOMIDE FOR PARTICIPANTS WITH RELAPSED/REFRACTORY DIFFUSE LARGE B CELL LYMPHOMA NOT ELIGIBLE FOR STEM CELL TRANSPLANTATION}}

12. Clinical Framework (Oncology Specific)

Disease Areas: {{Diffuse Large B-Cell Lymphoma}} **Diagnosis Threshold:** {{Confirmed DLBCL with \$\\ge 1\$ prior systemic therapy}} **Medical Methodology:** {{Combination Targeted Immunotherapy and Immunomodulatory Therapy}} **Optimization Target:** {{Phase 2 Dose Selection and Objective Response Rate (ORR)}}

13. Study Procedures & Methodology

Management Pathway: {{Standard oncology clinical pathway for R/R DLBCL}} **Education Protocol (HCP):** {{Investigator training on DLT criteria and infusion protocols}} **Education Protocol (Patient):** {{Informed consent and cycle-based treatment schedule education}} **Quality Audit Mechanism:** {{Routine clinical site monitoring and centralized data review}}

14. Observation & Follow-Up Schedule

Total Duration: {{Treatment continued until disease progression or unacceptable toxicity}} **Onsite Visit Cadence:** {{Weekly for the first 3 cycles, then bi-weekly}} **Remote Monitoring Frequency:** {{As per site monitoring plan}} **Checklist Review Interval:** {{Every 28-day cycle}}

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				