

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 02-Apr-2026

1. Administrative & Study Identification

Brief Study Title: Tofacitinib in Adult Patients With Moderate to Severe Ulcerative Colitis **Full Study Title:** Evaluation of the Clinical Benefit of ToFACitinib Treatment in Patients With Moderate to Severe Ulcerative Colitis Under Real-life Conditions of Use: TOFAst Study **Short Title/Acronym:** TO FAST Study **Protocol Number:** A3921360 **NCT Number:** NCT04424303 **Posting ID:** 666633849431694640 **Sponsor Name:** Pfizer (Pfizer Inc.) **Investigational Product:** Tofacitinib (Trade Name: Xeljanz) **Phase of Development:** Not Applicable **Trial Status:** RECRUITING (Currently recruiting participants) **Medical Monitor:** Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To describe the clinical benefit of tofacitinib 1 year after its initiation for the treatment of moderate to severe Ulcerative Colitis (UC) in routine clinical practice in France. **Secondary Objectives:** To report the clinical benefit 2 years after its initiation, to search for predictors of clinical benefit, improve understanding of the efficacy of treatment in a real-life setting (response and speed of response), and describe the characteristics of patients starting treatment with tofacitinib, real-life patterns of use, and patient adherence.

2.2 Background & Rationale Ulcerative colitis (UC) is a chronic inflammatory bowel disease affecting millions worldwide. Tofacitinib is an oral small-molecule Janus kinase (JAK) inhibitor approved for moderately to severely active UC. This non-interventional study in France (TO FAST) aims to generate real-world evidence on the drug's symptomatic remission capabilities independent of the administration of other common treatments like 5-ASA or corticosteroids during the observation period.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational (Non-interventional), Prospective cohort **Control Type:** Single-arm (Real-world cohort) **Blinding:** Open-label **Randomization:** N/A (Observational)

3.2 Planned Enrollment & Duration Target \$N\$: 280 patients (Targeted Enrollment) **Number of Sites:** Multiple clinics and

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Patients 18 years old or above. 2. Patients with a confirmed diagnosis of moderate to severe ulcerative colitis. 3. Patients for whom a gastroenterologist decides to initiate treatment with tofacitinib as per the French summary of product characteristics (SmPC). 4. Patients informed about the study procedures and who have received an information letter signed by the investigator.

4.2 Exclusion Criteria 1. Patients who have already received tofacitinib treatment before baseline. 2. Patients participating in another interventional clinical trial that could impact the outcome assessments. 3. Patients with contraindications to tofacitinib as per the current French SmPC (e.g., active severe infections, severe hepatic impairment, pregnancy). 4. Patients unwilling or unable to comply with the observational data collection schedule.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Tofacitinib (Xeljanz) oral administration prescribed at the discretion of the treating gastroenterologist, routinely reflecting SmPC dosing (e.g., 10 mg twice daily for induction, followed by 5 mg twice daily for maintenance). **Route of Administration:** Oral **Duration of Treatment:** Up to 2 years of observed follow-up.

5.2 Concomitant & Prohibited Therapy Permitted: Routine concomitant UC treatments including 5-ASA and corticosteroids as per normal clinical practice. **Prohibited:** Use of other concurrent biologic therapies or unapproved experimental drugs outside of standard care guidelines.

6. Study Timeline & Visit Schedule

Visit ID	Window	Key Activities/Procedures	:---	:---
:---	Screening/Baseline	Day 0	Diagnosis confirmation, tofacitinib initiation, Baseline Partial Mayo Score	Interim
Months 3-6	Routine follow-up, symptom tracking, safety and adverse event monitoring	Year 1 Follow-up	1 Year	Primary Endpoint Assessment (PR02 score), concomitant medication review
End of Study	2 Years	Secondary Efficacy Assessment, long-term adherence evaluation, Exit		

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Proportion of patients with clinical benefit one year after initiation of tofacitinib treatment. The definition of clinical benefit is independent of the discontinuation or not of tofacitinib treatment during the observation period. Clinical benefit at year 1 is defined on the basis of symptomatic remission evaluated with the PRO2 score ≤ 1 (absence of rectal bleeding and a stool frequency score between 0 and 1). *Note:* Patients who died or who had a colectomy or used another biologic/anti-JAK/immunosuppressant will be considered to be non-responders, as well as patients who used oral corticosteroids for UC, (regardless of the treatment duration) during the 3 months preceding the end of the observation period. The clinical benefit is independent of the administration or not of 5-ASA, or corticosteroids (not complying with the above definition) between 0 and 1 year. **Measurement Method:** Clinical Patient-Reported Outcomes (PRO2) collected in routine clinical practice.

7.2 Secondary & Exploratory Endpoints 1. Proportion of patients with clinical benefit of tofacitinib at 2 years. 2. Predictors of the clinical benefit at one year identified from the available baseline data. 3. Proportion of patients in clinical remission and still receiving tofacitinib. 4. Patient adherence and real-world treatment patterns. 5. Proportion of patients with serious and non-serious adverse events.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard real-world collection via patient interviews and medical chart reviews at routine visits. **Serious Adverse Events (SAEs):** Pharmacovigilance reporting per local regulatory standards (serious and non-serious AEs reported). **Data Monitoring Committee (DMC):** Managed through Pfizer pharmacovigilance and observational research protocols.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: The analysis set comprises all enrolled patients who received at least one dose of tofacitinib and have baseline assessments. **Sample Size Justification:** A target $N = 280$ is sufficient to provide a descriptive estimation of the proportion of real-world patients achieving clinical benefit at 1 year with a precise confidence interval. **Handling of Missing Data:** Missing data handled as observed in real-world scenarios (e.g., descriptive stats on available data, standard observational imputation methods where specified).

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Epidemiological Practice (GEP) guidelines and the Declaration of Helsinki. **Monitoring Plan:** Centralized monitoring of the registry and clinical records ensuring data fidelity from French sites. **Audit Trail:** Secure database tracking for source data verification and real-world electronic health record lock.

11. Study Identifiers & Governance

Primary ID: A3921360 **Registry Number:** NCT04424303 **Posting ID:** 666633849431694640 **Sponsor/Lead Organization:** Pfizer **Brief Study Title:** Tofacitinib in Adult Patients With Moderate to Severe Ulcerative Colitis **Official Regulatory Title:** Evaluation of the Clinical Benefit of ToFACitinib Treatment in Patients With Moderate to Severe Ulcerative Colitis Under Real-life Conditions of Use: TOFAst Study

12. Clinical Framework (Ulcerative Colitis Specific)

Primary Condition: Moderate to Severe Ulcerative Colitis (UC) **Diagnosis Threshold:** Clinically and endoscopically confirmed UC requiring systemic therapy **Medical Methodology:** Standard of Care following French SmPC for Tofacitinib **Optimization Target:** Symptomatic remission (PRO2 score ≤ 1 and reduction in partial Mayo score)

13. Study Procedures & Methodology

Management Pathway: Routine real-world clinical management of UC **Education Protocol (HCP):** Observational registry training and PRO2 scoring integration **Education Protocol (Patient):** Standard clinical guidance for oral JAK inhibitor compliance and safety monitoring **Quality Audit Mechanism:** Electronic Case Report Form (eCRF) logic checks and real-world data validation

14. Observation & Follow-Up Schedule

Total Duration: 2 Years **Onsite Visit Cadence:** Baseline, 1 Year, and 2 Years (alongside routine intermediate care) **Remote Monitoring Frequency:** N/A (Standard clinical contacts as needed) **Checklist Review Interval:** Annual clinical benefit review

15. Sign-off & Approval

Role	Name	Signature	Date		:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]					
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]					Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]					