

1. Administrative & Study Identification

Full Study Title: Phase 3, Multicenter, Randomized, Double-blind, Group Sequential, Placebo-controlled Study to Assess Efficacy and Safety of Rimegepant for the Treatment of Migraine (With or Without Aura) in Children and Adolescents ≥ 6 to < 18 Years of Age **Short Title/Acronym:** Randomized Study in Children and Adolescents With Migraine: Acute Treatment **Protocol Number:** BHV3000-311 **NCT Number:** NCT04649242 **Posting ID:** 694246504092247777 **Trial Status:** RECRUITING **Sponsor Name:** Pfizer (formerly Biohaven Pharmaceuticals, Inc.) **Investigational Product:** Rimegepant (Trade Name: Nurtec; ATC Code: N02CD06) / BHV-3000 (Oral dissolving tablet) **Phase of Development:** Phase 3 **Medical Monitor:** [Name and Contact Information]

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To test the safety and efficacy of the study medicine (rimegepant/BHV-3000) versus placebo in the acute treatment of moderate or severe migraine attacks in children and adolescents. **Secondary Objectives:** To evaluate secondary clinical endpoints such as pain relief, absence of the most bothersome symptom, and the long-term safety and tolerability of the drug.

2.2 Background & Rationale Migraine is a debilitating neurological disorder that affects about 7.7% to 9.1% of children and up to 15% of adolescents. Despite the significant burden, pediatric migraine represents a population with substantial unmet medical needs, as many conventional treatments lack robust efficacy or present unfavorable safety profiles for youth. This study aims to provide high-quality data on calcitonin gene-related peptide (CGRP) receptor antagonists (gepants) for safe and rapid acute treatment in this specific age group.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Placebo-controlled **Blinding:** Double-blind (incorporating a single-blind placebo run-in phase) **Randomization:** Randomized, Group Sequential

3.2 Planned Enrollment & Duration Target N : Estimated 2,100 enrolled participants (to achieve roughly 1,440 efficacy evaluable participants: ~1,200 adolescents and ~240 children under 12) **Number of Sites:** Approximately 190 sites globally (in

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Male and female participants ≥ 6 to < 18 years of age (participants must not reach their 18th birthday during the study). 2. History of migraine (with or without aura) for > 6 months before Screening, according to the IHS Classification ICHD-319 specifications for pediatric migraine. History may be verified using both medical records and recall. 3. History of 1 to 8 moderate or severe attacks per month during the 2 months prior to enrollment, with attacks lasting > 3 hours without treatment, and attacks occurring at intervals > 24 hours. 4. Prophylactic migraine medications are permitted if the dose has been stable for at least 12 weeks prior to Baseline, and is not expected to change. Participants may remain on one medication with prophylactic effects, excluding CGRP antagonists. Concomitant use of a CGRP antagonist (e.g., erenumab or fremanezumab) is prohibited. Previously discontinued prophylactics must have been stopped at least 90 days prior to Screening. 5. Ability to verbally distinguish between migraine and other types of headaches. 6. Weight > 40 kg at the Screening Visit. 7. Adequate venous access for blood sampling.

4.2 Exclusion Criteria 1. History of cluster headache or hemiplegic migraine headache. 2. Confounding and clinically significant pain syndrome that may interfere with participation. 3. Current psychiatric condition that is uncontrolled and/or untreated for a minimum of 6 months prior to Screening; lifetime history of psychosis and/or mania, suicidal behavior, or major psychiatric disorder. 4. Current diagnosis or history of substance abuse; positive drug test at Screening. 5. History of moderate or severe head trauma, seizure disorder, or other neurological disorder or systemic medical disease likely to affect central nervous system functioning. 6. Recent or planned surgery, requiring general anesthesia, < 8 weeks prior to the Screening Visit. 7. Participant has had gastrointestinal surgery that interferes with physiological absorption and motility (i.e., gastric bypass, duodenectomy, or gastric banding). 8. Current diagnosis of viral hepatitis or a history of liver disease. 9. Conditions considered clinically relevant in the context of the study such as uncontrolled hypertension (high blood pressure), diabetes, or a life-threatening allergy.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Single dose to be taken at the onset of a moderate or severe migraine attack. **Route of Administration:** Oral (dissolves under the tongue). **Duration of**

Treatment: Acute (single attack per observed period), with a subsequent follow-up period.

5.2 Concomitant & Prohibited Therapy Permitted: Stable doses of specific permitted medications (e.g., stable vitamins/supplements, one stable non-CGRP prophylactic). **Prohibited:** Concomitant use of CGRP antagonists (e.g., erenumab, fremanezumab), changes in prophylactic doses, or presence of gastrointestinal surgery altering absorption.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
1	Screening	Variable Informed Consent, Medical History, physical examination, ECG, drug test, verify eligibility criteria.
2	Baseline/Run-in	Single-blind phase Placebo run-in to evaluate baseline pain reporting and mitigate placebo effect.
3	Treatment	Acute Attack Randomization (1:1 rimegepant or placebo), medication administration, eDiary entry.
4	Follow-Up	Post-Treatment Safety labs, Efficacy Assessment, Questionnaires, AE reporting.

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Efficacy of rimegepant compared with placebo in the acute treatment of migraine in the adolescent population (≥ 12 to < 18 years of age). Measured by pain freedom using the number of patients reporting no pain, assessed on a 4-point numeric rating scale (0=none, 1=mild, 2=moderate, 3=severe).

7.2 Secondary & Exploratory Endpoints

1. Compare rimegepant with placebo for pain freedom in children and adolescents combined.
2. Freedom from the most bothersome symptom (MBS) associated with migraine in adolescents.
3. Probabilities of requiring rescue medication in adolescents.
4. Overall safety and tolerability including tracking of Adverse Events (AEs).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Collection via physical examinations, vital signs, ECG, laboratory samples, and participant reporting during study visits. **Serious Adverse Events (SAEs):** Reported continuously per standard pharmacovigilance timeline (typically 24 hours). **Data Monitoring Committee (DMC):** Utilized to monitor safety and efficacy across the group sequential design.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) for primary efficacy measurements; Safety Set for all subjects receiving at least one dose of the investigational product. **Sample Size Justification:** The trial aims to enroll ~2,100 subjects to ensure roughly 1,440 efficacy evaluable participants, providing adequate statistical power to differentiate rimegepant from placebo in the pediatric cohort. **Handling of Missing Data:** Standard imputation (e.g., LOCF or multiple imputation) will be applied for missing questionnaire and eDiary entries.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Scheduled onsite and remote monitoring of study centers (up to 190 locations). **Audit Trail:** Detailed electronic data capture (eDiary/eCRF) to ensure valid data on migraine characteristics and medication timing.

11. Study Identifiers & Governance

Primary ID: {{BHV3000-311}} **Registry Number:** {{NCT04649242}}
Posting ID: {{69424650409224777}} **Trial Status:** {{RECRUITING}}
Sponsor/Lead Organization: {{Pfizer / Biohaven Pharmaceuticals}}
Study Title: {{Randomized Study in Children and Adolescents With Migraine: Acute Treatment}} **Official Regulatory Title:** {{Phase 3, Multicenter, Randomized, Double-blind, Group Sequential, Placebo-controlled Study to Assess Efficacy and Safety of Rimegepant for the Treatment of Migraine (With or Without Aura) in Children and Adolescents ≥ 6 to < 18 Years of Age}}

12. Clinical Framework (Migraine Specific)

Primary Condition: {{Pediatric Migraine}} **Diagnosis Threshold:** {{History of migraine ≥ 6 months; 1-8 moderate or severe attacks per month}} **Medical Methodology:** {{Acute target-specific CGRP receptor antagonist administration}} **Optimization Target:** {{Pain freedom and absence of most bothersome symptoms at 2 hours}}

13. Study Procedures & Methodology

Management Pathway: {{Acute treatment of pediatric migraine episodes}} **Education Protocol (HCP):** {{Investigator training on ICHD-3 pediatric criteria and protocol adherence}} **Education Protocol (Patient):** {{Instructions for identifying qualifying

