

1. Administrative & Study Identification

Full Study Title: An Observational Study on RoActemra/Actemra (Tocilizumab) in Clinical Practice in Patients With Rheumatoid Arthritis (TRUST) **Short Title/Acronym:** TRUST **Protocol Number:** ML25728 **NCT Number:** NCT01394276 **Sponsor Name:** Hoffmann-La Roche **Investigational Product:** RoActemra / Actemra (Tocilizumab) **Phase of Development:** Observational / Phase IV Equivalent **Medical Monitor:** Hoffmann-La Roche Clinical Operations / Medical Monitor

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the efficacy and safety of RoActemra/Actemra (tocilizumab) in real-world clinical practice in patients with moderate to severe rheumatoid arthritis. **Secondary Objectives:** To evaluate the maintenance of low disease activity or disease remission, quality of life improvements (via the Health Assessment Questionnaire - HAQ), and to monitor the incidence of adverse events (AEs) and serious adverse events (SAEs).

2.2 Background & Rationale To address the need for real-world evidence regarding the effectiveness and safety of tocilizumab in clinical practice for patients with moderate to severe Rheumatoid Arthritis (RA). This study focuses on tracking disease activity improvements and safety profiles in patients who are inadequate responders to Disease Modifying Anti-Rheumatic Drugs (DMARDs) and anti-Tumor Necrosis Factor (anti-TNF) agents over a 12-month period.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational **Control Type:** Single-arm **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target \$N\$: Actual enrollment based on real-world clinical intake during the study window. **Number of Sites:** Multicenter (predominantly in the European Union / Italy). **Estimated Study Start:** May 2011 **Estimated Primary Completion:** April 2013

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Age ≥ 18 years. 2. Confirmed diagnosis of moderate to severe Rheumatoid Arthritis (RA). 3.

Inadequate responders to Disease Modifying Anti-Rheumatic Drugs (DMARDs) and anti-Tumor Necrosis Factors (anti-TNFs) agents, or patients who had received RoActemra treatment for 6 months prior to initiation of the study.

4.2 Exclusion Criteria 1. Current serious infection. 2. Hypersensitivity to the active component (tocilizumab) or any of the excipients. 3. Pregnant women.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Tocilizumab (TCZ) 8 milligrams per kilogram (mg/kg) body weight, once every 4 weeks. **Route of Administration:** Intravenous (IV). **Duration of Treatment:** 12 months, according to European Union (EU) approved dosage and Summary of Product Characteristics (SmPC).

5.2 Concomitant & Prohibited Therapy Permitted: Standard concomitant medications for RA (e.g., conventional DMARDs) as prescribed in real-world clinical practice per SmPC. **Prohibited:** Concurrent administration of live vaccines or other biological disease-modifying agents contraindicated by the tocilizumab SmPC.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
1	Screening / Baseline	Day 0 Informed Consent, Medical History, Baseline DAS28, HAQ, ESR/CRP Labs
2	Interim Observation	Every 4 Weeks IV Infusion of Tocilizumab, routine safety monitoring, AE/SAE tracking
3	End of Study	Month 12 Final Efficacy Assessment (DAS28, HAQ), Final Safety Evaluation, Exit Documentation

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Change in Disease Activity Score 28 (DAS28) and percentage of participants achieving low disease activity or disease remission (defined as DAS28 < 2.6) after 12 months. **Measurement Method:** Clinical evaluation combining 28 tender joint count, 28 swollen joint count, ESR/CRP measurements, and patient's global assessment of health via VAS (Visual Analog Scale).

7.2 Secondary & Exploratory Endpoints * Improvement in Health Assessment Questionnaire (HAQ) component scores. * Incidence, severity, and causality of Adverse Events (AEs) and Serious Adverse Events (SAEs).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Routine clinical data collection during observation; defined as any untoward medical occurrence in a participant whether or not related to the treatment. **Serious Adverse Events (SAEs):** Defined as events resulting in death, life-threatening conditions, hospitalization, disability, or congenital anomalies. Reported per standard pharmacovigilance tracking requirements. **Data Monitoring Committee (DMC):** Overseen by Hoffmann-La Roche safety and pharmacovigilance teams.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all patients who receive at least one dose of study medication) and Efficacy/Intent-to-Treat Set (patients with baseline and follow-up clinical assessments). **Sample Size Justification:** Observational in nature; sample size driven by real-world patient intake matching the inclusion criteria during the study window. **Handling of Missing Data:** Standard observational data conventions (e.g., observed cases or Last Observation Carried Forward [LOCF] for HAQ/DAS28 scores).

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP), the Declaration of Helsinki, and local observational study guidelines. **Monitoring Plan:** Observational data extraction via standard patient records/electronic Case Report Forms (eCRFs) across participating clinical sites. **Audit Trail:** Controlled eCRF data entry and standard query resolution processes maintained by the sponsor.

11. Study Identifiers & Governance

Primary ID: ML25728 **Registry Number:** NCT01394276 **Sponsor/Lead Organization:** Hoffmann-La Roche **Study Title:** TRUST **Official Regulatory Title:** Rheumatoid Arthritis Patients Treated With Tocilizumab in Real Clinical Practice: Effectiveness and Safety (TRUST Study)

12. Clinical Framework (Rheumatoid Arthritis Specific)

Primary Condition: Moderate to Severe Rheumatoid Arthritis (RA) **Diagnosis Threshold:** Inadequate response to DMARDs and anti-TNFs; elevated DAS28 **Medical Methodology:** EU Summary of Product Characteristics (SmPC) Guidelines / Standard Clinical Practice **Optimization Target:** Disease Remission (DAS28 < 2.6)

13. Study Procedures & Methodology

Management Pathway: Standard clinical pathway for Rheumatoid Arthritis with IV biologic therapy. **Education Protocol (HCP):** Implementation of SmPC and standardized joint count (DAS28) evaluations. **Education Protocol (Patient):** HAQ completion instructions and routine AE self-reporting guidance. **Quality Audit Mechanism:** Real-world observational data tracking and routine pharmacovigilance adherence.

14. Observation & Follow-Up Schedule

Total Duration: 12 Months **Onsite Visit Cadence:** Every 4 weeks (coinciding with the 8 mg/kg IV infusion schedule) **Remote Monitoring Frequency:** N/A (Predominantly onsite observational collection) **Checklist Review Interval:** Routine intervals per local clinical practice standards (typically baseline, mid-point, and 12 months)

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				