

1. Administrative & Study Identification

Full Study Title: A PHASE 1, RANDOMIZED, MULTI-CENTER, DOUBLE-BLIND, SPONSOR OPEN, PLACEBO-CONTROLLED, SINGLE DOSE-ESCALATION AND MULTIPLE DOSE STUDY TO EVALUATE THE SAFETY, TOLERABILITY, AND PHARMACOKINETICS OF PF-07985631 IN HEALTHY ADULT PARTICIPANTS
Short Title/Acronym: A Study to Learn How Different Amounts of the Study Medicine Called PF-07985631 Are Tolerated and Act in the Body of Healthy Adults
Protocol Number: Pfizer Internal Protocol (ID pending)
NCT Number: NCT06994897
Sponsor Name: Pfizer
Investigational Product: PF-07985631
Phase of Development: Phase 1
Medical Monitor / Clinical Operations Lead: Pfizer Clinical Operations Lead
Principal Investigator: [Name]
Lead Statistician: [Name]

2. Study Synopsis & Rationale

2.1 Study Objective
Primary Objective: To evaluate the safety, tolerability, and pharmacokinetics of single and multiple ascending doses of PF-07985631 or placebo in healthy adult participants.
Secondary Objectives: To evaluate pharmacokinetic (PK) parameters, including Area Under the Curve (AUC) and maximum observed concentration (Cmax) of the study medicine.

2.2 Background & Rationale
The purpose of this study is to learn about the safety and effects of the study medicine (called PF-07985631). PF-07985631 is being investigated for the possible future treatment of a kidney disease called IgA nephropathy. This Phase 1 study in healthy adults is required to determine the optimal dosage, how the drug is tolerated, and how it acts in the human body before proceeding to test in patient populations.

3. Study Design & Methodology

3.1 Design Framework
Study Type: Interventional
Control Type: Placebo-controlled
Blinding: Double-blind, Sponsor open
Randomization: Randomized

3.2 Planned Enrollment & Duration
Target N\$: 92 healthy adult participants
Number of Sites: Multi-center
Estimated Study Start: Q3 2025
Estimated Primary Completion: 08-Dec-2027

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Male or female between 18 and 45 years of age (up to 55 years of age for Japanese/Chinese/multiple dose participants). 2. Participants must be overtly healthy. 3. Japanese/Chinese cohorts only: Participants must have 4 biological Japanese/Chinese grandparents who were born in Japan/China.

4.2 Exclusion Criteria 1. Evidence or history of clinically significant hematological, renal, endocrine, pulmonary, gastrointestinal, cardiovascular, hepatic, psychiatric, neurological, or allergic disease. 2. Screening supine blood pressure (BP) ≥ 140 mm Hg (systolic) or ≥ 90 mm Hg (diastolic). 3. Estimated glomerular filtration rate (eGFR) < 75 mL/min/1.73 m². 4. History of infections requiring treatment within 28 days prior to Day 1, active/latent tuberculosis, or active infection at Day 1. 5. Chest X-ray showing any active disease in the chest, or pulmonary nodules > 0.5 cm in diameter that have not been previously evaluated, cavitary lesions or evidence of bronchiectasis. 6. Standard 12 lead ECG that demonstrates clinically relevant abnormalities that may affect participant safety or interpretation of study results.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Single dose-escalation (one time) or multiple ascending doses (once a month for 3 months) of PF-07985631 or matched placebo. **Route of Administration:** Subcutaneous (small needle injection in the abdomen, thigh, or back of the arm) or Intravenous (IV) infusion. **Duration of Treatment:** Single dose for most participants; up to 3 months of dosing for the multiple-dose cohort.

5.2 Concomitant & Prohibited Therapy Permitted: Standard allowed therapies per Phase 1 healthy volunteer unit protocols. **Prohibited:** Any vaccination within 6 weeks prior to Day 1 dosing or planned vaccination during the study.

6. Study Timeline & Visit Schedule

The study schedule includes Screening (Day -28 to -1) involving Informed Consent, Medical History, eGFR/BP check, Chest X-Ray, and ECG; Baseline (Day 1) involving Randomization, First Dose (IV or SC), and Safety Checks; Interim (Weeks 1-12) involving In-clinic stays (11-14 days for single dose; 4 days/month x 3 for multiple dose); and End of Study (Month 6) involving Final 8-10 outpatient visits, blood sampling, safety reviews, and PK analysis.

Visit ID	Window (Days)	Key Activities/Procedures	
Screening	Day -28 to -1	Informed Consent, Medical History, eGFR/BP check, Chest X-Ray, ECG	
Baseline	Day 1	Randomization, First Dose (IV or SC), Safety Checks	
Interim	Weeks 1-12	In-clinic stays (11-14 days for single dose; 4 days/month x 3 for multiple dose)	
End of Study	Month 6	Final 8-10 outpatient visits, blood sampling, safety reviews, and PK analysis	

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Number of Participants with Adverse Events (AEs) related to safety and tolerability from baseline to end of the study. **Measurement Method:** Clinical observation, vital signs, continuous safety reviews, and laboratory assessments.

7.2 Secondary & Exploratory Endpoints Area Under the Curve (AUC) for evaluating drug exposure over time. Maximum observed drug concentration (Cmax).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous daily monitoring during in-clinic stays and at the 8-10 follow-up outpatient visits.

Serious Adverse Events (SAEs): Must be reported to the sponsor and Institutional Review Board typically within 24 hours of site awareness. **Data Monitoring Committee (DMC):** Internal safety monitoring committee to review dose-escalation criteria and halt protocols if intolerable AEs occur.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all randomized participants receiving at least one dose of study medication/placebo). **Sample Size Justification:** $N=92$ is determined to be adequately powered to assess early Phase 1 safety, tolerability signals, and PK dose-proportionality. **Handling of Missing Data:** Standard handling for PK studies; participants withdrawing early may be replaced to ensure sufficient evaluable PK profiles.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Onsite Phase 1 unit monitoring to verify protocol adherence, dosing accuracy, and sample processing. **Audit**

Trail: eCRF entry tracking and source data verification (SDV) for all primary safety and PK data endpoints.

11. Study Identifiers & Governance

Primary ID: NCT06994897 **Registry Number:** NCT06994897 **Sponsor/Lead Organization:** Pfizer **Study Title:** A Study to Learn How Different Amounts of the Study Medicine Called PF-07985631 Are Tolerated and Act in the Body of Healthy Adults **Official Regulatory Title:** A PHASE 1, RANDOMIZED, MULTI-CENTER, DOUBLE-BLIND, SPONSOR OPEN, PLACEBO-CONTROLLED, SINGLE DOSE-ESCALATION AND MULTIPLE DOSE STUDY TO EVALUATE THE SAFETY, TOLERABILITY, AND PHARMACOKINETICS OF PF-07985631 IN HEALTHY ADULT PARTICIPANTS

12. Clinical Framework (IgA Nephropathy / Phase 1 Specific)

Primary Condition: Healthy Volunteers (Evaluating Investigational Drug for IgA Nephropathy) **Diagnosis Threshold:** Overtly healthy adults; normal baseline eGFR (≥ 75 mL/min/1.73 m²), normal BP (< 140/90 mm Hg) **Medical Methodology:** Phase 1 Pharmacokinetics & Safety Evaluation **Optimization Target:** Identifying Maximum Tolerated Dose (MTD) and optimal pharmacokinetic exposure

13. Study Procedures & Methodology

Management Pathway: Clinical Research Unit (CRU) Healthy Volunteer Confinement **Education Protocol (HCP):** Investigator site training on dose-escalation and safety halting rules **Education Protocol (Patient):** Strict informed consent orientation detailing in-clinic confinement rules and side-effect reporting **Quality Audit Mechanism:** Phase 1 unit Source Data Verification (SDV) & PK sampling timeline adherence tracking

14. Observation & Follow-Up Schedule

Total Duration: Up to 6 months total participation for multiple-dose cohorts **Onsite Visit Cadence:** Confinement periods (11-14 consecutive days or 4 days/month $\times$ 3) plus 8-10 follow-up visits **Remote Monitoring Frequency:** N/A (Study relies strictly on in-clinic monitoring and scheduled outpatient physical follow-ups) **Checklist Review Interval:** Daily during in-clinic stays

15. Sign-off & Approval

The final report will be signed and dated by the Principal Investigator, Clinical Operations Lead, and Lead Statistician to confirm approval.

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				