

1. Administrative & Study Identification

Full Study Title: A Safety and Efficacy Extension Study of Pertuzumab in Patients With Solid Tumors Previously Enrolled in a Hoffmann-La Roche-Sponsored Pertuzumab Clinical Trial **Official Title:** A Single-Arm Open-Label Multi-Centre Extension Study of Pertuzumab Administered As a Single Agent or in Combination With Other Anti-Cancer Therapies in Patients Previously Enrolled in a Hoffmann-La Roche-Sponsored Pertuzumab Study **Short Title/Acronym:** PerEx **Protocol Number:** M029406 (Other IDs: 2014-002048-42, 2023-505102-42-00) **NCT Number:** NCT02320435 **Sponsor Name:** Hoffmann-La Roche (Roche) **Investigational Product:** Pertuzumab (Perjeta / R04368451) **Phase of Development:** Phase 3 **Trial Status:** Active, not recruiting **Medical Monitor:** [Hoffmann-La Roche Clinical Operations Lead]

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the long-term safety and efficacy of pertuzumab over time when used alone or in combination with other cancer treatments in patients with solid tumors who have previously taken part in a parent pertuzumab clinical trial. **Secondary Objectives:** To evaluate investigator-assessed Objective Response Rate (ORR), Time to Progression (TTP), and Duration of Response.

2.2 Background & Rationale To provide continued access to pertuzumab for patients who are still receiving clinical benefit from the study treatment at the end of their respective parent studies. This extension study allows the sponsor to systematically collect long-term safety, tolerability, and efficacy data for pertuzumab-based therapies in patients with HER2-positive solid tumors.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Single-arm **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target \$N\$: 154 **Number of Sites:** Multi-center (Global) **Estimated Study Start:** 02-Feb-2015 **Estimated Primary Completion:** 31-Dec-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Informed consent. 2. Prior eligibility for, and receiving pertuzumab as an investigational medicinal product in, a Roche-sponsored study (either as a single agent or in combination with other anti-cancer drugs used in the Parent study) at the time of the Parent study closure. 3. Investigator's opinion that the patient continues to benefit from treatment.

4.2 Exclusion Criteria 1. Meets any of the exclusion criteria of the Parent protocol at the time the patient is considered for entry in the extension study. 2. Evidence of disease progression assessed according to Parent protocol before enrollment into the extension study. 3. Permanent discontinuation of pertuzumab for any reason during the Parent study, or between the end of the Parent study and before enrollment into the extension study. 4. Any unresolved or irreversible toxicities during the Parent study that require permanent discontinuation of pertuzumab, according to Parent protocol or local label. Delay of treatment to wait for resolution of toxicities is allowed as long as it is within the guidelines of the respective Parent protocol and does not contradict exclusion criterion below. 5. More than 9 weeks between the last dose of pertuzumab in the Parent study and the first dose of pertuzumab in the extension study. 6. Left ventricular ejection fraction $\leq 50\%$. 7. Any serious uncontrolled concomitant disease that would contraindicate the use of pertuzumab or that would put the patient at high risk for treatment-related complications. 8. Treatment with any anti-cancer treatment (other than any treatment given as permitted in the Parent protocol) in the time period between last treatment in the Parent study and the first dose pertuzumab in the extension study (i.e. up to 9 weeks). 9. Positive serum pregnancy test. 10. Women of child-bearing potential and men with partners of childbearing potential who do not agree to use a highly-effective non-hormonal form of contraception or two effective forms of non-hormonal contraception by the patient and/or partner for the duration of study treatment and for at least 7 months after the last dose of study medication. Male patients who do not agree to refrain from donating sperm during this same period. Male patients whose partner is pregnant who do not agree to use condoms for the duration of the pregnancy. 11. Concurrent participation in any therapeutic clinical trial (other than the Parent study). 12. Assessed by the Investigator to be unable or unwilling to comply with the requirements of the protocol.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: 420 mg of pertuzumab administered as an intravenous infusion every 3 weeks (following the dose, schedule, and guidance of the parent protocol). **Route of Administration:** Intravenous (IV) infusion. **Duration of Treatment:** Until disease progression, unacceptable

toxicity, investigator/patient decision, patient non-compliance, patient death, patient request to withdraw, or study termination by the Sponsor, whichever occurs first.

5.2 Concomitant & Prohibited Therapy Permitted: Trastuzumab, Vinorelbine, or other approved combination anti-cancer therapies as guided by the parent protocol. **Prohibited:** Any investigational therapies or therapeutic clinical trials outside of the extension study parameters.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -14 to 0	Informed Consent, verification of Parent Study completion, LVEF baseline review
Baseline	Day 0	Enrollment into Extension, First Dose
Interim	Every 3 Weeks	Pertuzumab Infusion, Safety Labs, Efficacy Assessment (Tumor Scans), AE/SAE monitoring
End of Study	At Progression	Final Vital Signs, LVEF Assessment, Exit Interview, IP Accountability

7. Outcome Measures (Endpoints)

7.1 Primary Outcomes 1. Progression-Free Survival (PFS). 2. Overall Survival (OS). 3. Number of Participants with Adverse Events by Severity, Classified According to the National Cancer Institute Common Terminology Criteria for Adverse Events (NCI-CTCAE) Version 4.03.

7.2 Secondary & Exploratory Endpoints * Investigator-assessed Objective Response Rate (ORR). * Duration of Response. * Time to Progression (TTP).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard collection of AEs evaluated in the safety population, graded by severity via the National Cancer Institute Common Terminology Criteria for Adverse Events (NCI-CTCAE) Version 4.03. **Serious Adverse Events (SAEs) / AESIs:** Monitoring for Adverse Events of Special Interest (AESIs) including asymptomatic decline in left ventricular ejection fraction (LVEF) requiring treatment/discontinuation, potential drug-induced liver injury (Hy's law), and suspected transmission of an infectious agent by the study drug. **Data Monitoring Committee (DMC):** Internal safety monitoring mechanisms managed by Hoffmann-La Roche.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) population for efficacy outcomes and Safety Set for patients receiving at least one dose of study treatment. **Sample Size Justification:** Descriptive statistics. Driven by the number of patients from parent studies who continue to derive clinical benefit (Target: 154 patients). **Handling of Missing Data:** [e.g., Last Observation Carried Forward (LOCF) for continuous variables; descriptive analysis parameters].

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Routine remote and onsite monitoring to ensure compliance with the parent protocol dosing and safety evaluations. **Audit Trail:** eCRF locking and adherence tracking to extension protocol safety requirements.

11. Study Identifiers & Governance

Primary ID: M029406 **Registry Number:** NCT02320435 **Sponsor/Lead Organization:** Hoffmann-La Roche (Roche) **Study Title:** PerEx **Official Regulatory Title:** A Single-Arm Open-Label Multi-Centre Extension Study of Pertuzumab Administered As a Single Agent or in Combination With Other Anti-Cancer Therapies in Patients Previously Enrolled in a Hoffmann-La Roche-Sponsored Pertuzumab Study

12. Clinical Framework (Oncology Specific)

Primary Condition: Solid Tumors (Cancer) / Solid Neoplasm **Diagnosis Threshold:** Confirmed solid tumors with previously established HER2 positivity and prior clinical benefit. **Medical Methodology:** HER2 Dimerization Inhibition. **Optimization Target:** Maintenance of therapeutic response, extended Progression-Free Survival (PFS) and Overall Survival (OS).

13. Study Procedures & Methodology

Management Pathway: Maintenance infusion pathway based on the established parent trial guidelines. **Education Protocol (HCP):** LVEF and AE monitoring protocol guidelines. **Education Protocol (Patient):** Information regarding long-term extension participation. **Quality Audit Mechanism:** Investigator site files, safety reporting compliance, and drug accountability.

14. Observation & Follow-Up Schedule

Total Duration: Until disease progression or December 2026 (estimated study completion). **Onsite Visit Cadence:** Every 3 weeks (Q3W) correlating with IV infusion schedule. **Remote Monitoring Frequency:** Between treatment cycles as required for AE follow-ups. **Checklist Review Interval:** Every 3 weeks prior to subsequent dosing.

15. Sign-off & Approval

Role	Name	Signature	Date		:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]					
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]					Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]					