

1. Administrative & Study Identification

Full Study Title: Zavicefta Combination for Intravenous Infusion Special Investigation - Surveillance on Patients With Sepsis or Renal Impairment (Creatinine Clearance $\leq$ 50 mL/Min)
Short Title/Acronym: A Study to Learn About the Study Medicine - Zavicefta in Patients With Sepsis or Loss of Kidney Function in Japan
Protocol Number: Not specified (Post-marketing surveillance)
NCT Number: NCT06864585
Trial Status: Not Yet Recruiting **Sponsor Name:** Pfizer Inc. / Pfizer Japan, Inc.
Investigational Product: Zavicefta (Avibactam sodium and Ceftazidime hydrate) [Preferred UMLS: Ceftazidime-Avibactam; ATC Code: J01DD02; Other Trade Names: Tazidime, Tazicef, Fortaz, Ceptaz, Avycaz] **Phase of Development:** Not Applicable **Medical Monitor:** Pfizer CT.gov Call Center

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the safety and effectiveness of Zavicefta under actual clinical practice in Japan in patients with sepsis or renal impairment. **Secondary Objectives:** To monitor real-world performance, patient outcomes, and safety profiles in critical conditions without intervention allocation or masking.

2.2 Background & Rationale To understand how Zavicefta performs in real-world clinical settings, potentially offering new treatment insights for serious infections and kidney function loss. The study targets serious health conditions like sepsis and kidney function loss, which require critical and effective treatment options.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational (Cohort model, prospective time perspective)
Control Type: Single-arm (No control group)
Blinding: Open-label (No masking)
Randomization: N/A

3.2 Planned Enrollment & Duration Target N\$: 59 Number of Sites: Multiple clinical sites across Japan
Estimated Study Start: 15-Sep-2025
Estimated Primary Completion: ~2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Patients who received Zavicefta for the first time after the launch of Zavicefta. 2. Patients who received Zavicefta for an infectious disease indicated for Zavicefta. 3. Patients with a diagnosis of sepsis and/or renal impairment (creatinine clearance ≤ 50 mL/min) at the start of the treatment with Zavicefta. 4. Individuals who understand the nature of this study and give consent for the provision of the information collected in this study to third parties and the use of the information for other than intended use.

4.2 Exclusion Criteria There are no exclusion criteria for this study.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Typically 2.5 g administered three times daily (as per standard labeling). **Route of Administration:** Intravenous (IV) Infusion. **Duration of Treatment:** Real-world administration, followed prospectively from Day 1 to Day 28.

5.2 Concomitant & Prohibited Therapy Permitted: Standard of care treatments for sepsis, renal impairment, and underlying comorbidities according to the treating physician's judgment. **Prohibited:** Any medications contraindicated in the local product label for Zavicefta.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Baseline	Day 1 Start of Zavicefta administration, Patient Consent, Medical History, Baseline Labs
Observation	Day 1 to 28	Clinical evaluation, Safety & Efficacy Monitoring under real-world practice	End of Study
Day 28	Final safety and clinical outcome assessment, AE/SAE recording		

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Number of Participants with adverse drug reactions (ADRs). **Measurement Method:** Clinical observation and physician-reported outcomes spanning from Day 1 to Day 28.

7.2 Secondary Endpoints 1. Number of Participants with Clinical Effectiveness. 2. Number of Participants with Clinical Effectiveness at the time of Test of Cure.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard collection via physician reporting during the 28-day prospective observation period.

Serious Adverse Events (SAEs): Routine pharmacovigilance and expedited reporting per local Japanese regulatory guidelines.

Data Monitoring Committee (DMC): Managed through Pfizer's internal pharmacovigilance and surveillance groups.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all enrolled patients who received at least one dose of Zavicefta and have post-dose safety data). **Sample Size Justification:** Target enrollment of 59 participants, determined by real-world clinical usage during the surveillance period rather than power calculations for hypothesis testing. **Handling of Missing Data:** Descriptive statistics typical for post-marketing surveillance; missing data will generally not be imputed.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Post-marketing Study Practice (GPSP) and local Japanese regulatory requirements for observational special investigations.

Monitoring Plan: Standardized collection of real-world data without protocol-mandated interventions. **Audit Trail:** Electronic data capture (eCRF) with tracking of physician reports and data clarification forms.

11. Study Identifiers & Governance

Posting ID: 86646833812350412 **Primary ID:** Pfizer Internal Special Investigation ID **Registry Number:** NCT06864585 **Sponsor/Lead**

Organization: Pfizer Japan, Inc. **Study Title:** A Study to Learn About the Study Medicine - Zavicefta in Patients With Sepsis or Loss of Kidney Function in Japan **Official Regulatory Title:** Zavicefta Combination for Intravenous Infusion Special Investigation - Surveillance on Patients With Sepsis or Renal Impairment (Creatinine Clearance $\leq$ 50 mL/Min)

12. Clinical Framework (Infectious Disease / Sepsis Specific)

Primary Condition: Infectious Diseases (Sepsis and/or Renal Impairment) **Diagnosis Threshold:** Creatinine clearance $\leq$ 50 mL/min for the renal impairment cohort **Medical Methodology:** Intravenous antibiotic therapy (Avibactam sodium/Ceftazidime hydrate) **Optimization Target:** Clinical cure of severe infection and stabilization of patient status

13. Study Procedures & Methodology

Management Pathway: Standard of care for severe bacterial infections in Japan. **Education Protocol (HCP):** Routine post-marketing drug safety instructions. **Education Protocol (Patient):** Informed consent detailing observational data collection. **Quality Audit Mechanism:** Real-world data verification and pharmacovigilance tracking.

14. Observation & Follow-Up Schedule

Total Duration: 28 Days **Onsite Visit Cadence:** Dictated by standard clinical practice (hospitalization/outpatient care). **Remote Monitoring Frequency:** N/A (Data collected via treating physicians). **Checklist Review Interval:** Final evaluation at Day 28 post-initiation.

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]				Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]				