

1. Administrative & Study Identification

Full Study Title: A Phase 3, Open-label Extension Study to Evaluate the Long-term Safety and Tolerability of KarXT + KarX-EC for the Treatment of Agitation Associated With Alzheimer's Disease (ADAGIO-3) **Short Title/Acronym:** ADAGIO-3 **Protocol Number:** CN012-0025 **NCT Number:** NCT06937229 **Sponsor Name:** Bristol-Myers Squibb **Investigational Product:** KarXT (xanomeline tartrate/trospium chloride immediate release capsule) + KarX-EC (xanomeline enteric capsules) **Phase of Development:** Phase 3 **Medical Monitor:** BMS Clinical Trials Contact Center

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the long-term efficacy, safety, and tolerability of the combined formulation of KarXT and KarX-EC in participants with agitation associated with Alzheimer's Disease who completed the parent studies CN012-0023 or CN012-0024. **Secondary Objectives:** To monitor and assess the long-term incidence of Treatment Emergent Adverse Events (TEAEs) and maintain symptomatic relief of agitation.

2.2 Background & Rationale Behavioral and psychological symptoms of dementia, specifically agitation, significantly impact patient outcomes and caregiver burden. Current pharmacologic treatments are limited by efficacy and severe safety concerns. KarXT (Cobenfy) offers a novel mechanism of action by modulating muscarinic receptors rather than traditional dopamine/serotonin pathways, potentially providing a much safer and effective alternative to manage Alzheimer's-related agitation.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Single-arm (Open-label extension) **Blinding:** Open-label **Randomization:** Non-Randomized (Single Group Assignment)

3.2 Planned Enrollment & Duration Target \$N\$: 600 participants **Number of Sites:** Approximately 241 locations globally **Estimated Study Start:** 02-Dec-2025 **Estimated Primary Completion:** 22-Jun-2029

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Participants must have completed parent study CN012-0023 or CN012-0024 per protocol. 2. Participants must have one identified caregiver who has sufficient contact (approximately 10 hours a week or more) and is willing to attend visits and report on the participant's status. 3. Age 55 to 90 years.

4.2 Exclusion Criteria 1. Clinically significant cardiovascular disease (e.g., untreated or unstable hypertension, clinically significant tachycardia). 2. Clinically significant pulmonary, renal, hematologic, GI, endocrine, immunologic, dermatologic, neurologic, or oncologic disease. 3. Any other condition that, in the investigator's opinion, would jeopardize the safety of the participant or the validity of the study results.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Combined formulation of KarXT (xanomeline tartrate/trospium chloride) and KarX-EC administered daily. **Route of Administration:** Oral (Immediate release and Enteric capsules). **Duration of Treatment:** Long-term extension, assessed up to approximately Week 30 for primary safety metrics.

5.2 Concomitant & Prohibited Therapy Permitted: Stable background medications for Alzheimer's Disease (must be stable for at least 12 weeks prior to parent trial enrollment). **Prohibited:** Medications that significantly prolong QT intervals, strong anticholinergics, or concurrent investigational treatments.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures		:---
:---	:---	Screening / Roll-over	Day -14 to 0	Consent (Participant & Caregiver), Verification of CN012-0023/0024 completion, Baseline Safety Labs
		Baseline	Day 0	Open-label treatment initiation, Vital Signs, Caregiver Assessment
	Interim	Weeks 4 to 24	Safety Labs, Efficacy Assessment (Agitation scales), AE/TEAE tracking	
	End of Primary Assessment	Week 30	Final Vital Signs, Treatment Emergent Adverse Event (TEAE) Evaluation, Drug Accountability	

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Number of Participants With Treatment Emergent Adverse Events (TEAEs) from baseline up to approximately Week 30. **Measurement Method:** Clinical assessment, physical examinations, and caregiver-reported events.

7.2 Secondary & Exploratory Endpoints Long-term reduction and maintenance of relief in agitation symptoms associated with Alzheimer's Disease. Evaluation of long-term tolerability (e.g., tracking of typical muscarinic side effects like dry mouth or constipation).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring via caregiver interviews, clinical exams, and laboratory assessments at every visit. **Serious Adverse Events (SAEs):** Reported to the sponsor (BMS) and regulatory authorities within 24 hours of site awareness. **Data Monitoring Committee (DMC):** Independent Data Monitoring Committee assigned to ensure the long-term safety of the geriatric Alzheimer's population.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all participants who receive at least one dose of open-label KarXT + KarX-EC). **Sample Size Justification:** Target enrollment of $N=600$ is based on the anticipated roll-over volume from parent phase 3 trials (CN012-0023 and CN012-0024). **Handling of Missing Data:** Mixed-Model for Repeated Measures (MMRM) for efficacy and Last Observation Carried Forward (LOCF) where applicable.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Periodic onsite and remote monitoring visits conducted by BMS Clinical Operations. **Audit Trail:** Secure eCRF systems with automated audit trails and data clarification forms.

11. Study Identifiers & Governance

Primary ID: CN012-0025 **Registry Number:** NCT06937229 **Sponsor/Lead Organization:** Bristol-Myers Squibb **Study Title:** ADAGIO-3 **Official Regulatory Title:** A Phase 3, Open-label Extension Study to Evaluate the Long-term Safety and Tolerability of KarXT + KarX-EC for the Treatment of Agitation Associated With Alzheimer's Disease (ADAGIO-3)

12. Clinical Framework (Alzheimer's Disease Specific)

Primary Condition: Alzheimer's Disease with Agitation **Diagnosis Threshold:** Completion of parent trial (history of agitation

meeting IPA consensus definition) **Medical Methodology:** Muscarinic Receptor Modulation (M1/M4 targeting) **Optimization Target:** Sustained reduction of agitation symptoms and behavioral stability

13. Study Procedures & Methodology

Management Pathway: Long-term open-label pharmacological management **Education Protocol (HCP):** Site initiation visits and Phase 3 specific protocol training **Education Protocol (Patient):** Caregiver instructions for daily dosing, monitoring, and side-effect reporting **Quality Audit Mechanism:** Centralized data review and routine site monitoring

14. Observation & Follow-Up Schedule

Total Duration: Minimum 30 weeks (Long-term extension) **Onsite Visit Cadence:** Monthly to quarterly physical and cognitive assessments **Remote Monitoring Frequency:** Caregiver phone check-ins during interval weeks **Checklist Review Interval:** Concomitant medications and TEAEs reviewed at every contact

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				