

1. Administrative & Study Identification

Full Study Title: A Rollover Study of Alectinib in Patients With Anaplastic Lymphoma Kinase (ALK)-Positive or Rearranged During Transfection (RET)-Positive Cancer **Short Title/Acronym:** Alectinib/Crizotinib Rollover Study **Protocol Number:** B039694 **NCT Number:** NCT03194893 **Sponsor Name:** Roche (Hoffmann-La Roche) **Investigational Product:** Alectinib (Alecensa) / Crizotinib (Xalkori) **Phase of Development:** Phase III **Trial Status:** Active, not recruiting **Medical Monitor:** Clinical Operations Lead, Hoffmann-La Roche

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To provide continued treatment with alectinib or crizotinib as applicable to participants with ALK- or RET-positive cancer who were previously enrolled in any Roche-sponsored alectinib study and who are deriving continued clinical benefit from alectinib or crizotinib in the parent trial at the time of parent trial closure. **Secondary Objectives:** To assess the long-term safety, tolerability, and continued efficacy of alectinib and crizotinib therapy in this patient population.

2.2 Background & Rationale To ensure uninterrupted access to life-prolonging targeted therapies (alectinib or crizotinib) for patients with ALK-positive or RET-positive malignancies (primarily Non-Small Cell Lung Cancer) transitioning from parent clinical trials. This rollover study is medically necessary to bridge the gap for patients where a switch to commercial drug supply is not yet feasible, ensuring continuity of care and continued tumor suppression.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional (Rollover study) **Control Type:** Single-arm (No placebo group; all participants receive active treatment) **Blinding:** Open-label **Randomization:** N/A (Rollover)

3.2 Planned Enrollment & Duration Target \$N\$: 200 participants **Number of Sites:** Global multicenter **Estimated Study Start:** May 16, 2017 **Estimated Primary Completion:** Q2 2024 / Ongoing

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Participants enrolled in a Roche-sponsored alectinib trial who are experiencing a clinical benefit from alectinib or crizotinib treatment at the time of discontinuation from the parent trial and for whom a switch to commercial supply is not feasible. 2. Collected study termination data, including efficacy and safety data, as required by the parent study on the electronic Case Report Form (eCRF). 3. For women who are not postmenopausal (≥ 12 months of non-therapy-induced amenorrhea) or surgically sterile (absence of ovaries and/or uterus): agreement to remain abstinent or use single or combined contraceptive methods that result in a failure rate of $< 1\%$ per year during the treatment period and for at least 3 months after the last dose of study drug. 4. For men: agreement to remain abstinent or use a contraceptive method that results in a failure rate of $< 1\%$ per year during the treatment period and for at least 3 months after the last dose of study drug.

4.2 Exclusion Criteria 1. Evidence of lack of clinical benefit in the parent trial during the screening phase of this rollover study. 2. Permanent discontinuation of alectinib or crizotinib for any reason during the parent study or before the first dose of study drug in the rollover study. 3. Evidence of an adverse event for which the parent protocol stipulates permanent discontinuation. 4. Pregnant or breastfeeding women. 5. Ongoing serious adverse event that has not resolved to baseline level or Grade ≤ 1 prior to the first dose of study treatment in the rollover study. 6. Treatment interruption for more than 21 days due to an adverse event since the last administration of alectinib or crizotinib in the parent trial. Any ongoing adverse events that require temporary treatment interruption must be resolved to baseline grade or assessed as stable and not requiring further treatment interruption by the investigator. 7. Administration of strong/potent cytochrome P450 (CYP) 3A inhibitors or inducers within 14 days prior to the first dose of treatment on this study and while on treatment with crizotinib. 8. Any psychological, familial, sociological, or geographical condition potentially hampering compliance with the study protocol requirements and/or follow-up procedures; these conditions should be discussed with the participant before trial entry.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Alectinib (e.g., 600 mg twice daily) or Crizotinib, administered based on the previously established safe dose in the parent trial. **Route of Administration:** Oral. **Duration of Treatment:** Continued until disease progression/loss of clinical benefit, unacceptable toxicity, withdrawal of consent, or commercial availability of the respective drug.

5.2 Concomitant & Prohibited Therapy Permitted: Standard supportive care medications for oncological symptoms. **Prohibited:** Strong or potent CYP3A inhibitors/inducers while on crizotinib therapy (requires at least a 14-day washout period prior to dosing).

6. Study Timeline & Visit Schedule

Visit ID	Window	Key Activities/Procedures
Screening	Parent Trial Exit	Assessment of continued clinical benefit, baseline safety checks, and termination data collection from the parent trial.
Baseline Day 0	Rollover enrollment, first dose in the rollover study, continuation of TKI therapy.	
Interim	Standard Care Intervals	Routine clinical oncology visits for drug dispensing, safety monitoring, and efficacy assessments.
End of Study	N/A	Final visit upon transition to commercial supply, unacceptable toxicity, or disease progression.

7. Outcome Measures (Endpoints)

7.1 Primary Endpoints

- 1. Number of Patients with Serious Adverse Events (SAEs), Non-serious Adverse Events (non-SAEs) and Adverse Events of Special Interest:** An AE is considered any unfavorable and unintended sign, symptom, or disease associated with use of study drug. Adverse events of special interest are cases of potential drug-induced liver injury that include elevated ALT or AST in combination with either elevated bilirubin or clinical jaundice and suspected transmission of an infectious agent.
- 2. Number of Patients With Clinically Significant Laboratory Values as per Protocol for Selected Safety Laboratory Parameters:** Selected parameters include alanine transaminase (ALT), aspartate transaminase (AST), total bilirubin, alkaline phosphatase (ALP), and blood creatine phosphokinase (CPK).

7.2 Secondary Endpoints

- 1. Number and Causes of Death Occurring on Study.** (*Exploratory evaluations of continued efficacy, including Disease-Free Survival (DFS) / Progression-Free Survival (PFS) may also be conducted*).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring during scheduled clinical visits, specifically tracking known TKI-associated toxicities (e.g., visual disturbances for crizotinib, hepatic/renal parameters for alectinib). **Serious Adverse Events (SAEs):** Prompt reporting to the sponsor required (typically within 24 hours). **Data Monitoring Committee (DMC):** Internal safety evaluation per Hoffmann-La Roche guidelines.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all enrolled patients who receive at least one dose of study medication in the rollover protocol). **Sample Size Justification:** $N = 200$. Enrollment is pragmatic, based strictly on the number of eligible patients rolling over from parent trials; the study is not statistically powered for primary hypothesis testing. **Handling of Missing Data:** Descriptive statistics will be utilized to report safety data, with no planned imputation for missing data.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study is conducted in strict accordance with Good Clinical Practice (GCP) guidelines and the Declaration of Helsinki. **Monitoring Plan:** Routine site monitoring to ensure patient safety, protocol compliance, and proper management of transition/rollover documentation. **Audit Trail:** Robust tracking of safety data and electronic Case Report Form (eCRF) completion.

11. Study Identifiers & Governance

Primary ID: B039694 **Registry Number:** NCT03194893 **Posting ID:** 970528783239893352 **Sponsor/Lead Organization:** Roche **Study Title:** A Rollover Study of Alectinib in Patients With Anaplastic Lymphoma Kinase (ALK)-Positive or Rearranged During Transfection (RET)-Positive Cancer **Official Regulatory Title:** A Multicenter, International, Rollover Study of Alectinib in Patients With Anaplastic Lymphoma Kinase (ALK)-Positive or Rearranged During Transfection (RET)-Positive Cancer

12. Clinical Framework (Oncology Specific)

Primary Condition / Disease Areas: Neoplasms (Preferred UMLS Name: Oncogene-Associated Disease) / ALK-positive or RET-positive Solid Tumors (NSCLC) **Diagnosis Threshold:** Histologically and genetically confirmed ALK or RET gene rearrangement **Medical Methodology:** Targeted Tyrosine Kinase Inhibitor (TKI) Therapy **Optimization Target:** Maintenance of clinical benefit and long-term tumor suppression

13. Study Procedures & Methodology

Management Pathway: Maintenance targeted therapy rollover pathway **Education Protocol (HCP):** Investigator guidelines on long-term AE monitoring and safe commercial drug transition **Education Protocol (Patient):** Counseling on strict medication adherence and prompt

adverse event symptom reporting **Quality Audit Mechanism:** Regular safety and efficacy data verification via eCRFs

14. Observation & Follow-Up Schedule

Total Duration: Until commercial availability, unacceptable toxicity, or disease progression **Onsite Visit Cadence:** Aligned with standard oncological care and TKI drug dispensing intervals **Remote Monitoring Frequency:** As clinically indicated between treatment cycles **Checklist Review Interval:** Standard AE and concomitant medication review at each clinic visit

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]				Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]				