

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 02-Apr-2026

1. Administrative & Study Identification

Full Study Title: Ondexxya for Intravenous Injection 200mg Drug Use Result Investigation (All Case Investigation) **Short Title/ Acronym:** Ondexxya All-Case Investigation **Protocol Number:** D9603C00001 **NCT Number:** NCT05454787 **Sponsor Name:** AstraZeneca **Investigational Product:** Ondexxya (andexanet alfa) for Intravenous Injection 200mg **Phase of Development:** Observational / Post-Marketing Surveillance (Phase IV equivalent) **Medical Monitor:** AstraZeneca Clinical Operations Lead **Study Contact:** AstraZeneca Clinical Study Information Center

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To investigate the status of occurrence of the safety specifications set for "Safety specification" (Thrombotic events, Infusion reaction, re-bleeding) in patients who received Ondexxya Intravenous Injection 200 mg under actual use conditions. In addition, this survey will be implemented to understand the following by collecting safety and effectiveness information under actual use conditions: detection of unknown adverse drug reactions; incidence of adverse drug reactions under actual use conditions of the drug; and factors that may affect safety or effectiveness. **Secondary Objectives:** To collect and understand effectiveness information (neutralization of FXa inhibitors and achievement of hemostasis) in a real-world clinical setting.

2.2 Background & Rationale To address the clinical need for real-world safety and effectiveness data regarding the use of Ondexxya as a reversal agent. This survey is conducted to monitor patients treated with Ondexxya to neutralize the anticoagulant effects of Factor Xa (FXa) inhibitors (apixaban, rivaroxaban, or edoxaban) when experiencing life-threatening or unarrestable bleeding episodes.

3. Study Design & Methodology

3.1 Design Framework Study Type: Observational (Drug Use Result Investigation) **Control Type:** Single-arm (All-case investigation) **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target N\$: 300 evaluable patients **Number of Sites:** Multicenter **Study Status:** Recruiting **Actual Study Start Date:** 02-Jun-2022 **Estimated Primary Completion Date:** 31-Aug-2026 **Estimated Study Completion Date:** 31-Aug-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. All patients treated with this drug (Ondexxya) to neutralize the anticoagulant effect of an FXa inhibitor (apixaban, rivaroxaban, or edoxaban). 2. Patients treated at the onset of a life-threatening bleeding episode or an unarrestable bleeding episode.

4.2 Exclusion Criteria 1. N/A (As an "All Case Investigation" in a post-marketing setting, all patients receiving the drug are intended to be included).

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Ondexxya for Intravenous Injection 200mg (Administered via low or high dose regimens based on the specific FXa inhibitor, its dose, and time since last administration). **Route of Administration:** Intravenous (IV) **Duration of Treatment:** Single administration sequence (initial IV bolus followed by a continuous IV infusion) per bleeding event.

5.2 Concomitant & Prohibited Therapy Permitted: Standard of care medical and surgical interventions for bleeding control and hemodynamic support. **Prohibited:** Routine re-administration of FXa inhibitors until hemostasis is fully secured and it is deemed clinically safe.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening / Day 0	Day 0	Bleeding event identification, baseline characteristics, FXa inhibitor history, Informed Consent (if applicable).
Baseline	Day 0	Administration of Ondexxya bolus and infusion, initial safety monitoring.
Interim	Hours 1-24	Acute Safety Labs, Hemostatic Efficacy Assessment.
End of Study	Day 30	Final Vital Signs, 30-day safety follow-up (AE/SAE recording), IP Accountability.

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Incidence of Adverse Events (AEs) and Serious Adverse Events (SAEs) as defined by the drug's safety specifications (Thrombotic events, Infusion reaction, re-bleeding) under actual use conditions. Detection of unknown adverse drug reactions and factors that may affect safety or

effectiveness. **Measurement Method:** Clinical evaluation and HCP reporting via Electronic Case Report Forms (eCRF).

7.2 Secondary & Exploratory Endpoints Real-world clinical effectiveness, defined as the proportion of patients achieving excellent or good hemostasis following Ondexxya administration. Incidence of thrombotic events within the 30-day follow-up period.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous post-marketing surveillance collection via HCP assessment, specifically investigating unknown adverse drug reactions, incidence of ADRs under actual use conditions, and identifying factors affecting safety. **Serious Adverse Events (SAEs):** Reported within 24 hours of site awareness, particularly monitoring for the predefined safety specifications: thromboembolic risks (thrombotic events), infusion reactions, and re-bleeding. **Data Monitoring Committee (DMC):** AstraZeneca internal pharmacovigilance and safety review board.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (All enrolled patients who received at least one dose of Ondexxya). **Sample Size Justification:** $N = 300$, determined to be adequate for detecting the occurrence status of predefined safety specifications during actual clinical use. **Handling of Missing Data:** Handled via standard observational practices (e.g., observed cases analysis without imputation for missing safety data).

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Vigilance Practice (GVP), Good Clinical Practice (GCP), and local regulatory requirements for post-marketing studies. **Monitoring Plan:** Centralized remote data monitoring to verify reporting accuracy and completeness from participating institutions. **Audit Trail:** Secure eCRF locking and documented data clarification forms to ensure data integrity.

11. Study Identifiers & Governance

Primary ID: D9603C00001 **Registry Number:** NCT05454787 **EudraCT / CTIS Identifier:** N/A **Sponsor/Lead Organization:** AstraZeneca **Study Title:** Ondexxya for Intravenous Injection 200mg Drug Use Result Investigation (All Case Investigation) **Official Regulatory Title:**

12. Clinical Framework (Reversal Agent Specific)

Primary Condition: Life-threatening or unarrestable bleeding episode **Conditions Studied:** Life Threatening Bleeding, Factor Xa Inhibitor **Diagnosis Threshold:** Clinically significant bleed in a patient actively taking an FXa inhibitor (apixaban, rivaroxaban, edoxaban) **Medical Methodology:** Direct Factor Xa inhibitor reversal **Optimization Target:** Rapid neutralization of anticoagulant effect and achievement of clinical hemostasis

13. Study Procedures & Methodology

Management Pathway: Acute emergency bleeding management protocol **Education Protocol (HCP):** Site-level training on appropriate Ondexxya dosing algorithms (low vs. high dose) based on prior FXa inhibitor timing. **Education Protocol (Patient):** N/A (Emergency administration; retrospective consent or legally authorized representative consent typically utilized). **Quality Audit Mechanism:** Continuous safety monitoring and pharmacovigilance tracking.

14. Observation & Follow-Up Schedule

Total Duration: Approximately 30-day safety follow-up post-administration **Onsite Visit Cadence:** Acute hospital stay (Days 0-X) **Remote Monitoring Frequency:** Standard safety check at Day 30 **Checklist Review Interval:** Continuous monitoring during acute admission

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	____	[DD-MMM-YYYY]				