

1. Administrative & Study Identification

Full Study Title: AN INTERVENTIONAL, PHASE 3, DOUBLE-BLIND, RANDOMIZED STUDY TO EVALUATE THE EFFICACY AND SAFETY OF PF-08634404 IN COMBINATION WITH CHEMOTHERAPY VERSUS BEVACIZUMAB IN COMBINATION WITH CHEMOTHERAPY IN TREATMENT-NAÏVE PARTICIPANTS WITH METASTATIC COLORECTAL CANCER **Short Title/Acronym:** PF-08634404 in Combination With Chemotherapy in mCRC **Protocol Number:** C6461003 **NCT Number:** NCT07222800 **Posting ID:** 983576661696837234 **Sponsor / Company Name:** Pfizer **Investigational Product:** PF-08634404 (in combination with mFOLFOX6 chemotherapy) **Phase of Development:** Phase 3 **Trial Status:** NOT_YET_RECRUITING (Not yet recruiting participants) **Medical Monitor:** Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To demonstrate that the investigational medicine PF-08634404 combined with mFOLFOX6 chemotherapy is superior to the standard of care (bevacizumab + mFOLFOX6) in prolonging Progression-Free Survival (PFS) and Overall Survival (OS). **Secondary Objectives:** To evaluate the Objective Response Rate (ORR), Duration of Response (DOR), and overall safety and tolerability of the combination therapy in adult participants.

2.2 Background & Rationale The goal of this study is to address the need for more effective first-line treatments for metastatic colorectal cancer (mCRC). The study evaluates if PF-08634404, when combined with approved chemotherapy, can better assist the immune system in finding and attacking cancer cells, and slowing down tumor growth, compared to the current standard active comparator, bevacizumab.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Active Comparator (Bevacizumab + Chemotherapy) **Blinding:** Double-blind **Randomization:** 1:1 (Randomized to one of 2 different treatment arms)

3.2 Planned Enrollment & Duration Target **NS:** 800 evaluable participants **Number of Sites:** Global multicenter trial **Estimated Study Start:** December 2025 **Estimated Primary Completion:** March 2030

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Age $\geq$ 18 years or older. 2. Diagnosed with metastatic (stage IV) colorectal cancer that has spread to other parts of the body. 3. Treatment-naïve for metastatic disease (have not previously received any anticancer treatment for mCRC).

4.2 Exclusion Criteria 1. Major surgery or severe trauma within 4 weeks prior to the first dose, or planned major surgery during the study. 2. History of allogeneic organ transplantation and allogeneic hematopoietic stem cell transplantation. 3. Any Grade ≥ 3 bleeding/hemorrhage events within 28 days of Cycle 1 Day 1, or prior history of clinically significant bleeding events.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Administered in cycles. Arm A receives PF-08634404 + chemotherapy; Arm B receives bevacizumab (trade names include Avastin, Vegzelma, Alymsys, Zirabev, Mvasi) + chemotherapy. **Route of Administration:** Intravenous (IV) Infusions. **Duration of Treatment:** Approximately 2 years, or until disease progression/unacceptable toxicity (overall study length is approx. 33 months per participant).

5.2 Concomitant & Prohibited Therapy Permitted: Standard approved chemotherapy for mCRC (mFOLFOX6). **Prohibited:** Concurrent administration of other investigational therapies or non-protocol systemic treatments for active autoimmune diseases.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Eligibility Checks, Baseline Imaging
Treatment Cycles	Per Cycle	IV Infusions, health checks, safety and tolerability monitoring
End of Treatment	Day 30 to 37 (Post-Last Dose)	Final health check, vital signs, review of any side effects
Follow-up	Every 12 Weeks	Phone/in-person review of health status, survival, and new treatments

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Progression-Free Survival (PFS) and Overall Survival (OS). **Measurement Method:** PFS per Response Evaluation Criteria in Solid Tumors 1.1 (RECIST 1.1) assessed by Blinded Independent Central Review (BICR).

7.2 Secondary & Exploratory Endpoints 1. Objective Response Rate (ORR). 2. Duration of Response (DOR).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Conducted continuously by trained medical staff during and after each IV treatment visit, continuing into the 12-week follow-up intervals. **Serious Adverse Events (SAEs):** Protocol-defined immediate reporting timelines (typically 24 hours). **Data Monitoring Committee (DMC):** Independent Data Monitoring Committee instituted per standard Phase 3 global oncology trial guidelines.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) for primary efficacy measures (OS and PFS); Safety Set for adverse events. **Sample Size Justification:** $N = 800$ appropriately powered to detect statistical superiority in PFS and OS between the PF-08634404 arm and the bevacizumab control arm. **Handling of Missing Data:** Standard survival analysis methodologies (e.g., right-censoring for survival outcomes) in alignment with FDA/EMA oncology guidelines.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Centralized and onsite monitoring by the sponsor. **Audit Trail:** Validated Electronic Data Capture (EDC) system with automated audit trails and Data Clarification Forms (DCFs).

11. Study Identifiers & Governance

Primary ID: C6461003 **Registry Number:** NCT07222800 **Posting ID:** 983576661696837234 **Sponsor/Lead Organization:** Pfizer **Study Title:** A Study to Learn About the Study Medicine Called PF-08634404 in Combination With Chemotherapy in Adult Participants With Metastatic Colorectal Cancer **Official Regulatory Title:** AN INTERVENTIONAL, PHASE 3, DOUBLE-BLIND, RANDOMIZED STUDY TO EVALUATE THE EFFICACY AND SAFETY OF PF-08634404 IN COMBINATION WITH CHEMOTHERAPY VERSUS BEVACIZUMAB IN COMBINATION WITH CHEMOTHERAPY IN TREATMENT-NAÏVE PARTICIPANTS WITH METASTATIC COLORECTAL CANCER

12. Clinical Framework (Metastatic Colorectal Cancer Specific)

Primary Condition: Metastatic Colorectal Cancer (mCRC) **Preferred UMLS Name:** Colorectal Neoplasms **Disease Areas:** Intestinal Neoplasms, Gastrointestinal Neoplasms, Digestive System Neoplasms, Neoplasms by Site, Digestive System Diseases, Gastrointestinal Diseases, Colonic Diseases, Intestinal Diseases, Rectal Diseases, Colorectal Neoplasms **Diagnosis Threshold:** Confirmed Stage IV disease **Medical Methodology:** Systemic target-based and cytotoxic combination therapy **Optimization Target:** Anti-tumor immunity activation / angiogenesis inhibition

13. Study Procedures & Methodology

Management Pathway: Standard oncologic systemic therapy cycles. **Education Protocol (HCP):** Site initiation visits and safety training for managing immunotherapy/chemotherapy adverse events. **Education Protocol (Patient):** Informed consent detailing the 50% randomization chance, IV infusion schedules, and symptom reporting. **Quality Audit Mechanism:** Electronic medical record reconciliation and rigorous Blinded Independent Central Review (BICR) of imaging.

14. Observation & Follow-Up Schedule

Total Duration: Approximately 33 months per participant **Onsite Visit Cadence:** Regular visits scheduled per treatment cycle lengths **Remote Monitoring Frequency:** Final end-of-treatment check followed by long-term follow-up **Checklist Review Interval:** Every 12 weeks (Q12W) via phone, in-person, or health records

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				