

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase Ib, Multicenter, Open-label Platform Study of Select Drug Combinations in Adult Patients With Lower Risk (Very Low, Low, or Intermediate Risk) Myelodysplastic Syndrome
Short Title:	Phase Ib Study of Select Drug Combinations in Patients With Lower Risk MDS
Protocol Number:	100479642675566773
NCT Number:	NCT04810611
Sponsor Name:	Novartis
Investigational Product:	canakinumab
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Dose interruption reduction: Dose tolerability 2. Incidence of DLTs: Incidence of dose limiting toxicities (DLTs) during the first 2 cycle of treatment during the dose escalation/confirmation part 3. Dose intensity: Dose tolerability
Secondary Obj.:	1. Reduction in red blood cell (RBC) / platelet transfusions from baseline in transfusion dependent patients 2. Duration of transfusion independence lasting for ≥ 8 weeks, ≥ 12 weeks, ≥ 16 weeks, ≥ 24 weeks in transfusion dependent patients 3. Change from baseline in hemoglobin (Hb) in transfusion dependent and transfusion independent patients

2.2 Background & Rationale

The purpose of this study was to characterize the safety, tolerability and confirm the dose for select single agents and combinations in patients with lower risk (very low, low, and intermediate risk) MDS.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, Status: TERMINATED
Target N:	33

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- Signed informed consent must be obtained prior to participation in the study.
- Patients must be ≥ 18 years of age at the time of signing the informed consent form (ICF).
- Patients must have a diagnosis prior to participation in the study of IPSS-R very low, low, or intermediate risk MDS with $\geq 10\%$ bone marrow blasts and one or more of the following:

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Symptomatic anemia with hemoglobin ≤ 10 g/dL that has relapsed after or is refractory to ESAs (or the patient is intolerant to ESAs)
2. Symptomatic anemia with hemoglobin ≤ 10 g/dL that is ESA-naïve with EPO level ≥ 500 /uL
3. Thrombocytopenia with platelets $\leq 30,000$ /uL or with clinically significant bleeding or bruising and platelets $\leq 50,000$ /uL
4. Neutropenia with an absolute neutrophil count (ANC) ≤ 500 / μ L or with recurrent and/or severe infections and an ANC that is ≤ 1000 / μ L and amenable to response assessments by International Working Group (IWG) response criteria in myelodysplasia (Cheson et al 2006)
4. Patients who are refractory to, intolerant of, or ineligible/unable to receive SOC therapeutic options including lenalidomide
5. Eastern Cooperative Oncology Group (ECOG) performance status (PS) ≥ 2
6. Patient must be a candidate for serial bone marrow aspirate and/or biopsy according to the institutions' guidelines and be willing to undergo a bone marrow aspirate and/or biopsy at screening, during and at the end of therapy on this study -

Key Exclusion Criteria:

1. Systemic antineoplastic therapy (including cytotoxic chemotherapy, alpha-interferon, kinase inhibitors or other targeted small molecules, and toxin-immunoconjugates) or any experimental therapy within 14 days or 5 half-lives, whichever is longer, before the first dose of study treatment.
2. History of hypersensitivity to any of the study treatments or its excipients or to drugs of similar chemical classes.
3. Patients with chronic myelomonocytic leukemia (CMML) or myelodysplastic/myeloproliferative neoplasms (MDS/MPN)
4. Use of hematopoietic colony-stimulating growth factors (e.g. G-CSF, GM-CSF, M-CSF), thrombopoietin mimetics or ESAs anytime ≥ 2 weeks (or 5 half-lives, whichever is longer) prior to start of study treatment.
5. Systemic chronic corticosteroid therapy (>10 mg/day prednisone or equivalent) or any immunosuppressive therapy within 7 days of first dose of study treatment. Topical, inhaled, nasal and ophthalmic steroids are allowed.
6. For arms containing canakinumab: Patients with ANC ≤ 500 / μ L

4.2 Exclusion Criteria

1. Systemic antineoplastic therapy (including cytotoxic chemotherapy, alpha-interferon, kinase inhibitors or other targeted small molecules, and toxin-immunoconjugates) or any experimental therapy within 14 days or 5 half-lives, whichever is longer, before the first dose of study treatment.
2. History of hypersensitivity to any of the study treatments or its excipients or to drugs of similar chemical classes.
3. Patients with chronic myelomonocytic leukemia (CMML) or myelodysplastic/myeloproliferative neoplasms (MDS/MPN)
4. Use of hematopoietic colony-stimulating growth factors (e.g. G-CSF, GM-CSF, M-CSF), thrombopoietin mimetics or ESAs anytime ≥ 2 weeks (or 5 half-lives, whichever is longer) prior to start of study treatment.
5. Systemic chronic corticosteroid therapy (>10 mg/day prednisone or equivalent) or any immunosuppressive therapy within 7 days of first dose of study treatment. Topical, inhaled, nasal and ophthalmic steroids are allowed.
6. For arms containing canakinumab: Patients with ANC ≤ 500 / μ L

11. Study Identifiers & Governance

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Primary ID: 100479642675566773
Registry Number: NCT04810611
Sponsor: Novartis
Study Title: Phase Ib Study of Select Drug Combinations in Patients With Lower Risk MDS

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____