

# Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

## 1. Administrative & Study Identification

|                          |                                                                                                                                                                                                                                             |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Full Study Title:        | A Phase 1 Open-label, Multicenter Study to Assess the Safety, Tolerability, Pharmacokinetics, and Pharmacodynamics of MK-4700 as Monotherapy and in Combination With Pembrolizumab in Participants With Advanced or Metastatic Solid Tumors |
| Short Title:             | A Study of MK-4700 Alone or With Pembrolizumab in Participants With Advanced or Metastatic Solid Tumors (MK-4700-001)                                                                                                                       |
| Protocol Number:         | 316411419878127298                                                                                                                                                                                                                          |
| NCT Number:              | NCT06894771                                                                                                                                                                                                                                 |
| Sponsor Name:            | Merck                                                                                                                                                                                                                                       |
| Investigational Product: | pembrolizumab                                                                                                                                                                                                                               |
| Phase:                   | PHASE1                                                                                                                                                                                                                                      |
| Medical Monitor:         | [Not Provided in Posting]                                                                                                                                                                                                                   |

## 2. Study Synopsis & Rationale

### 2.1 Study Objective

|                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|--------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Primary Objective: | <p>1. Dose-Limiting Toxicity (DLT): The occurrence of any of the following toxicities during Cycle 1 will be considered a DLT, if assessed by the investigator related to study intervention administration:</p> <ul style="list-style-type: none"><li>* Grade 4 nonhematologic toxicity</li><li>* Grade 4 hematologic toxicity lasting ≥7 days, except thrombocytopenia</li><li>* Any nonhematologic AE ≥Grade 3 in severity, with exceptions</li><li>* Any Grade 3 or Grade 4 nonhematologic laboratory value, as with pre-specified exceptions</li><li>* Any Grade 3 or Grade 4 laboratory abnormalities, with the exceptions</li><li>* Febrile neutropenia Grade 3 or Grade 4</li><li>* Prolonged delay (&gt;2 weeks) in initiating Cycle 2 due to intervention-related toxicity</li><li>* Any study drug toxicity that causes the participant to discontinue study drug during Cycle 1</li><li>* Missing &gt;25% of MK-4700 doses as a result of drug-related AEs during the first cycle</li><li>* Grade 5 toxicity</li></ul> <p>2. Percentage of Participants Who Experience an Adverse Event (AE): An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention.</p> <p>3. Percentage of Participants who Discontinue Study Treatment Due to an AE: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention.</p> |
|--------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

# Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

## Secondary Obj.:

1. Area Under the Concentration-Time Curve of MK-4700
2. Maximum Plasma Concentration (Cmax) of MK-4700
3. Minimum Plasma Concentration (Cmin) of MK-4700

## 2.2 Background & Rationale

The goal of this study is to learn about the safety of different doses of MK-4700 and if people tolerate them. The study will also measure what happens in a person's body over time when MK-4700 is given alone or with pembrolizumab (MK-3475) in order to find a dose that is safe, tolerated, and may work to treat certain types of cancer.

## 3. Study Design & Methodology

**Design Framework:** Type: INTERVENTIONAL, Phase: PHASE1, Status: ACTIVE\_NOT\_RECRUITING

**Target N:** 82

## 4. Subject Selection (Eligibility Criteria)

### 4.1 Inclusion Criteria

The key inclusion criteria include but are not limited to the following:

- \* Has histologically or cytologically confirmed advanced/metastatic solid tumor by pathology report who have experienced disease progression on or after prior anti-cancer treatments, or been intolerant to, or refused all treatment known to confer clinical benefit
- \* Has head and neck squamous cell carcinoma (HNSCC), melanoma (cutaneous), non-small cell lung cancer (NSCLC), cervical cancer, triple negative breast cancer (TNBC), urothelial carcinoma (UC), or renal cell carcinoma (RCC; clear cell, papillary)
- \* If human immunodeficiency virus (HIV) infected, must have well controlled HIV on antiretroviral therapy (ART)
- \* Has normal cardiac function

### 4.2 Exclusion Criteria

The key exclusion criteria include but are not limited to the following:

- \* If HIV-infected with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease
- \* Has known additional malignancy that is progressing or has required active treatment within the past 2 years
- \* Has known active central nervous system (CNS) metastases and/or carcinomatous meningitis
- \* Has active autoimmune disease that has required systemic treatment in the past 2 years except replacement therapy
- \* Has history of (noninfectious) pneumonitis/interstitial lung disease that required steroids
- \* Has current pneumonitis/interstitial lung disease
- \* Has active infection requiring systemic therapy
- \* Has known history of Hepatitis B (defined as Hepatitis B surface antigen reactive) or known active Hepatitis C virus infection
- \* Has had an allogeneic tissue/solid organ transplant in the last 5 years

## 11. Study Identifiers & Governance

# Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

**Primary ID:** 316411419878127298  
**Registry Number:** NCT06894771  
**Sponsor:** Merck  
**Study Title:** A Study of MK-4700 Alone or With Pembrolizumab in Participants With Advanced or Metastatic Solid Tumors (MK-4700-001)

## 15. Sign-off & Approval

Principal Investigator: \_\_\_\_\_ Date: \_\_\_\_\_  
Clinical Operations Lead: \_\_\_\_\_ Date: \_\_\_\_\_  
Lead Statistician: \_\_\_\_\_ Date: \_\_\_\_\_