

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1 Study of MK-5890 as Monotherapy and in Combination With Pembrolizumab in Participants With Advanced Solid Tumors
Short Title:	Study of Boserolimab (MK-5890) as Monotherapy and in Combination With Pembrolizumab (MK-3475) in Adults With Advanced Solid Tumors (MK-5890-001)
Protocol Number:	357436295497771169
NCT Number:	NCT03396445
Sponsor Name:	Merck
Investigational Product:	carboplatin
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants Who Experienced a Dose-Limiting Toxicity (DLT): DLTs were assessed using the National Cancer Institute Common Terminology Criteria for Adverse Events version 5. A DLT was defined as any of the following events: Grade 4 nonhematologic toxicity; Grade 4 hematologic toxicity lasting ≥ 7 days, except thrombocytopenia; Grade 4 thrombocytopenia; Grade 3 thrombocytopenia requiring a platelet transfusion; Nonhematologic adverse event (AE) Grade ≥ 3, with exceptions; Grade 3 or 4 nonhematologic laboratory abnormality; Prolonged delay (≥ 2 weeks) in initiating Cycle 2 due to treatment-related toxicity; Treatment-related toxicity resulting in study drug discontinuation during the DLT evaluation period; Missing $\geq 25\%$ of any study drug during the DLT evaluation period due to a treatment-related AE; or Grade 5 toxicity. The number of participants who experienced one or more DLTs is reported. 2. Number of Participants With One or More AEs: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study treatment, whether or not considered related to the study treatment. The number of participants who experienced one or more AEs is reported. 3. Number of Participants Who Discontinued Study Treatment Due to an AE: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study treatment, whether or not considered related to the study treatment. The number of participants who discontinued study treatment due to an AE is reported.
Secondary Obj.:	1. Area Under the Concentration-Time Curve From Time 0 to Last Quantifiable Concentration (AUC0-last) of Boserolimab 2. Area Under the Concentration-Time Curve From Time 0 to 3 Weeks (AUC0-3w) of Boserolimab 3. Area Under the Concentration-Time Curve From Time 0 to 6 Weeks (AUC0-6w) of Boserolimab

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

2.2 Background & Rationale

The purpose of this study is to assess the safety and pharmacokinetics of boserolimab (MK-5890) when administered alone and in combination with pembrolizumab (MK-3475) in adults. Boserolimab monotherapy or boserolimab plus pembrolizumab combination therapy will be administered in adults with advanced solid tumors, including endometrial cancer, for up to 35 administrations (approximately 2 years). The safety and pharmacokinetics of boserolimab when administered with pembrolizumab, pemetrexed and carboplatin in adults with non-squamous non-small cell lung cancer (NSCLC) and boserolimab when administered with pembrolizumab and nab-paclitaxel in adults with triple-negative breast cancer (TNBC) will also be assessed.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, Status: COMPLETED

Target N: 182

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Arms 1 & 2: Histologically or cytologically confirmed advanced/metastatic solid tumor by pathology report and has received or been intolerant to all treatment known to confer clinical benefit
- * Arm 3: Histologically or cytologically confirmed diagnosis of stage IV (M1a or M1b per current American Joint Committee on Cancer criteria) non-squamous NSCLC
- * Arm 4: Triple-negative breast cancer (TNBC) that is locally recurrent, inoperable, not previously treated with chemotherapy, and which cannot be treated with curative intent OR metastatic disease not previously treated with chemotherapy
- * Measurable disease by Response Evaluation Criteria in Solid Tumors Version 1.1 (RECIST 1.1) as assessed by the local site investigator/radiologist. Target lesions situated in a previously irradiated area are considered measurable if progression has been demonstrated in such lesions
- * Adequate organ function
- * Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1
- * Male participants must agree to use adequate contraception during the treatment period and for at least 120 days after the last dose of boserolimab or pembrolizumab OR 180 days after the last dose of chemotherapeutic agents and refrain from donating sperm during this period
- * Female participants must not be pregnant or breastfeeding and agree to follow use adequate contraception during the treatment period and for at least 120 days after the last dose of boserolimab or pembrolizumab OR 180 days after the last dose of chemotherapeutic agents
- * Submit an evaluable baseline tumor sample for analysis (either a newly obtained or archival tumor sample)

4.2 Exclusion Criteria

- * History of a second malignancy, unless potentially curative treatment has been completed with no evidence of malignancy for 2 years
- * Clinically active central nervous system metastases and/or carcinomatous meningitis
- * Has had a severe hypersensitivity reaction to treatment with a monoclonal antibody (mAb) and/or other components of

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

the study treatment

- * Active infection requiring systemic treatment
- * History of interstitial lung disease
- * History of (noninfectious) pneumonitis that required steroids or current pneumonitis
- * Symptomatic ascites or pleural effusion
- * Previously had a stem cell or bone marrow transplant
- * Previously had a solid organ transplant
- * Active autoimmune disease that has required systemic treatment in the past 2 years (i.e., with use of disease-modifying agents, corticosteroids, or immunosuppressive drugs) except vitiligo or resolved childhood asthma/atopy
- * Known human immunodeficiency virus (HIV) and/or active and acute Hepatitis B or C infections
- * Not fully recovered from any effects of major surgery without significant detectable infection
- * Pregnant or breastfeeding, or expecting to conceive or father children within the projected duration of the study
- * Had chemotherapy, definitive radiation, or biological cancer therapy within 4 weeks (2 weeks for palliative radiation) before the first dose of study treatment, or has not recovered to Grade ?1 or better from any AEs that were due to cancer therapeutics administered more than 4 weeks earlier
- * Expected to require any other form of antineoplastic therapy while participating in this study
- * On chronic systemic steroid therapy in excess of replacement doses (e.g., exceeding 10 mg/day of prednisone equivalent), or on any other form of immunosuppressive medication
- * Regular user (including "recreational use") of any illicit drugs at the time of signing informed consent, or has a recent history (within the last year) of substance abuse (including alcohol), as determined by the treating investigator. Participants who use cannabis for medicinal purposes or to treat specific symptoms will not be excluded unless it is being abused in the opinion of the treating investigator
- * Received a live-virus vaccine within 28 days before the first dose of study treatment
- * Currently participating and receiving study treatment in a study of an investigational agent or has participated and received study treatment in a study of an investigational agent or has used an investigational device within 28 days before the first dose of study treatment

Additional Exclusion Criteria for Participants in Arm 3:

- * Has received radiation therapy to the lung that is ≥ 30 Gray (Gy) within 6 months before the first dose of study treatment
- * Is unable to interrupt aspirin or other nonsteroidal anti-inflammatory drugs (NSAIDs), other than an aspirin dose ≥ 1.3 g per day, for a 5-day period (8-day period for long-acting agents, such as piroxicam).
- * Is unable or unwilling to take folic acid or vitamin B12 supplementation

Additional Exclusion Criteria for Participants in Arm 4:

- * Has a known history of hypersensitivity or allergy to nab-paclitaxel or any of its components
- * Has neuropathy $\geq$ Grade 2

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Has a history of class II-IV congestive heart failure or myocardial infarction within 6 months of randomization
- * Has received previous treatment with immune checkpoint inhibitor(s) (eg, Programmed Cell Death Receptor 1 (PD-1)/PD-L1)

11. Study Identifiers & Governance

Primary ID: 357436295497771169

Registry Number: NCT03396445

Sponsor: Merck

Study Title: Study of Boserolimab (MK-5890) as Monotherapy and in Combination With Pembrolizumab (MK-3475) in Adults With Advanced Solid Tumors (MK-5890-001)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____