

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase Ib, Multi-arm, Open-label, Study of HDM201 in Combination With MBG453 or Venetoclax in Adult Subjects With Acute Myeloid Leukemia (AML) or High-risk Myelodysplastic Syndrome (MDS)
Short Title:	HDM201 in Combination With MBG453 or Venetoclax in Patients With Acute Myeloid Leukemia (AML) or High-risk Myelodysplastic Syndrome (MDS)
Protocol Number:	734407662148635645
NCT Number:	NCT03940352
Sponsor Name:	Novartis
Investigational Product:	venetoclax
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Incidence of Adverse Events (AEs) and Serious Adverse Events (SAEs) as a measure of safety: Month 24 is assumed to be study end 2. Severity of Adverse Events (AEs) and Serious Adverse Events (SAEs) as a measure of safety: Month 24 is assumed to be study end 3. Incidence of dose limiting toxicities (DLTs) of treatment: end of first cycle
Secondary Obj.:	1. Overall Response Rate (ORR) 2. Best Overall Response (BOR) 3. Event Free Survival (EFS) for AML (Cheson 2003) or Progression Free Survival (PFS) for MDS (Cheson 2006)

2.2 Background & Rationale

This was a phase 1b, multi-arm, open-label study of HDM201 in combination with MBG453 or venetoclax in subjects with AML or high-risk MDS.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, Status: TERMINATED
Target N:	52

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

* Male or female patients ? 18 years of age at the date of ICF signature who present with one of the following:

1. Relapsed/refractory AML following ?1 prior therapies (but ?3 prior therapies) who have relapsed or exhibited

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

refractory disease (primary failure) and are deemed by the Investigator not to be candidates for standard therapy, including re-induction with cytarabine or other established chemotherapy regimens for patients with AML (patients who are suitable for standard re-induction chemotherapy or hematopoietic stem cell transplantation and willing to receive it are excluded)

2. First line AML patient unfit for standard induction chemotherapy (includes both de novo and secondary AML), except in countries where approved therapies are available. Patients who are suitable for hematopoietic stem cell transplantation and willing to receive it are excluded.

3. High-risk MDS patient (high and very high-risk groups according to rIPSS) who have failed hypomethylating agent therapy.

* ECOG performance status ? 1

* TP53wt tumor. At minimum exons 5, 6, 7 and 8 in the TP53 gene must be sequenced and determined to contain no mutations. The TP53 status must be obtained from a bone-marrow sample, collected no longer than 3 months before signing the main ICF.

* Patient must be a candidate for serial bone marrow aspirate and/or biopsy according to the institutional guidelines and be willing to undergo a bone marrow aspirate and/or biopsy at screening, during and at the end of therapy on this study. Exceptions may be considered after documented discussion with Novartis.

Main Exclusion Criteria:

Patients eligible for this study must not meet any of the following criteria:

* Prior combination treatment with compounds having the same mode of action:

* mdm2 or mdm4 inhibitors combined with TIM-3 inhibitors (for patients enrolled in treatment arm1)

* mdm2 or mdm4 inhibitors combined with Bcl-2 inhibitor (for patients enrolled in treatment arm2)

* History of severe hypersensitivity reactions to any ingredient of study drug(s) and other monoclonal antibodies (mAbs) and/or their excipients.

* Patients with acute promyelocytic leukemia with PML-RARA.

* Allogeneic stem cell transplant (HSCT) within last 6 months and/or active GvHD requiring systemic immunosuppressive therapy.

* GI disorders impacting absorption of oral HDM201 or venetoclax.

* Evidence of active bleeding or bleeding diathesis or major coagulopathy (including familial).

* Patients with active, known or suspected autoimmune disease (treatment arm 1 only).

Other eligibility criteria apply.

4.2 Exclusion Criteria

Patients eligible for this study must not meet any of the following criteria:

* Prior combination treatment with compounds having the same mode of action:

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * mdm2 or mdm4 inhibitors combined with TIM-3 inhibitors (for patients enrolled in treatment arm1)
- * mdm2 or mdm4 inhibitors combined with Bcl-2 inhibitor (for patients enrolled in treatment arm2)
- * History of severe hypersensitivity reactions to any ingredient of study drug(s) and other monoclonal antibodies (mAbs) and/or their excipients.
- * Patients with acute promyelocytic leukemia with PML-RARA.
- * Allogeneic stem cell transplant (HSCT) within last 6 months and/or active GvHD requiring systemic immunosuppressive therapy.
- * GI disorders impacting absorption of oral HDM201 or venetoclax.
- * Evidence of active bleeding or bleeding diathesis or major coagulopathy (including familial).
- * Patients with active, known or suspected autoimmune disease (treatment arm 1 only).

Other eligibility criteria apply.

11. Study Identifiers & Governance

Primary ID: 734407662148635645

Registry Number: NCT03940352

Sponsor: Novartis

Study Title: HDM201 in Combination With MBG453 or Venetoclax in Patients With Acute Myeloid Leukemia (AML) or High-risk Myelodysplastic Syndrome (MDS)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____