

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase I, Open-Label, Multiple-Dose Study of the Effect of Evobrutinib on the Pharmacokinetics of a Combined Oral Contraceptive in Healthy Female Participants
Short Title:	Effect of Evobrutinib on Pharmacokinetics of a Combined Oral Contraceptive
Protocol Number:	766317915096357473
NCT Number:	NCT07215806
Sponsor Name:	Merck
Investigational Product:	norethisterone
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Area Under the Plasma Concentration-time Curve From Time Zero Extrapolated to Infinity (AUC0-inf) of Ethinyl Estradiol and Norethisterone: AUC0-inf was calculated by combining AUC0-t and AUCextra. AUCextra represents an extrapolated value obtained by $C_{last\ pred}/\lambda_z$, where $C_{last\ pred}$ was the calculated serum concentration at the last sampling time point at which the measured serum concentration is at or above the Lower Limit of quantification (LLOQ) and λ_z was the apparent terminal rate constant determined by log-linear regression analysis of the measured serum concentrations of the terminal log-linear phase. 2. Maximum Observed Plasma Concentration (Cmax) of Ethinyl Estradiol and Norethisterone: Cmax was obtained directly from the concentration versus time curve.
Secondary Obj.:	1. Number of Participants With Treatment- Emergent Adverse Events (TEAEs) 2. Number of Participants With Treatment- Emergent Adverse Events (TEAEs) by Severity 3. Change From Baseline in Hematology Parameter: Hemoglobin at Day 18

2.2 Background & Rationale

The purpose of this study was to assess the effect of M2951 on the pharmacokinetics (PK) of a combined oral contraceptive [Ethinyl estradiol/Norethisterone (EE/NET)] in healthy female participants.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, Status: COMPLETED
Target N:	20

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Participants are overtly healthy as determined by medical evaluation, including no clinically significant abnormality identified on physical examination or laboratory evaluation and no active clinically significant disorder, condition, infection or disease that would pose a risk to participant safety or interfere with the study evaluation, procedures, or completion
- * Participants have a body weight within 50.0 and 100.0 kilograms (kg) (inclusive) and body mass index within the range 19.0 and 30.0 kilograms per square meter (kg/m²) (inclusive)
- * Participants are nonsmokers for at least 6 months preceding Screening
- * Female participants who are not a Woman of Childbearing Potential (WOCBP)
- * Other protocol defined inclusion criteria could apply

4.2 Exclusion Criteria

- * Participants with history or presence of clinically relevant respiratory, gastrointestinal, renal, hepatic, hematological, lymphatic, neurological, cardiovascular, psychiatric, musculoskeletal, genitourinary, immunological, dermatological, connective tissue diseases or disorders, as determined by medical evaluation
- * Participants with diagnosis of hemochromatosis, Wilson's disease, alpha 1 antitrypsin deficiency, or any other chronic liver disease including Gilbert's disease will be excluded from the study
- * Participants with prior history of splenectomy or any clinically relevant surgery within 3 months prior to Screening
- * Participants with history of any malignancy
- * Participants with history of chronic or recurrent acute infection or any bacterial, viral, parasitic, or fungal infections within 30 days prior to Screening and at any time between Screening and admission, or hospitalization due to infection within 6 months prior to Screening
- * Participants with history of shingles within 12 months prior to Screening
- * Administration of live vaccines or live-attenuated virus vaccines within 3 months prior to Screening. Administration of other types of vaccines [e.g., Severe acute respiratory syndrome coronavirus 2 (SARSCoV2) vaccines] is allowed until 4 weeks before admission to CRU, thereafter it is prohibited until the end of the study
- * Note: In case of clinical symptoms, the participant should be symptom-free for at least 1 week prior to admission to Clinical Research Unit (CRU)
- * Other protocol defined exclusion criteria could apply

11. Study Identifiers & Governance

Primary ID: 766317915096357473

Registry Number: NCT07215806

Sponsor: Merck

Study Title: Effect of Evobrutinib on Pharmacokinetics of a Combined Oral Contraceptive

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____