

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase IB, Open-Label, Multicenter, Single Arm Study Evaluating the Preliminary Efficacy, Safety, and Pharmacokinetics of Glofitamab in Combination With Rituximab Plus Ifosfamide, Carboplatin Etoposide Phosphate in Patients With Relapsed/Refractory Transplant or CAR-T Therapy Eligible Diffuse B-Cell Lymphoma
Short Title:	A Study Evaluating the Efficacy, Safety, and Pharmacokinetics of Glofitamab in Combination With Rituximab Plus Ifosfamide, Carboplatin Etoposide Phosphate in Participants With Relapsed/Refractory Transplant or CAR-T Therapy Eligible Diffuse B-Cell Lymphoma
Protocol Number:	793434791615708648
NCT Number:	NCT05364424
Sponsor Name:	Roche
Investigational Product:	etoposide
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Objective response rate (ORR), defined as the proportion of participants that achieves a CR or PR within three cycles of glofit-R-ICE, as determined by the investigator according to Lugano criteria
Secondary Obj.:	1. Event-free survival (EFS) after enrollment 2. Progression-free survival (PFS) after enrollment 3. Mobilization-adjusted response rate (MARR)

2.2 Background & Rationale

The purpose of this study is to evaluate the preliminary efficacy, safety, and pharmacokinetics of glofitamab (glofit) in combination with rituximab plus ifosfamide, carboplatin, and etoposide (R-ICE) in participants with relapsed or refractory (R/R) diffuse large B-cell lymphoma (DLBCL), who have failed one prior line of therapy incorporating an anti-cluster of differentiation (CD) 20 antibody (i.e., rituximab) and an anthracycline, and who are transplant or chimeric antigen receptor T-cell (CAR-T) therapy eligible, defined as being medically eligible for intensive platinum-based salvage therapy followed by autologous stem cell transplantation (ASCT) or for CAR-T therapy.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, Status: ACTIVE_NOT_RECRUITING
Target N:	43

4. Subject Selection (Eligibility Criteria)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4.1 Inclusion Criteria

- * Life expectancy $\geq$ 12 weeks
- * Histologically confirmed B-cell lymphoma
- * One line of prior systemic therapy including an anti-CD20 monoclonal antibody (i.e. rituximab) and an anthracycline
- * Relapsed or refractory disease after first-line chemoimmunotherapy
- * Eastern Cooperative Oncology Group (ECOG) Performance Status of 0 or 1
- * Participant must be a candidate for high-dose chemotherapy followed by ASCT or CAR-T therapy

4.2 Exclusion Criteria

- * Treatment with more than one prior line of therapy for DLBCL
- * Primary mediastinal B-cell lymphoma
- * Prior treatment with glofitamab or other bispecific antibodies targeting both CD20 and CD3
- * Peripheral neuropathy assessed to be Grade $\geq$ 1 according to National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE) v5.0 at enrollment
- * Treatment with radiotherapy, chemotherapy, immunotherapy, immunosuppressive therapy, or any investigational agent for the purposes of treating cancer within 2 weeks prior to first study treatment
- * Treatment with monoclonal antibodies for the purposes of treating cancer within 4 weeks prior to first study treatment
- * Primary or secondary CNS lymphoma at the time of enrollment or history of CNS lymphoma
- * Current or history of CNS disease, such as stroke, epilepsy, CNS vasculitis, or neurodegenerative disease
- * Known or suspected history of hemophagocytic lymphohistiocytosis (HLH)
- * Known history of progressive multifocal leukoencephalopathy
- * Adverse events from prior anti-cancer therapy that have not resolved to Grade 1 or better (with the exception of alopecia and anorexia, or as otherwise permitted by inclusion criteria)
- * Prior solid organ transplantation
- * Prior allogeneic stem cell transplant
- * Prior ASCT for lymphoma
- * Prior autologous stem cell transplant for any indication other than lymphoma, within 5 years from the start of study treatment
- * Active autoimmune disease requiring treatment
- * Prior treatment with systemic immunosuppressive medications (including, but not limited to, cyclophosphamide, azathioprine, methotrexate, thalidomide, and anti-tumor necrosis factor agents), within 4 weeks prior to first dose of study treatment
- * Ongoing corticosteroid use $\geq$ 30 mg/day of prednisone or equivalent. Participants who received corticosteroid treatment with $\geq$ 30 mg/day of prednisone or equivalent must be documented to be on a stable dose of at least 4 weeks' duration prior to Cycle 1 Day 1. Participants may have received a brief ($\leq$ 7 days) course of systemic steroids ($\leq$ 100 mg prednisone equivalent per day) prior to initiation of study therapy for control of lymphoma-related symptoms
- * Recent major surgery (within 4 weeks before the first study treatment) other than for diagnosis
- * Clinically significant history of cirrhotic liver disease

11. Study Identifiers & Governance

Primary ID: 793434791615708648

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Registry Number: NCT05364424

Sponsor: Roche

Study Title: A Study Evaluating the Efficacy, Safety, and Pharmacokinetics of Glofitamab in Combination With Rituximab Plus Ifosfamide, Carboplatin Etoposide Phosphate in Participants With Relapsed/Refractory Transplant or CAR-T Therapy Eligible Diffuse B-Cell Lymphoma

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____