

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A PHASE 1B, OPEN-LABEL STUDY OF ELRANATAMAB IN COMBINATION WITH IBERDOMIDE IN PARTICIPANTS WITH RELAPSED REFRACTORY MULTIPLE MYELOMA
Short Title:	A Study to Learn About the Effects of the Combination of Elranatamab (PF-06863135) and Iberdomide in Patients With Relapsed or Refractory Multiple Myeloma (MagnetisMM-30)
Protocol Number:	952664735435573374
NCT Number:	NCT06215118
Sponsor Name:	Pfizer
Investigational Product:	elranatamab
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Part 1: Number of participants with dose limiting toxicity (DLT): Dose limiting toxicity rate based on dose limiting toxicity evaluable participants 2. Part 2: Number of participants with Adverse Events (AE) by Seriousness and Relationship to Treatment: Number of participants with AE among participants who take at least 1 dose of study intervention. AEs are categorized by seriousness and relationship to treatment. Relatedness to study drug is assessed by investigator.
Secondary Obj.:	1. Part 1: Number of participants with Adverse Events (AE) by Seriousness and Relationship to Treatment 2. Part 1 and Part 2: Number of Participants with Adverse Events (AE) characterized by type, frequency, severity 3. Part 1 and Part 2: Number of Participants with Clinically Significant Change from Baseline in Laboratory Abnormalities

2.2 Background & Rationale

The main purpose of the study is to understand how safe and tolerable is elranatamab when given along with iberdomide.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, Status: RECRUITING
Target N:	87

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Prior diagnosis of multiple myeloma as defined by IMWG criteria
- * Measurable disease based on IMWG criteria as defined by at least 1 of the following:
 - * Serum M-protein ≥ 0.5 g/dL by SPEP
 - * Urinary M-protein excretion ≥ 200 mg/24 hour by UPEP
 - * Serum immunoglobulin FLC ≥ 10 mg/dL (≥ 100 mg/L) AND abnormal serum immunoglobulin kappa to lambda FL ratio (< 0.26 or > 1.65)
- * Part 1: Received 2-4 prior lines of therapy for multiple myeloma, consisting of at least 1 immunomodulatory drug and 1 proteasome inhibitor.
- * Part 2: Received 1-3 prior lines of therapy for multiple myeloma, consisting of at least 1 immunomodulatory drug and 1 proteasome inhibitor.
- * ECOG performance status 0-1
- * Resolved acute effects of any prior therapy to baseline severity or CTCAE Grade ≤ 1

4.2 Exclusion Criteria

- * Plasma cell leukemia, Smoldering multiple myeloma, Waldenström's macroglobulinemia, Amyloidosis, POEMS Syndrome
- * Impaired cardiovascular function or clinically significant cardiovascular diseases
- * Stem cell transplant within 12 weeks prior to enrollment or active graft vs host disease
- * Participants with any active, uncontrolled bacterial, fungal, or viral infection
- * Any other active malignancy within 3 years prior to enrollment, except for adequately treated basal cell or squamous cell skin cancer, or carcinoma in situ
- * Previous treatment with:
 - * BCMA-directed or CD3 redirecting therapy
 - * Iberdomide (CC-220) or Mezigdomide
- * Administration of strong inhibitor or inducer of CYP3A4/5 within 2 weeks prior to dosing and during the study
- * Administration with an investigational product within 30 days preceding the first dose of study intervention
- * Participant is unable or unwilling to undergo protocol required thromboembolism prophylaxis

11. Study Identifiers & Governance

Primary ID: 952664735435573374

Registry Number: NCT06215118

Sponsor: Pfizer

Study Title: A Study to Learn About the Effects of the Combination of Elranatamab (PF-06863135) and Iberdomide in Patients With Relapsed or Refractory Multiple Myeloma (MagnetisMM-30)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____