

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III, Multicenter, Randomized, Double-Blind, Placebo-Controlled Trial Comparing the Efficacy and Safety of Polatuzumab Vedotin in Combination With Rituximab and CHP (R-CHP) Versus Rituximab and CHOP (R-CHOP) in Previously Untreated Patients With Diffuse Large B-Cell Lymphoma
Short Title:	A Study Comparing the Efficacy and Safety of Polatuzumab Vedotin With Rituximab-Cyclophosphamide, Doxorubicin, and Prednisone (R-CHP) Versus Rituximab-Cyclophosphamide, Doxorubicin, Vincristine, and Prednisone (R-CHOP) in Participants With Diffuse Large B-Cell Lymphoma
Protocol Number:	187939002460679087
NCT Number:	NCT03274492
Sponsor Name:	Roche
Investigational Product:	cyclophosphamide
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression-Free Survival (PFS) as Assessed by the Investigator, Using the Lugano Response Criteria for Malignant Lymphoma
Secondary Obj.:	1. Percentage of Participants With Complete Response (CR) as Assessed by Fluorodeoxyglucose Positron Emission Tomography (FDG-PET) by Blinded Independent Central Review (BICR) 2. Event-Free Survival-Efficacy (EFSeff) as Assessed by the Investigator, Using the Lugano Response Criteria for Malignant Lymphoma 3. Percentage of Participants Who are Progression Free as Assessed by the Investigator, Using the Lugano Response Criteria for Malignant Lymphoma

2.2 Background & Rationale

This Phase III, randomized, double-blind, placebo-controlled study will compare the efficacy, safety, and pharmacokinetics of polatuzumab vedotin plus R-CHP versus R-CHOP in participants with previously untreated diffuse large B-cell lymphoma (DLBCL).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	1000

4. Subject Selection (Eligibility Criteria)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4.1 Inclusion Criteria

- * Previously untreated participants with cluster of differentiation 20 (CD20)-positive DLBCL, including one of the following diagnoses by 2016 World Health Organization (WHO) classification of lymphoid neoplasms: DLBCL, not otherwise specified (NOS) including germinal center B-cell type, activated B-cell type; T-cell/histiocyte-rich large B-cell lymphoma; Epstein-Barr virus-positive DLBCL, NOS; anaplastic lymphoma kinase (ALK)-positive large B-cell lymphoma; human herpesvirus-8 (HHV8)-positive DLBCL, NOS; High-grade B-cell lymphoma with MYC and B-cell lymphoma 2 (BCL2) and/or B-cell lymphoma 6 (BCL6) rearrangements (double-hit or triple-hit lymphoma); High-grade B-cell lymphoma, NOS
- * Availability of archival or freshly collected tumor tissue before study enrolment
- * International Prognostic Index (IPI) score of 2-5
- * Eastern Cooperative Oncology Group (ECOG) Performance Status of 0, 1, or 2
- * Life expectancy greater than or equal to ($\geq$) 12 months
- * Left ventricular ejection fraction (LVEF) $\geq$ 50 percent (%) on cardiac multiple-gated acquisition (MUGA) scan or cardiac echocardiogram (ECHO)
- * Adequate hematologic function
- * Female participants: Agreement to remain abstinent (refrain from heterosexual intercourse) or use contraceptive methods and refrain from donating eggs.
- * Male participants: agreement to remain abstinent (refrain from heterosexual intercourse) or use a condom and agreement to refrain from donating sperm.

4.2 Exclusion Criteria

- * History of severe allergic or anaphylactic reactions to humanized or murine monoclonal antibodies or known sensitivity or allergy to murine products
- * Contraindication to any of the individual components of CHOP, including prior receipt of anthracyclines
- * Prior organ transplantation
- * Current Grade greater than ($>$) 1 peripheral neuropathy by clinical examination
- * Demyelinating form of Charcot-Marie-Tooth disease
- * History of indolent lymphoma
- * History of follicular lymphoma grade 3B
- * B-cell lymphoma, unclassifiable, with features intermediate between DLBCL and classical Hodgkin lymphoma (grey-zone lymphoma)
- * Primary mediastinal (thymic) large B-cell lymphoma
- * Burkitt lymphoma
- * Prior treatment with cytotoxic drugs within 5 years of screening for any condition (example [e.g.], cancer, rheumatoid arthritis) or prior use of any anti-CD20 antibody
- * Prior use of any monoclonal antibody within 3 months of the start of Cycle 1
- * Prior therapy for DLBCL, with the exception of nodal biopsy
- * Corticosteroid use $\geq$ 30 mg/day of prednisone or equivalent, for purposes other than lymphoma symptom control
- * Participants with central nervous system (CNS) lymphoma (primary or secondary involvement), primary effusion DLBCL, and primary cutaneous DLBCL
- * Vaccination with live vaccines within 28 days prior to the start of Cycle 1

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Any investigational therapy within 28 days prior to the start of Cycle 1
- * History of other malignancy that could affect compliance with the protocol or interpretation of results
- * Evidence of significant, uncontrolled, concomitant diseases that could affect compliance with the protocol or interpretation of results, including significant cardiovascular disease or pulmonary disease
- * Recent major surgery (within 4 weeks prior to the start of Cycle 1), other than for diagnosis
- * History or presence of an abnormal electrocardiogram (ECG) that is clinically significant in the investigator's opinion, including complete left bundle branch block, second- or third-degree heart block, or evidence of prior myocardial infarction
- * Known active bacterial, viral, fungal, mycobacterial, parasitic, or other infection (excluding fungal infections of nail beds) at study enrollment or significant infections within 2 weeks before the start of Cycle 1
- * Clinically significant liver disease, including active viral or other hepatitis, current alcohol abuse, or cirrhosis
- * Prior radiotherapy to the mediastinal/pericardial region
- * Participants with suspected active or latent tuberculosis
- * Positive test results for chronic hepatitis B and hepatitis C infection
- * Known history of human immunodeficiency virus (HIV) seropositive status
- * Positive results for the human T-lymphotrophic 1 virus (HTLV-1)
- * Participants with a history of progressive multifocal leukoencephalopathy

11. Study Identifiers & Governance

Primary ID: 187939002460679087

Registry Number: NCT03274492

Sponsor: Roche

Study Title: A Study Comparing the Efficacy and Safety of Polatuzumab Vedotin With Rituximab-Cyclophosphamide, Doxorubicin, and Prednisone (R-CHP) Versus Rituximab-Cyclophosphamide, Doxorubicin, Vincristine, and Prednisone (R-CHOP) in Participants With Diffuse Large B-Cell Lymphoma

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____