

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Multicenter Study of Secukinumab, With a Randomized Double-blind, Placebo-controlled Withdrawal-retreatment Period, to Evaluate Maintenance of Response in Participants With Non-radiographic Axial Spondyloarthritis Who Achieved Remission
Short Title:	A Study of Secukinumab to Evaluate Maintenance of Response in Participants With Non-radiographic Axial Spondyloarthritis Who Achieved Remission
Protocol Number:	309611177861007345
NCT Number:	NCT05622708
Sponsor Name:	Novartis
Investigational Product:	secukinumab
Phase:	PHASE4
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective: 1. The proportion of participants remaining flare-free during Treatment Period 2: The primary efficacy endpoint is the proportion of participants in the randomized withdrawal population remaining flare-free at Week 120. A flare is defined as ASDAS-CRP $\geq$ 2.1 at 2 consecutive visits, or ASDAS-CRP $\geq$ 3.5 at any visit during Treatment Period 2, starting at Week 60.

Parameters used for ASDAS-CRP include:

- * Spinal pain (BASDAI question 2),
- * Patient's global assessment of disease activity,
- * Peripheral pain/swelling (BASDAI question 3),
- * Duration of morning stiffness (BASDAI question 6)
- * C-reactive protein (CRP) in mg/L

Secondary Obj.: 1. Time to flare during Treatment Period 2 2. Number of participants with Adverse Events

2.2 Background & Rationale

This study will establish whether prolonged chronic dosing with secukinumab is needed in participants with Non-radiographic axial spondyloarthritis, (nr-axSpA) who have achieved remission. Remission is defined as Ankylosing Spondylitis Disease Activity Score - C-reactive protein (ASDAS-CRP) Inactive Disease (ID) response (ASDAS-CRP $\leq$ 1.3). Maintenance of remission on continued secukinumab treatment will be evaluated compared to placebo using a randomized withdrawal design. The primary outcome measure for this study is the proportion of participants remaining

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

flare-free at Week 120.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE4, Status: RECRUITING

Target N: 340

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Male or non-pregnant, non-lactating female participants at least 18 years of age
- * Clinical diagnosis of axSpA AND according to ASAS axSpA criteria:
 1. Inflammatory back pain for at least 6 months
 2. Onset before 45 years of age
 3. Sacroiliitis on MRI (magnetic resonance imaging) (as assessed by central reader) with ? 1 SpA feature OR HLA-B-27 positive with ?2 SpA features
- * Objective signs of inflammation at screening, evident by either MRI with Sacroiliac Joint inflammation (as assessed by central reader) AND / OR hsCRP \> ULN (as defined by the central lab)
- * Active axSpA as assessed by total BASDAI ? 4 cm (0-10 cm) at baseline.
- * Spinal pain as measured by BASDAI question #2 ? 4 cm (0-10 cm) at baseline.
- * Total back pain as measured by VAS (visual analog scale) ? 40 mm (0-100 mm) at baseline.
- * Participants should have been on at least 2 different NSAIDs (non-steroidal anti-inflammatory drugs) at the highest recommended dose for at least 4 weeks in total prior to baseline with an inadequate response or failure to respond, or less if therapy had to be withdrawn due to intolerance, toxicity or contraindications.

4.2 Exclusion Criteria

- * Participants with radiographic evidence for sacroiliitis, grade ? 2 bilaterally or grade ? 3 unilaterally (radiological criterion according to the modified New York diagnostic criteria for AS) as assessed by central reader.
- * Participants taking high potency opioid analgesics (e.g., methadone, hydromorphone, morphine).
- * Previous exposure to secukinumab or any other biologic drug directly targeting IL-17 or IL-17 receptor or previous treatment with immunomodulatory biologic agents including those targeting TNF? (tumor necrosis factor ?) (unless participants discontinued the treatment with TNF? inhibitor due to a reason other than efficacy \[primary or secondary lack of efficacy, inadequate response\] and only after appropriate wash-out period prior to baseline was observed).
- * History of hypersensitivity to the study drug or its excipients or to drugs of similar chemical classes.
- * Active ongoing inflammatory diseases other than nr-axSpA that might confound the evaluation of the benefit of secukinumab therapy, including uveitis.
- * Active inflammatory bowel disease.
- * History of ongoing, chronic or recurrent infectious disease or evidence of tuberculosis infection.

11. Study Identifiers & Governance

Primary ID: 309611177861007345

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Registry Number: NCT05622708

Sponsor: Novartis

Study Title: A Study of Secukinumab to Evaluate Maintenance of Response in Participants With Non-radiographic Axial Spondyloarthritis Who Achieved Remission

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____